

ACCEPTED
 by Self Cert. Plan
 SEP 9 1987
 EPA Reg. No. 239-579

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
DANGER
 Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Avoid inhalation of dust. Remove and separately launder clothing before reuse.
STATEMENT OF PRACTICAL TREATMENT
 If swallowed contact physician, poison control center, or emergency clinic before inducing vomiting. Give water or milk to drink and induce vomiting by touching back of throat with finger, or by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person. If medical advice cannot be obtained, take person and a copy of the label or technical bulletin to the nearest emergency treatment facility. In case of eye contact, immediately flush eyes with plenty of clear water and get medical attention. For skin contact, wash with plenty of soap and water. See a doctor if irritation occurs or rash develops. If inhaled, remove person from exposure area.
 Note to Physicians: Emergency Information—call (415) 233-3737.
ENVIRONMENTAL HAZARDS
 This product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

STORAGE AND DISPOSAL
PROHIBITIONS
 Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.
STORAGE
 Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Protect from excessive heat. For help with any spill, leak, fire or exposure involving this material, call day or night (415) 233-3737.
PESTICIDE DISPOSAL
 This product is acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.
CONTAINER DISPOSAL
 Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Do not reuse container. Dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron. Fieldworkers and harvesters must wear chemical resistant gloves if entering treated areas within 4 days following application. Leather or fabric gloves are not acceptable. A dust mask must be worn when mixing and loading this product. AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

ORTHOCIDE 75 Seed Protectant is a special formulation containing ORTHOCIDE (Captan) for seed treatment application. ORTHOCIDE 75 Seed Protectant now contains a binder which helps to adhere the treaters to the seed so that there is minimum loss during processing and planting. Has no adverse effect on seed germination or planting rate. It is used in preventing seed rot and damping-off diseases on germinating seeds and seedlings and smut on onion and hulled sorghum seeds.

USE PRECAUTION
 When using formulations that do not contain dye, to comply with 21 CFR Chapter 1, Section 2.5, all seed treated with a pesticide must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.

SLURRY TREATMENT: When used in slurry treaters, do not allow mixtures to stand overnight. Clean out equipment thoroughly between treating operations.

GLADIOLUS CORMS: For Prevention of Bulb Rot—Use 1 1/2 oz. in 1 gal. water (1 lb. to 10 gals.). Soak bulbs in mixture 20 to 30 minutes.


GRASS SEED (Range Grasses such as Wheatgrass and Bromegrass): To control *Podosporiella verticillata*: 12 oz. ORTHOCIDE 75 Seed Protectant per 100 lbs. seed. Apply as slurry treatment.

ONION SEED PELLETTING: Use 1 lb. ORTHOCIDE 75 Seed Protectant per 1 lb. of seed plus 1/2 pt. methyl cellulose 5% solution.



ORTHO
OR
75 S

(FUNGICIDE)
 A Slurry Product
 Prevents Seed F



KEEP
 SEE SIDE PANEL FO

NET WEIGHT

BEST AVAILABLE COPY

3. L
 3. LC

 1. COI
 If neces
 NAME
 J.
 TITLE
 SU
 2. SIGN.
 4. TYPE
 J.
 EPA Form

Handwritten notes and numbers: 579, TS, S



ORTHO

ORTHO[®]CIDE

75 Seed Protectant

(FUNGICIDE)
A Slurry Product — CONTAINS BINDER
Prevents Seed Rot and Damp-off Diseases



Active Ingredients	By Wt.
*Captan	73.3%
Related Derivatives	17%
Inert Ingredients	25.0%

*N-Trichloromethyl thio-4-cyctahexene-1,2-dicarbaximide.

**KEEP OUT OF REACH OF CHILDREN
DANGER**
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NET WEIGHT _____

BEST AVAILABLE COPY

Directions. Do not get in eyes. Wear mask and goggles. Remove and separate...

...giving five water or milk to drink... Do not induce vomiting or give... and a copy of the label or technical literature... flush eyes with plenty of clear water... a doctor if irritation occurs or rash...

...of wastes. In the event that treated...

...AND DIRECTIONS, AND WITH AP...

...3-3737

...is a violation of Federal Law... State Pesticide or Environmental... for guidance

...Empty residue into application... by State and local authorities

...long pants or a coverall that covers... or a face shield and a chemically... within 4 days following application.

...contaminated clothing. Wash protective... must be laundered separately... with captan must be destroyed

...for seed treatment application... the seed so that there is minimum... It is used in preventing seed rot... and other fungal diseases.

...U.S. all seed treated with a pesticide... lead for animals.

...Clean out equipment thoroughly

...Soak bulbs in mixture 20 to 30

...vaccillata: 12 oz. ORTHOCIDE

...methyl cellulose 5% solution

CONTAINS DYE

TYPE

THE FOLLOWING TABLE gives amount of ORTHOCIDE 75 Seed Protectant to use on various seeds. Do not use on baby lima beans.

SEED	Dosage Rate	Lbs. Seeds Treated/ lb. of Product
CORN — Field	1 1/2 oz./100 lbs. 1/4 oz./bu	1280
CORN — Sweet	2 1/2 oz./100 lbs. 1 1/2 oz./bu	600
WHEAT	1 1/2 oz./100 lbs. 1/4 oz./bu	1280
OATS	2 oz./100 lbs. 1/2 oz./bu	800
BARLEY	1 1/2 oz./100 lbs. 1/4 oz./bu	1070
FLAX	2 1/2 oz./100 lbs. 1 1/2 oz./bu	640
BEANS (Except Baby Lima)	1 1/2 oz./100 lbs. 1 oz./bu	910
PEAS	1 1/2 oz./100 lbs.	910
LENTILS (Washington & Idaho Only)	1 1/2 oz./100 lbs.	910
CARROT	6 oz./100 lbs.	265
CUCUMBERS, CANTALOUPE	2 oz./100 lbs.	800
SQUASH, PUMPKIN, WATERMELON	1 1/2 oz./100 lbs.	1200
BEETS, SWISS CHARD	8 oz./100 lbs.	200
RUTABAGA	3 oz./100 lbs.	530
SUGAR BEETS	8 oz./100 lbs.	200
CLOVER	5 1/2 oz./100 lbs.	290
ALFALFA	3 1/2 oz./bu	290
SOYBEANS	1 1/2 oz./100 lbs. 1 oz./bu	950
TURNIP	2 oz./100 lbs.	800
COTTON — Acid Delinted	2 oz./100 lbs.	800
COTTON — Machine Delinted Fuzzy (not delinted)	3 oz./100 lbs.	530
BLUE GRASS	5 1/2 oz./100 lbs.	290
CABBAGE, CAULIFLOWER, RADISH, BROCCOLI, BRUSSELS SPROUTS	1 oz./100 lbs.	1600
SPINACH, EGGPLANT	4 oz./100 lbs.	400
SORGHUM hulled (Kernel Small)	3 oz./100 lbs.	530
PEPPERS	2 oz./100 lbs.	800
COWPEAS	2 oz./100 lbs. 1 1/2 oz./bu	800
RYE	1 1/2 oz./100 lbs.	910
SESAME	1 oz./100 lbs.	1600

Do not feed treated seed to animals. Do not use treated seed for human consumption. Seeds treated with ORTHOCIDE 75 Seed Protectant, sold or shipped as feed or foodstuffs may be considered adulterated under state and federal laws. Treated seed must be labeled as follows: "TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL"

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

**RQ Hazardous substance, solid,
N.O.S., NA 9188, (Captan)**

ORM-E

Chevron Chemical Company © 1984

Ortho Agricultural Chemicals Division/San Francisco CA 94120-7144
Form 9699-H Product 2006 Made in U.S.A. EPA Reg. No. 239-579-AA EPA Est. 239 MO-1