ORTHOCIDE 7.5 Dust

Chevron ORTHO

ACCEPTED
239-604

(FUNGICIDE)

Active Ingredient	By Wt.
*Captan	7.5%
Inert Ingredients	92.5%

*N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

STORE IN COOL, DRY PLACE
PROTECT FROM EXCESSIVE HEAT

This product is formulated for application by ground or air equipment.

ALMONDS: Brown Rot, Shot Hole, Scab, Leaf Blight—Use 45 lbs. per acre. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. May be used up to 12 days of harvest if hulls are not fed to dairy animals or animals being finished for slaughter.

APPLES: Apple Scab, Brooks Fruit Spot, Flyspecking, Bitter Rot, Black Rot, Botryosphaeria, Black Pox—25 to 40 lbs. per acre. Make first application at delayed dormant period. Repeat applications for each infection period or at weekly intervals as long as necessary.

APRICOTS: Brown Rot, Jacket Rot—50 to 60 lbs. per acre. Apply at red bud stage and repeat at 75 per cent petal fall.

BEANS AND PEAS: Damp-Off on Beans, Root Rot on Beans and Peas—50 to 60 lbs. per acre. Work into top 3 or 4 inches of soil before planting.

CANTALOUPES, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELON, PUMPKINS: Downy Mildew, Angular Leaf Spot, Anthracnose—Apply 25 to 30 lbs. per acre on young plants, increasing to 40 lbs. per acre on mature plants. Apply at first sign of disease and continue at weekly intervals as necessary.

NET WEIGHT 50 POUNDS

Chevron Chemical Company
Ortho Division/San Francisco, Calif. 94119
Richmond, California Fresno, California
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 2976 Q2 Product 1558 Made in U.S.A.
EPA Reg. No. 239-604-AA

CARROTS: Septoria Leafspot, Early Blight (Cercospora Blight)—25 lbs. per acre. Apply at first sign of disease. Repeat at 6 to 10-day intervals.

CELERY: Pink Rot, Damp-Off Diseases (Field)—26 lbs. per acre. **(Seed Bed)**—40 to 45 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

CHERRIES: Cherry Leaf Spot, Brown Rot—50 to 60 lbs. per acre. Apply at popcorn, blossom, and petal fall stages, continue at 10 to 14-day intervals.

GRAPES: Mold Control on Fruit on Drying Trays in Field—Penicillium spp., Aspergillus spp., Botrytis Cinerea, Rhizopus spp.—13 lbs. per acre. Apply to fruit on trays for protection of fruit and prevention of above molds. **Apply to give even coverage.** During damp weather make a second application after trays are turned. Apply by hand, ground duster or airplane. If applied by plane increase to 20 lbs. per acre.

MANGO: Cercospora Spot or Blotch—50 to 60 lbs. per acre. Begin applications at first sign of disease. Repeat at regular intervals as necessary to maintain control.

ORNAMENTALS: ROSES—Black Spot. CARNATIONS—Rust, Leaf Spot. CHRYSANTHEMUMS—Flower Blight—Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10-day intervals as necessary.

PEACH DISEASES: Brown Rot, Peach Scab—50 to 60 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

PEARS: Pear Scab (except Pacific Northwest)—25 to 40 lbs. per acre. Make first application during early finger and petal fall stages. Repeat applications for each infection period or at weekly intervals as long as necessary. Russeting may be produced on Bosc Pears. Do not use on D'Anjou Pears.

CAUTION

READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Avoid inhalation of dust. Avoid contact with skin. Do not store or transport near feed or food. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

[illegible] equipment or [illegible]

POTATOES (Seed Pieces): Fusarium Seed Rot, Verticillium Wilt and Damp-Off—Dust at 10 ounces per bushel cut seed pieces (16 ozs. or 1 lb. to 100 lbs. cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 to 3 days in open crates before planting. Do not use treated seed for feed or food purposes.

POTATOES: Early and Late Blight—Young Plants—30 to 35 lbs. per acre. **Mature Plants**—40 lbs. per acre. Begin applications when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals as long as disease threatens.

RASPBERRIES: Fruit Rot 35 lbs. per acre making applications 3 to 5 days before harvest starts, at mid-harvest and 8 to 10 days after mid-harvest.

SOIL AND GREENHOUSE BENCH TREATMENT: Preplanting for Damp-Off and Root Rot Diseases on Seedling or Transplants of Roses (and other shrubs and trees), Many flowers and Vegetables (such as Beans, Peas, Cucurbits) and Lawn Seedbeds—2½ lbs. per 1000 sq. ft. (75 to 100 pounds per acre). Work into upper 3 or 4 inches of soil before planting.

STRAWBERRIES: Botrytis Rot—40 lbs. per acre. Make first application when new growth starts in spring before fruit starts to form. Repeat applications weekly. **Severe Infection Conditions**—Continue through harvest period treating immediately after each picking.

TOMATOES: Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot—Young Plants—25 to 35 lbs. per acre. **Mature Plants**—Increase to 40 to 50 lbs. per acre. Apply at first sign of disease. Repeat at 7 to 10-day intervals.

TURF AND LAWN: Brown Patch, Leaf Spot, Damp-Off, Root Rot—2 to 3 lbs. per 1000 sq. ft. Begin treatment before infection occurs. Repeat at 7 to 10-day intervals throughout summer.

COMPATIBILITY: Do [illegible] spray.

CONDITIONS OF SALE: [illegible] MERCHANTABILITY, [illegible] connection with the use of [illegible] caused by fault to other plants or crops [illegible] directions are followed. Buyer and user agree to assume all risks and liability resulting therefrom and hereby [illegible] material.