

PRECAUTIONARY STATEMENTS - 12 pt

HAZARDS TO HUMANS & DOMESTIC ANIMALS - 12 pt

CAUTION

Causes eye irritation. May cause allergic skin reaction. Avoid contact with eyes, skin or clothing. Avoid inhalation of dust. Wash thoroughly after using.

STATEMENT OF PRACTICAL TREATMENT - 12 pt

In case of eye contact, wash eyes immediately with fresh water for at least 15 minutes. See a doctor if irritation persists. Note to Physicians: Emergency Information - call (415) 233-3737.

ENVIRONMENTAL HAZARDS - 12 pt

This product is toxic to fish. Do not apply directly to water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

REENTRY STATEMENT

Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Do not enter treated areas without protective clothing until sprays have dried.

Certain states may require more restrictive reentry intervals; consult your State Department of Agriculture for further information.

Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas about to be treated.

STORAGE AND DISPOSAL - 12 pt

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Do not store or transport near feed or food.

For help with any spill, leak, fire or exposure involving this material, call day or night (415) 233-3737.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Do not reuse container. Dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS

This product is formulated for application by ground or air equipment.

ALMONDS: Brown Rot, Shot Hole, Scab, Leaf Blight—Use 50 lbs per acre. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. May be used up to 12 days of harvest if hulls are not fed to dairy animals or animals being finished for slaughter.

APPLES: Apple Scab, Brooks Fruit Spot, Flyspecking, Bitter Rot, Black Rot, Botryospheria, Black Rot—30 to 50 lbs. per acre. Make first application at delayed dormant period. Repeat applications for each infection period or at weekly intervals as long as necessary.



ORTHO

ORTHO-CIDE® 5
Dust

CONFIDENTIAL

(FUNGICIDE)

PROPERTY OF

CHEVRON CHEMICAL COMPANY

Active Ingredients

| | |
|---------------------------|-------|
| *Captan | 4.7% |
| Related Derivatives | 0.3% |
| Inert Ingredients | 95.0% |

*N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide.

UNTIL ACCEPTED BY EPA

KEEP OUT OF REACH OF CHILDREN - 12 pt

CAUTION - 18 pt

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY LABELING.

ACCEPTED

AUG 29 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this product is registered under EPA Reg. No. 239-619

NET WEIGHT 50 POUNDS

Chevron Chemical Company
Ortho Agricultural Chemicals Division
San Francisco CA 94119 Richmond CA 94804
Product 1605 Made in U.S.A.
Form 3867-N2 EPA Est. 239-NJ-1
EPA Reg. No. 239-619-AA

BEANS AND PEAS: Damp-Off on Beans, Root Rot on Beans and Peas—70 to 80 lbs. per acre. Work into top 3 or 4 inches of soil before planting.

CANTALOUPE, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELON, PUMPKINS: Downy Mildew, Angular Leaf Spot, Anthracnose—Apply 25 to 30 lbs. per acre on young plants, increasing to 40 to 50 lbs. per acre on mature plants. Apply at first sign of disease and continue at weekly intervals as necessary.

CELERY: Pink Rot, Damp-Off Diseases—Field—40 lbs. per acre. (Seed Bed)—Use 40 to 50 pounds per acre. Apply when all seeds have germinated. Repeat as necessary.

GRAPES, RAISINS: Mold Control on Fruit on Drying Trays in Field—Penicillium spp., Aspergillus spp., Botrytis Cinerea, Rhizopus spp.—20 lbs. per acre. Apply to trays for protection of fruit and prevention of above molds. Apply to give even coverage. Repeat as necessary. Apply by hand, ground duster or airplane. If applied by plane increase to 30 lbs. per acre.

LETTUCE: Downy Mildew (California and Arizona)—40 to 50 lbs. per acre. Apply at first sign of disease and continue at weekly intervals as necessary.

ONIONS: Downy Mildew—30 lbs. per acre. Apply at first sign of disease and continue at weekly intervals as necessary.

ORNAMENTALS: ROSES—Black Spot, CARNATIONS—Rust and Leaf Spot, CHRYSANTHEMUMS—Flower Blight—Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10-day intervals as necessary.

PEARS: Pear Scab (except Pacific Northwest)—30 to 50 lbs. per acre. Make first application during early linger and petal fall stages. Repeat applications for each infection period or at weekly intervals as long as necessary. Rusting may be produced on Box Pears. Do not use on D'Anjou Pears.

POTATOES (Seed Pieces): Damp-Off—Dust of 15 ounces per bushel cut seed pieces (24 ozs. or 1 1/2 lbs. to 100 lbs. cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 8 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 days in open crates before bagging.

RASPBERRIES: Fruit Rot—45 lbs. per acre making applications 3 to 5 days before harvest starts, at mid harvest and 8 to 10 days after mid harvest.

SOIL AND GREENHOUSE BENCH TREATMENT: Preplanting for Damp-Off and Root Rot Diseases on Seedlings or Transplants of Roses (and other shrubs and trees), Many Flowers and Vegetables (such as Beans, Peas, Cucurbits) and Lawn Seedbeds—2 1/2 to 4 pounds per 1000 sq. ft. (100 to 120 lbs. per acre). Work into upper 3 or 4 inches of soil before planting.

STRAWBERRIES: Botrytis Rot—40 to 50 pounds per acre. Make first application when new growth starts in spring before fruit starts to form. Severe Infection Conditions—Continue applications through harvest period, weekly, immediately after each picking.

TOMATOES: Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot—On young plants apply 25 to 35 pounds per acre, increasing to 40 to 50 pounds per acre on mature plants. Begin applications at first sign of disease and continue at weekly intervals as necessary.

COMPATIBILITY: Do not use this product on foliage closely following oil sprays.

Do not apply under conditions involving possible drift to food, forage, or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

1 of 1
Page 21
239-619