


OPTHO ISOTOX° Seed Treater (F)



READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

ISOTOX Seed Treater (F) is a formulation for use in the treatment of seeds for protection against wireworms, seed corn maggots and other soil-borne insects, as well as added fungicidal protection at planting time. Also repellent to pheasants. Where Seedling Blight, Damp-Off or Seed Decay normally occurs, combine with a full dosage of protective fungicide. ISOTOX Seed Treater (F) may be used on corn seed previously treated with a protective fungicide.

DIRECTIONS

APPLY IN PLANTER BOX USING DESIGNATED AMOUNT PER 100 LBS. SEED. MEASURING CUP ENCLOSED—One level cup contains approximately one ounce of product.

CORN -8 ozs.

BEANS: (Use with a protective fungicide)

Baby Bush Limas - 2 ozs.

Bush, Snap and all other except Limas — 2 ozs.

Standard and Fordhook Large Limas —1 oz.

SUGAR BEETS-16 ozs. (1 lb.)

PEAS - 4 025.

CUCUMBERS, CANTALOUPES, WATERMELONS, SQUASH, PUMPKIN-4 ozs.

CARROTS, ONIONS – 4 ozs.

COTTON - 8 ozs.

MILO, SORGHUM, OKRA-4 ozs.

ALFALFA, CLOVER - 2 pzs.

BARLEY - 4 ozs.

SPINACH, SUNFLOWER-4 ozs.

OATS - 2 ozs.

SAFFLOWER, RYE, SUDAN GRASS

-- 2 ozs.

FLAX - 2 ozs.

SOYBEAN - 2 ozs.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS

SPROUTS, RADISHES – 2 ozs.

WHEAT - 2 ozs.

ACCEPTED

UNDER THE FEDERAL INSECTICIDE FUNCICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO 207-6.77 SUBJECT TO Accordance Comments.

WARNING: May Be Fatal If Swallowed! Do not breathe dust. May Be Absorbed Through Skin! Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Work in a well-ventilated room. Do not feed treated seed to animals. Treated seed should be labeled as follows: "TREATED SEED. DO NOT USE FOR FOOD. FEED OR OIL." If soil temperature is extremely high due to low moisture, real hot weather and midsummer planting time, reduce dosage by one-hall. Avoid lengthy storage of treated seed to prevent loss of lindane. Where seed decay occurs, due to soil organism damage, combine with a protective fungicide. When container is empty, immediately wash thoroughly and destroy. Never re-use.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife. Dispose of excess treated seeds by burial. Do not contaminate water by cleaning of equipment, or disposal of wastes.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied. concerning this material

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

Chevron Chemical Company Ortho Division/San Francisco, Calif. 94120

Ortho Division/San Francisco, Calif. 94120
Richmond, Calif. Portland, Ore. Fresno, Calif. Des Moines, towa
Cherry Hill, N.J. Honolulu, Hawaii Orlando, Fla.
Form 6397-J USDA Reg. No. 239-677 Product 1860 Made in U.S.A.

ىر

9