


ORTHORIX[®] Foliage Spray



ORTHO

(FUNGICIDE-MITICIDE)

Active Ingredient By Wt.
Calcium Polysulfides 27.5%
Inert Ingredients 72.5%
Density Baume at 68°F. 29.9. Contains calcium and
sulfur expressed as gypsum 2.8 lbs./gal. Total sulfur
2.29 lbs./gal. Other combined sulfur 1.8 lbs./gal.
10.5 Lbs. Per Gal.

DANGER: KEEP OUT OF REACH OF CHILDREN.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE
WITH LABEL DANGER STATEMENTS AND DIRECTIONS, AND
WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT
CONCENTRATE OR DILUTE INTO FOOD OR DRINK CON-
TAINERS.

WHEN EMPTY REPLACE BUNGS SECURELY AND RETURN TO
MANUFACTURER.

ORTHORIX Foliage Spray is a calcium polysulfide fungicide and miticide. It is a foliar spray for use on a wide variety of plants. It is a foliar spray for use on a wide variety of plants. It is a foliar spray for use on a wide variety of plants.

NET CONTENTS 28 GALS. LIQUID

Chevron Chemical Company

Ortho Division/San Francisco CA 94119 Richton and CA 94804
Product 1872 Made in U.S.A.
Form 2480-W
EPA Reg. No. 239-709-AA

DIRECTIONS

APPLES AND PEARS: Scab and Mildew Control. Use 2 qts. in 100 gals. of water. Apply in Prebloom, Bloom and Calyx periods. **Powdery Mildew:** Continued spraying is required during growing season. Do not use ORTHORIX Foliage Spray on d'Anjou pears after the calyx period. FLOTOX Wettable Sulfur may be added to this spray at the rate of 3 lbs. per 100 gals.

CANE BERRIES: Red Berry Mite, Leaf Spot, Cane Blight. 3 qts. per 100 gals. water. Apply as a delayed dormant spray.

GRAPES: Powdery Mildew. 1 pt. ORTHORIX Foliage Spray plus 4 lb. FLOTOX Wettable Sulfur in 100 gals. water.

PEARS (California and Northwest only): **POST HARVEST AND LATE FALL TREATMENTS:** Pear Bud Mites, Blister Mites—Apply 2 qts. per 100 gals. ORTHORIX Foliage Spray per acre in 100 gals. of water for a thorough coverage application. **Fall Spray** (September-October), just prior to dormancy. Apply 2 qts. ORTHORIX Foliage Spray per acre in 100 gals. of water. **DELAYED DORMANT SEASON TREATMENT:** Pear

Leaf Blister Mites, Pear Psylla, Scale Insects, Aphid and Mite Eggs. Apply 2 qts. ORTHORIX Foliage Spray per acre combined with 3 to 5 qts. FLEETOP Flowable or 1 1/2 to 2 qts. VOLOCK Supreme Spray in each 100 gals. water. Use sufficient spray for thorough coverage application.

RED CLOVER AND ALFALFA: Powdery Mildew. 1 qt. in 50 to 100 gals. water per acre at bud stage.

ROSES (Nursery Stock and Field Grown Ornamentals): Powdery Mildew, Rust, Black Spot, Red Spider Mites. Use 1 1/2 pts. per 100 gals. of water on tender new growth and 1 1/2 qts. in 100 gals. of water if mildew becomes severe. Pick or in flower before spraying to avoid discoloration.

STONE FRUITS (CHERRIES, PEACHES, NECTARINES, PRUNES AND PLUMS): Brown Rot, Leaf Spot. Use 2 qts. per 100 gals. water. Apply 3 to 5 weeks before harvest repeated at weekly intervals until just before picking. **Powdery Mildew**—2 qts. per 100 gals. water, applied 2 weeks before use. Total of 4 lbs. sulfur per acre.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, quality or representation, expressed or implied, concerning this material. 2. Critical and unforeseeable factors beyond Chevron's control prevent it from assuming any risk in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though the directions are followed. Buyer and user assume entire and exclusive liability, except those assumed by Chevron, for the total liability from handling, storage, and use of this material.

DANGER:

Do not get in eyes. Do not get on face. Do not get on clothing. Do not get on skin. Do not get on hands. Do not get on feet. Do not get on shoes. Do not get on socks. Do not get on underwear. Do not get on pajamas. Do not get on nightgown. Do not get on robe. Do not get on towel. Do not get on blanket. Do not get on bedspread. Do not get on curtain. Do not get on rug. Do not get on carpet. Do not get on floor. Do not get on wall. Do not get on ceiling. Do not get on furniture. Do not get on appliance. Do not get on electronic equipment. Do not get on vehicle. Do not get on animal. Do not get on bird. Do not get on fish. Do not get on insect. Do not get on plant. Do not get on soil. Do not get on water. Do not get on air. Do not get on fire. Do not get on explosion. Do not get on poison. Do not get on disease. Do not get on death.

Note to Physicians: Emergency Information—(415) 233-7737. Do not use on animals as they are susceptible to anthrax. ORTHORIX Foliage Spray may burn foliage when temperatures high (85°F.) are made apparent on at least 3 times. When applied to crops, do not spray for 10 days before beginning treatments with ORTHORIX Foliage Spray and treatments with other spray. If frost occurs on surface of ORTHORIX Foliage Spray, break through. Crystallization does not impair product quality. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.