

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER

Corrosive to eyes. Concentrate causes eye damage and skin irritation. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid breathing spray mist.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, immediately flush eyes with fresh water for 15 minutes and see a doctor. Wash skin thoroughly with soap and water. If swallowed, give large quantities of water, induce vomiting and see a doctor immediately.

Note to Physicians: Emergency Information - call 415-233-3737.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

PHYSICAL OR CHEMICAL HAZARDS

MAY GIVE OFF HIGHLY TOXIC AND EXTREMELY FLAMMABLE HYDROGEN SULFIDE GAS IF MIXED WITH ACIDS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner not consistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

REENTRY STATEMENT

Keep unprotected persons out of operating areas or units where there may be danger of drift.

Do not enter treated areas without protective clothing until sprays have dried.

Certain states may require more restrictive reentry intervals. Consult your State Department of Agriculture for further information.

When using Ortho Foliage Spray, keep children and pets away from treated areas. Do not allow children or pets to play in treated areas until the spray has dried.



ORTHO

ORTHORIX[®]
Foliage Spray

(FUNGICIDE-MITICIDE)

Active Ingredient	By Wt.
Calcium Polysulfides	27.5%
Inert Ingredients	72.5%

Density Baume at 68° F. — 29.9. Contains calcium and sulfur expressed as gypsum—2.8 lbs./gal. Total sulfur 2.29 lbs./gal. Other combined sulfur 1.8 lbs./gal. 10.5 lbs. Per Gal.

KEEP OUT OF REACH OF CHILDREN

DANGER-PELIGRO

FOR AGRICULTURAL USE ONLY OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

NET CONTENTS
28 GALLONS

Chevron Chemical Company

Ortho Agricultural Chemicals Division
San Francisco CA 94120 7144 Richmond, CA 94804 3735
Product 1872 Made in U.S.A.

Form 2480 C2 EPA Est. 939-CA-1

EPA Reg. No. 239-709-AA

ORTHO

ORTHORIX® Foliage Spray

(FUNGICIDE-MITICIDE)

Active Ingredient By Wt.
Calcium Polysulfides 27.5%
Inert Ingredients 72.5%
Density Baume at 68° F. — 29.9. Contains calcium and sulfur expressed as gypsum—2.8 lbs./gal. Total sulfur 2.29 lbs./gal. Other combined sulfur 1.8 lbs./gal. 10.5 lbs. Per Gal.

**KEEP OUT OF REACH OF CHILDREN
DANGER-PELIGRO**

FOR AGRICULTURAL USE ONLY OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**NET CONTENTS
28 GALLONS**

Chevron Chemical Company

Ortho Agricultural Chemicals Division
San Francisco CA 94120-7144 Richmond CA 94804-0036
Product 1872 Made in U.S.A.
Form 2480-C2 EPA Est. 239-CA-
PA Reg. No. 239-709-AA

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment.
Open dumping is prohibited.

STORAGE

Keep pesticide in original container.
Do not put concentrate or dilute into food or drink containers.
For help with any spill, leak, fire or exposure involving this material, call day or night 415-233-3131.

PESTICIDE DISPOSAL

This product is acute. Follow proper disposal of excess pesticide, spray mixture, or residue in accordance with Federal Law. If these wastes cannot be disposed of in accordance with these instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Region Office for guidance.

CONTAINER DISPOSAL

Completely empty inner, striking and tapping sides and bottom to loosen clinging particles. Empty into application equipment. Do not reuse container. Dispose of container in accordance with instructions on label. State and local authorities should be contacted and container disposed of in the same manner.


ORTHORIX Foliage Spray is a balanced fungicide-insecticide composition containing special effective wetting agents. It is for Growing Season Spraying. For best results spray to cover all exposed plant parts and repeat as necessary.

Do not use on apricots as they are susceptible to sulfur injury. ORTHORIX Foliage Spray may burn foliage when temperature is high (85°). Do not make applications at such times. When applied to crops with foliage, at 30 days re-apply between treatments with ORTHORIX Foliage Spray and treatments with other spray, if crust occurs on surface of ORTHORIX Foliage Spray break through. Crystallization does not impair product quality.

DIRECTIONS

APPLES AND PEARS: Scab and Mildew Control—2 to 4 qt. per 100 gals. of water. Apply in Pre-bloom, Bloom and Post-bloom. **Powdery Mildew:** Continued spraying. Do not stop spraying until after harvest. Do not use ORTHORIX Foliage Spray on Pears after the calyx period.

CANE BERRIES: Red Berry Mite, Leaf Spot, Cane Blight—3 qt. per 100 gals. water. Apply in late bud and dormant spray.



GRAPES: Powdery Mildew—1 pt. ORTHORIX Foliage Spray plus 4 lbs. Wettable Sulfur in 100 gals. water.

PEARS (California and Northwest only): POST HARVEST AND LATE FALL TREATMENTS—Pear Bud Mites, Blister Mites—Immediately after harvest: Apply 2½ gals. ORTHORIX Foliage Spray per acre in sufficient water for a thorough coverage application on average size trees. **Fall Spray (September-October, just prior to Leaf Fall)—**Apply 2½ gals. ORTHORIX Foliage Spray per acre. Use sufficient spray for a thorough coverage application. **DE-LAYED DORMANT SEASON TREATMENT—Pear Leaf Blister Mites, Pear Psylla, Scale Insects, Aphid and Mite Eggs—**Apply 2½ gals. ORTHORIX Foliage Spray per acre combined with 3 to 5 gals. KLEENUP Flowable or 1½ to 2 gals. VOLCK Supreme Spray in each 100 gals. water. Use sufficient spray for thorough coverage application.

RED CLOVER AND ALFALFA: Powdery Mildew—1 gal. in 50 to 100 gals. water per acre at bud stage.

ROSES (Nursery Stock and Field Grown Ornamentals): Powdery Mildew, Rust, Black Spot, Red Spider Mites—Use 1½ pts. per 100 gals. of water on tender new growth and 1½ qts. in 100 gals. of water if mildew becomes severe. Pick open flowers before spraying to avoid discoloration.

STONE FRUITS (CHERRIES, PEACHES, NECTARINES, PRUNES AND PLUMS): Brown Rot, Leaf Spot—Use 2 qts. per 100 gals. water. Apply 3 to 5 weeks before harvest, repeating at weekly intervals until just before picking. **Powdery Mildew—**2 qts. per 100 gals. water, when disease appears, or use ½ pt. with 4 lbs. wettable sulfur.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed herein. Chevron neither makes, nor authorizes, any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Certain and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, and if completely killed, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all such and other risks except those assumed by Chevron under 1. No warranty is made by Chevron as to the use of this material.