


# ORTHO<sup>®</sup> 10 Dust



# ORTHO

## (FUNGICIDE)

Active Ingredient	By Wt.
Captan*	10%
Inert Ingredients	90%
*N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide.	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL, DRY PLACE  
PROTECT FROM EXCESSIVE HEAT

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING OR WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

This product is formulated for application by ground or air equipment.

**ALMONDS:** Brown Rot, Shot Hole, Scab, Leaf Blight—Use 40-50 lbs. per acre. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. May be used up to 12 days of harvest if hulls are not fed to dairy animals or animals being finished for slaughter.

**APPLES:** Apple Scab, Brooks Fruit Spot, Fly-specking, Bitter Rot, Black Rot, Botryosphaeria, Black Pox—20 to 30 lbs. per acre. Apply at delayed dormant period. Repeat for each infection period or at weekly intervals as necessary.

**APRICOTS:** Brown Rot, Jacket Rot—50 lbs. per acre. Apply at red bud stage and repeat at 75 per cent petal fall.

**BEANS AND PEAS:** Damp-Off on Beans, Root Rot on Beans and Peas—35 to 40 lbs. per acre. Work into top 3 or 4 inches of soil before planting.

## NET WEIGHT 50 LBS.

### Chevron Chemical Company

Ortho Division/San Francisco, California 94119

Richmond, California Fresno, California

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 3962 Q2

Product 1982

Made in U.S.A.

EPA Reg. No. 239-728-AA

**CANTALOUPE, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELON, PUMPKINS:** Downy Mildew, Angular Leaf Spot, Anthracnose—Apply at 20 to 25 pounds per acre on young plants, increasing to 30 lbs. per acre on mature plants. Apply at first sign of disease and continue at weekly intervals as necessary.

**CELERY:** Pink Rot, Damp-Off Diseases—(Field)—20 lbs. per acre. (Seed Bed)—35 to 40 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

**CHERRIES:** Cherry Leaf Spot, Brown Rot—50 to 60 lbs. per acre. Apply at popcorn, blossom, and petal fall stages, continue at 10 to 14-day intervals.

**GRAPES:** Mold Control on Fruit on Drying Trays in Field—*Penicillium spp.*, *Aspergillus spp.*, *Botrytis Cinerea*, *Rhizopus spp.*—10 lbs. per acre. Apply to fruit on trays for protection of fruit and prevention of above molds. Apply to give even coverage. During damp weather make a second application after trays are turned. Apply this dust by hand or ground dusters only.

**GRAPES:** Botrytis Rot (Bunch Rot)—40 to 50 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically making up to 3 cover applications before bunches close.

**LETTUCE:** Downy Mildew (California and Arizona)—20 to 25 lbs. per acre. Apply at first sign of disease and continue at weekly intervals as necessary.

**MANGO:** *Cercospora Spot* or Blotch—50 to 70 lbs. per acre. Begin applications at first sign of disease. Repeat as necessary.

**ORNAMENTALS:** ROSES—Black Spot, CARNATIONS—Rust and Leaf Spot, CHRYSANTHEMUMS—Flower Blight—Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10-day intervals.

**PEACH, NECTARINES:** Brown Rot, Peach Scab—50 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make pre-harvest application where necessary.



## READ THE LABEL

### KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Avoid inhalation of dust. Avoid contact with skin. Do not store or transport near feed or food. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish. Keep out of streams and ponds. Do not apply where wind blowing toward water. In windy weather, conditions favor drift from areas where application is made. Rinse thoroughly with clean water by cleaning of equipment or spraying with water.

**PEARS:** Pear Scab (except Pacific Northwest)—20 to 30 lbs. per acre. Make first applications during early finger and petal fall stages. Repeat for each infection period or at weekly intervals as long as necessary. Russeting may be produced on Bosc Pears. Do not use on D'Anjou Pears.

**POTATOES:** (Seed Pieces): *Fusarium Seed Rot*, *Verticillium Wilt* and *Damp-Off*—Dust at 7½ ounces per bushel cut seed pieces (12 ozs. or ¾ lb. to 100 lbs. cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 days in open crates before bagging. Do not use treated seed for food or feed purposes.

**PLUMS, PRUNES:** Brown Rot—40 to 50 lbs. per acre. Apply at popcorn, full bloom, petal-fall and in covers as necessary. Thorough coverage is essential for maximum control.

**SOIL AND GREENHOUSE BENCH TREATMENT:** Preplanting for Damp-Off and Root Rot Diseases on Seedling or Transplants of Roses (and other shrubs and trees), Many Flowers and Vegetables (such as Beans, Peas, Cucurbits) and Lawn Seedbeds—1¼ to 2 lbs. per 1000 sq. ft. (50 to 60 lbs. per acre). Work into upper 3 or 4 inches of soil before planting.

**STOCK:** Botrytis Rot—40 to 50 lbs. per acre. Apply when disease appears. Repeat at weekly intervals as needed.

**STRAWBERRIES:** Botrytis Rot—30-35 lbs. per acre. Apply when new growth starts in spring before fruit starts to form. Repeat weekly. Severe Infection Conditions—Continue through harvest period treating immediately after each picking.

**TOMATOES:** Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot—Young Plants—25 to 30 lbs. per acre. Mature Plants—35 to 40 lbs. per acre. Begin applications at first sign of disease and continue at weekly intervals as necessary.

**TURF, LAWN:** Brown Patch, Leaf Spot, Damp-Off, Root Rot—1 to 2 lbs. per 1000 sq. ft. Treat before infection occurs. Repeat at 7 to 10-day intervals throughout summer.

**COMPATIBILITY:** Do not mix this product or other dusts, fungicides or sprays.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Certain and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to untreated crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the use stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1. above) resulting from the purchase, storage, and use of this material.