




# ORTHO ISOTOX<sup>®</sup> Insect Spray



READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

CONTAINS SPREADER

SYSTEMIC ACTION ENABLES PART OF THE INSECTICIDE TO BE ABSORBED INTERNALLY THROUGH LEAF AND STEM SURFACES TO PROVIDE LONGER, MORE EFFECTIVE PROTECTION.

GENERAL DIRECTIONS

For all listed uses, mix 2 Tablespoonfuls (1 oz.) ISOTOX Insect Spray in a gallon of water, unless otherwise specified. Spray plants and both sides of foliage thoroughly. Repeat as necessary. Do not store diluted spray. Make new dilution for each use. For more specific directions and cautions, read remainder of label. For best results, use an ORTHO SPRAY-ETTE Garden Hose Sprayer.

NOTE: DO NOT USE ON FRUITS, VEGETABLES, Boston Ivy, Mock Orange, Hawthorn & Chinese Elm.

### USE

ROSES,  
CAMELLIAS,  
AZALEAS,  
GARDENIAS,  
HIBISCUS,  
ORNAMENTALS,  
EVERGREENS,  
SHADE TREES,  
NURSERY PLANTS

### INSECTS

Aphids, Red Spiders, Armyworms,  
Tent Caterpillar, Chrysanthemum  
Fly, Diabrotica Beetles, Earwigs,  
Elm Leaf Beetle, Flea Beetles,  
Japanese Beetle, Cucumber Beetles,  
Grasshoppers, Lace Bugs, Box  
Elder Bugs, Leafhoppers, Leaf  
Miners, Leaf Rollers, Thrips,  
Mealy Bug, Bagworms, Plant Bugs,  
Psyllids, Rust Mite, Saw Fly  
Larvae, Scale (Crawlers),  
Spittle Bugs, Sowbugs, Whitefly,  
Woolly Aphid, Rose Chafer

### COMMENTS

Use 2 Table-  
spoonfuls per  
1 gal. water.  
  
Apply as pests  
appear. Repeat  
as necessary  
or at 2-week  
intervals

AROUND BUILDINGS,  
LAWNS, PATIOS,  
PICNIC AREAS,  
PARKS, PORCHES,  
GARBAGE CANS

Clover Mite (Brown  
Mite, Clover Spider  
Mite, or Brown Almond  
Mite), Ants, Black  
Widow Spiders, Crickets,  
Sowbugs, Earwigs, Box  
Elder Bugs

3 Tablespoonfuls per gal  
water. Spray around build-  
ing foundations and along  
fences, shrubbery areas,  
and other infested areas,  
applying 1 gal. diluted  
spray per 100 sq. ft. area.  
Repeat as necessary

COMBINATION SPRAY WITH PHALTAN ROSE & GARDEN FUNGICIDE ON ROSES — For Powdery Mildew, Black Spot and Listed Insects: Add 1 Tablespoonful PHALTAN Rose & Garden Fungicide and 1 Tablespoonful ISOTOX Insect Spray for each gallon of diluted spray. Always mix designated amount PHALTAN Rose & Garden Fungicide in a little water to form a thin, smooth paste before final dilution. To improve powdery mildew control use combination weekly during spring and fall mildew seasons. During hot summer months, apply PHALTAN Rose & Garden Fungicide at regular intervals for Disease Control, add ISOTOX Insect Spray only when necessary for insect control. To foliage feed plants, twice a month, add ORTHO Rose & Flower Food 8-12-4 at 2 teaspoonfuls per gal. to ISOTOX Insect Spray or the above combination. Note cautions on product labels.

**WARNING: MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN.** Do not breathe spray mist. Do Not Take Internally. Hazardous by skin absorption. Do not get in eyes, on skin or on clothing. Use waterproof gloves and face shield or goggles when handling concentrate. In case of eye contact, flush immediately with plenty of fresh water and get medical attention if irritation persists. Wash thoroughly with soap and warm water after using. Wash Hands and Face Before Eating. Food utensils such as teaspoons and tablespoons should not be used for food purposes after use with pesticides. Wash clothing with soap and hot water before re-use. Keep children and domestic animals off treated areas until surfaces have dried. **Note for Physician:** Atropine is antidotal. Do not contaminate feed and foodstuffs. **WARNING: FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME!** When container is empty, immediately wash thoroughly and destroy. Never re-use.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply to any area not specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

## Chevron Chemical Company


Ortho Division, San Francisco, Calif. 94119  
Richmond, Calif. Lafayette, Calif. Des Plaines, Ill. Perth Amboy, N.J. Atlanta, Ga.  
Farm 7223 0 Product 1984 Made in U.S.A.

EPA Reg. No. 239-741-ZC

\*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate. \*\* META-SYSTOX-R is the registered trade mark of Farbenfabriken Bayer A. G., Chemagro Corporation licensee. \*\*\*T.M. Reg. U.S. Pat. Off. by Rohm & Haas Co.



# Insect Spray



Chevron

# ORTHO ISOTOX<sup>®</sup> Insect Spray

DEL WARN

NTRATE OR

ABSORBED  
E LONGER,

r in a gallon  
s of foliage  
ew dilution  
ler of label.

ck Orange.

MENTS  
Table-  
fuls per  
water

as pests  
r. Repeat  
essary  
2-week  
als

er gal  
id build  
l along  
reas  
areas  
ed  
area  
y

**COMBINATION SPRAY WITH PHALTAN ROSE & GARDEN FUNGICIDE ON ROSES - For Powdery Mildew, Black Spot and Listed Insects:** Add 1 Tablespoonful PHALTAN Rose & Garden Fungicide and 1 Tablespoonful ISOTOX Insect Spray for each gallon of diluted spray. Always mix designated amount PHALTAN Rose & Garden Fungicide in a little water to form a thin, smooth paste before final dilution. To improve powdery mildew control use combination weekly during spring and fall mildew seasons. During hot summer months, apply PHALTAN Rose & Garden Fungicide at regular intervals for Disease Control, add ISOTOX Insect Spray only when necessary for insect control. To foliage feed plants, twice a month, add ORTHO Rose & Flower Food 8 12 4 at 2 teaspoonfuls per gal. to ISOTOX Insect Spray or the above combination. Note cautions on product labels.

**WARNING: MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN.** Do not breathe spray mist. Do Not Take Internally. Hazardous by skin absorption. Do not get in eyes, on skin or on clothing. Use waterproof gloves and face shield or goggles when handling concentrate. In case of eye contact flush **immediately** with plenty of fresh water and get medical attention if irritation persists. Wash thoroughly with soap and warm water after using. Wash Hands and Face Before Eating. Food utensils such as teaspoons and tablespoons should not be used for food purposes after use with pesticides. Wash clothing with soap and hot water before reuse. Keep children and domestic animals off treated areas until surfaces have dried. **Note for Physician:** Atropine is antidotal. Do not contaminate feed and foodstuffs. **WARNING: FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.** When container is empty, immediately wash thoroughly and destroy. Never reuse.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply to any area not specified on the label.


This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from you. Cooperative Agricultural Extension Service.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

**Chevron Chemical Company**

EPA Reg. No. 239-741-76

\*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate. \*\*META SYSTOX R is the registered trade mark of Farbenfabriken Bayer A. G., Chemagro Corporation licensee. \*\*\*T.M. Reg. U.S. Pat. Off. by Rohm & Haas Co.



**SYSTEMIC ACTION - Kills Aphids, Thrips, Mites, Japanese Beetles, Tent Caterpillar and Certain Other Insects**