



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

DANGER

Cause irreversible eye damage. May cause allergic skin reactions. Harmful if swallowed or inhaled. Do not get in eyes. Wear goggles when handling. Avoid contact with skin and clothing or breathing dust or vapor. Remove and wash contaminated clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, immediately flush eyes with fresh water for 15 minutes and see a doctor. If swallowed, give a large amount of water to drink, make person vomit, and call a doctor. For skin contact, wash with plenty of soap and water. See a doctor if irritation occurs or rash develops.

Note to Physicians: Emergency Information - call (415) 233-3737.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE

Keep pesticide in original container. Store in cool, dry place. Protect from excessive heat. Do not store or transport near feed or food.

PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

CONTAINER DISPOSAL

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by state and local authorities.



ORTHO

**ORTHO[®]
15 Dust**

(FUNGICIDE)

Active Ingredients	By Wt.
*Captan	14.2%
Related Derivatives	0.8%
Inert Ingredients	85.0%
*N-[trichloromethyl]thio]-4-cyclohexene-1,2-dicarboximide.	

KEEP OUT OF REACH OF CHILDREN

DANGER

FOR AGRICULTURAL USE ONLY OR FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NET WEIGHT 50 POUNDS

Chevron Chemical Company

Ortho Agricultural Chemicals Division
San Francisco CA 94120-7144 Richmond CA 94804-0036
Product 2176 Made in U.S.A.
Form 4386-W EPA Est. 239-CA-9
EPA Reg. No. 239-788-AA

DIRECTIONS

This product is formulated for application by ground or air equipment.

ALMONDS: Brown Rot, Shot Hole, Scab, Leaf Blight—Use 40 lbs. per acre. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. May be used up to 12 days of harvest if hulls are not fed to dairy animals or animals being finished for slaughter.

APPLES: Apple Scab, Brooks Fruit Spot, Botrytisphaeria, Flyspecking, Bitter Rot—15 to 25 lbs. per acre. Make first application at delayed dormant period. Repeat applications for each infection period or at weekly intervals as long as necessary.

APRICOTS: Brown Rot, Jacket Rot—33 lbs. per acre. Apply at red bud stage and repeat at 75 per cent petal fall.

BEANS AND PEAS: Damp-Off on Beans, Root Rot on Beans and Peas—35 to 40 lbs. per acre. Work into top 3 or 4 inches of soil before planting.

CANTALOUPE, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELON, PUMPKINS: Downy Mildew, Angular Leaf Spot, Anthracnose—Apply 15 to 20 lbs. per acre. Begin applications at first sign of disease. Repeat at 5 to 10-day intervals until danger of infection is past.

CELERY: Pink Rot, Damp-Off Diseases—(Field)—13 lbs. per acre. (Seed Bed)—25 to 30 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

CHERRIES: Cherry Leaf Spot, Brown Rot—35 to 40 lbs. per acre. Apply at popcorn, blossom, and petal fall stages, continue at 10 to 14 day intervals.

GRAPES: Botrytis Rot (Bunch Rot)—33 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically making up to 3 cover applications before bunches close.

MANGO: Cercospora Spot or Blotch—35 to 40 lbs. per acre. Begin applications at first sign of disease. Repeat as necessary.

ORNAMENTALS: ROSES — Black Spot. CARNATIONS — Rust and Leaf Spot. CHRYSANTHEMUMS — Flower Blight — Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

PEACH DISEASE: Brown Rot, Peach Scab—33 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

PEARS: Pear Scab (except Pacific Northwest)—15 to 25 lbs. per acre. Make first applications during early finger and petal fall stages. Repeat applications for each infection period or at weekly intervals as long as necessary. Russeting may be produced on Bosc Pears. Do not use on D'Anjou Pears.

POTATOES: Early and Late Blight—Begin applications when plants are 2 to 4 inches high and repeat at 7 to 10 day intervals as long as disease threatens. On young plants, apply 15 to 20 lbs. per acre.

POTATOES (Seed Pieces): Damp-Off—Dust at 5 oz. per bushel cut seed pieces (8 oz. or ½ lb. to 100 lbs. cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 days in open crates before bagging. Do not use treated seed for food or feed purposes.

PRUNES: Brown Rot—25 to 30 lbs. per acre. Apply at popcorn, full bloom, petal-fall and in covers as necessary. Thorough coverage is essential for maximum control.

SOIL AND GREENHOUSE BENCH TREATMENT: Preplanting for Damp-Off and Root Rot Diseases on Seedlings or Transplants of Roses (and other shrubs and trees), Many Flowers and Vegetables (such as Beans, Peas, Cucurbits) and Lawn Seedbeds—1¼ lbs. per 1000 sq. ft. (35 to 50 lbs. per acre). Work into upper 3 or 4 inches of soil before planting.

STRAWBERRIES: Botrytis Rot—20 lbs. per acre. Make first application when new growth starts in spring before fruit starts to form. Repeat these applications weekly. Under Severe Infection Conditions—Continue applications through harvest period, treating immediately after each picking.

TOMATOES: Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot—On young plants apply 15 to 20 lbs. per acre, increasing to 25 lbs. per acre on mature plants. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.

TURF AND LAWN: Brown Patch, Leaf Spot, Damp-Off, Root Rot—1 to 1½ lbs. per 1000 sq. ft. Begin treatment before infection occurs. Repeat at 7 to 10 day intervals throughout summer.

COMPATIBILITY: Do not use this product on foliage closely following oil sprays.

Do not use under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

