

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. May cause allergic skin reactions. Harmful if swallowed or inhaled. Do not get in eyes. Wear goggles when handling. Avoid contact with skin and clothing or breathing dust or vapor. Remove and wash contaminated clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, immediately flush eyes with fresh water for 5 minutes and see a doctor. If swallowed, give a large amount of water to drink, make person vomit, and call a doctor. For skin contact, wash with plenty of soap and water. See a doctor if irritation occurs or rash develops.

Note to Physicians: Emergency Information - call (415) 233-3737.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

REENTRY STATEMENT

Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift.

Do not enter treated areas without protective clothing until dusts have settled.

Certain states may require more restrictive reentry intervals, consult your State Department of Agriculture for further information.

Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas about to be treated.

DIRECTIONS

This product is formulated for application by ground or air equipment.

ALMONDS: Brown Rot, Shot Hole, Scab, Leaf Blight—Use 40 lbs. per acre. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. May be used up to 12 days of harvest if hulls are not fed to dairy animals or animals being finished for slaughter.

APPLES: Apple Scab, Brooks Fruit Spot, Botryosphaeria, Flyspecking, Bitter Rot—15 to 25 lbs. per acre. Make first application at delayed dormant period. Repeat applications for each infection period or at weekly intervals as long as necessary.

APRICOTS: Brown Rot, Jacket Rot—33 lbs. per acre. Apply at red bud stage and repeat at 75 per cent petal fall.

BEANS AND PEAS: Damp-Off on Beans, Root Rot on Beans and Peas—35 to 40 lbs. per acre. Work into top 3 or 4 inches of soil before planting.

CANTALOUPES, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELON, PUMPKINS: Downy Mildew, Angular

ORTHO

ORTHOCIDE® 15 Dust

(FUNGICIDE)

Active Ingredients	By Wt.
*Captan	14.2%
Related Derivatives	0.8%
Inert Ingredients	85.0%
*N-[(trichloromethylthio)-4-cyclohexene-1,2-dicarboximide]	

KEEP OUT OF REACH OF CHILDREN

DANGER—PELIGRO

FOR AGRICULTURAL USE ONLY. DO NOT
STORE IN AREAS ACCESSIBLE TO CHILDREN.

PRECAUCION AL USUARIO: Si usted no lee
ingles, no use este producto hasta que la etiqueta
le haya sido explicada ampliamente.

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.

NET WEIGHT 30 POUNDS

Chevron Chemical Company © 1984

Ortho Agricultural Chemicals Division

San Francisco CA 94120-7144

Richmond CA 94804-0036

Product 2176

Made in U.S.A.

Form 4386-X

EPA Est. 239-CA-9

EPA Reg. No. 239-788-AA

Leaf Spot, Anthracnose—Apply 15 to 25 lbs. per acre at first sign of disease. Repeat until danger of infection is past.

CELERY: Pink Rot, Damp-Off Diseases (Seed Rot)—25 to 30 lbs. per acre. Apply at germination. Repeat as necessary.

CHERRIES: Cherry Leaf Spot, Brown Rot—Apply at popcorn, blossom, and petal fall stages. Repeat at 14 day intervals.

GRAPES: Botrytis Rot (Bunch Rot)—33 lbs. per acre before bloom and 1 application after bloom. Repeat periodically making up to 3 applications per season.

MANGO: Cercospora Spot or Blotch—33 lbs. per acre at first sign of disease. Repeat at 7 to 10 day intervals.

ORNAMENTALS: ROSES—Black Spot, Leaf Spot. **CHRYSANTHEMUMS**—Flea sign of disease to give even coverage on all surfaces. Repeat at 7 to 10 day intervals.

PEACH DISEASE: Brown Rot, Peach Scab—2 to 3 prebloom applications. Repeat covers as necessary. Make preharvest application.

PEARS: Pear Scab (except Pacific Northwest)—2 to 4 inches high and repeat at 7 to 10 day intervals as long as necessary. Russeting Pears. Do not use on D'Anjou Pears.

POTATOES: Early and Late Blight—Begin applications 2 to 4 inches high and repeat as long as disease threatens. On young plants, apply 15 to 20 lbs. per acre.

POTATOES (Seed Pieces): Damp-Off—Dust seed pieces (8 oz. or 1/2 lb. to 100 lbs. per acre) in convenient container or by duster. Seed pieces should be treated within 6 hours of harvest. Do not be delayed more than 1 or 2 days. Do not store for 2 days in open crates before treating seed for food or feed purposes.

PRUNES: Brown Rot—25 to 30 lbs. per acre. Apply at full bloom, petal-fall and in covers coverage is essential for maximum control.

SOIL AND GREENHOUSE BENCH TREES: Damp-Off and Root Rot Diseases on 5 to 10 inch seedlings (such as Beans, Peas, Cucurbits) and 1000 sq. ft. (35 to 50 lbs. per acre). Apply 15 to 20 lbs. per acre of soil before planting.

STRAWBERRIES: Botrytis Rot—20 lbs. per acre. Apply when new growth starts in spring. Repeat these applications weekly. Under conditions—Continue applications through harvest immediately after each picking.

TOMATOES: Early Blight, Late Blight, Spot—On young plants apply 15 to 20 lbs. per acre on mature plants, 25 to 30 lbs. per acre. Repeat at 7 to 10 day intervals.

TURF AND LAWN: Brown Patch, Leaf Spot—1 to 1 1/2 lbs. per 1000 sq. ft. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.

COMPATIBILITY: Do not use this product in combination with low oil sprays.

Do not apply under conditions involving rain or other plantings that might be affected. Do not use if rendered unfit for sale, use or disposal.

ORTHO

THOCIDE® Dust

IDE)

By Wt.
Inert Ingredients 14.2%
Active Ingredients 0.8%
Total 85.0%
Methylthio]-4-cyclohexene-1,2-dicarboximide.

OUT OF REACH OF CHILDREN
DANGER - PELIGRO

CULTURAL USE ONLY. DO NOT
AREAS ACCESSIBLE TO CHILDREN.

ON AL USUARIO: Si usted no lee
este producto hasta que la etiqueta
p explicada ampliamente.

DE PANEL FOR ADDITIONAL
CAUTIONARY STATEMENTS.

EIGHT 50 POUNDS

ical Company © 1984

Chemicals Division
4720-7144 Richmond, CA 94804-0036

Made in U.S.A.

EPA Est. 239-CA-9

239-788-AA

Leaf Spot, Anthracnose—Apply 15 to 20 lbs. per acre. Begin applications at first sign of disease. Repeat at 5 to 10-day intervals until danger of infection is past.

CELERY: Pink Rot, Damp-Off Diseases—(Field)—13 lbs. per acre. (Seed Bed)—25 to 30 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

CHERRIES: Cherry Leaf Spot, Brown Rot—35 to 40 lbs. per acre. Apply at popcorn, blossom, and petal fall stages, continue at 10 to 14 day interval.

GRAPES: Botrytis Rot (Bunch Rot)—33 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically making up to 3 cover applications before bunches close.

MAANGO: Cercospora Spot or Blotch—35 to 40 lbs. per acre. Begin applications at first sign of disease. Repeat as necessary.

ORNAMENTALS: ROSES — Black Spot. CARNATIONS — Rust and Leaf Spot. CHRYSANTHEMUMS — Flower Blight — Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

PEACH DISEASE: Brown Rot, Peach Scab—33 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

PEARS: Pear Scab (except Pacific Northwest)—15 to 25 lbs. per acre. Make first applications during early finger and petal fall stages. Repeat applications for each infection period or at weekly intervals as long as necessary. Russeting may be produced on Bosc Pears. Do not use on D'Anjou Pears.

POTATOES: Early and Late Blight—Begin applications when plants are 2 to 4 inches high and repeat at 7 to 10 day intervals as long as disease threatens. On young plants, apply 15 to 20 lbs. per acre.

POTATOES (Seed Pieces): Damp-Off—Dust at 5 oz. per bushel cut seed pieces (8 oz. or ½ lb. to 100 lbs. cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 days in open crates before bagging. Do not use treated seed for food or feed purposes.

PRUNES: Brown Rot—25 to 30 lbs. per acre. Apply at popcorn, full bloom, petal-fall and in covers as necessary. Thorough coverage is essential for maximum control.

SOIL AND GREENHOUSE BENCH TREATMENT: Preplanting for Damp-Off and Root Rot Diseases on Seedlings or Transplants of Roses (and other shrubs and trees), Many Flowers and Vegetables (such as Beans, Peas, Cucurbits) and Lawn Seedbeds—1½ lbs. per 1000 sq. ft. (35 to 50 lbs. per acre). Work into upper 3 or 4 inches of soil before planting.

STRAWBERRIES: Botrytis Rot—20 lbs. per acre. Make first application when new growth starts in spring before fruit starts to form. Repeat these applications weekly. Under Severe Infection Conditions—Continue applications through harvest period, treating immediately after each picking.

TOMATOES: Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot—On young plants apply 15 to 20 lbs. per acre, increasing to 25 lbs. per acre on mature plants. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.

TURF AND LAWN: Brown Patch, Leaf Spot, Damp-Off, Root Rot—1 to 1½ lbs. per 1000 sq. ft. Begin treatment before infection occurs. Repeat at 7 to 10 day intervals throughout summer.

COMPATIBILITY: Do not use this product on foliage closely following oil sprays.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Open dumping is prohibited.

STORAGE

Keep pesticide in original container

Store in cool, dry place.

Protect from excessive heat.

Do not store or transport near feed or food

For help with any spill, leak, fire or exposure involving this material, call day or night (415) 233-3737

PESTICIDE DISPOSAL

This product is acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance

CONTAINER DISPOSAL

Completely empty bag into application equipment. Do not re-use container. Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

