



ORTHO

ORTHO-KLOR[®]10 Chlordane Dust

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DIRECTIONS: CAN BE USED AS A SPRAY—Mix 2 Tablespoonfuls in a little water to make a thin paste, add to 1 gallon water and agitate during application. Do not store diluted spray. Make new dilution for each use.

HOUSEHOLD

Ants: OUTSIDE USE—Dust or spray along runways, in cracks in cement walks and patios, around garbage containers, under porches, around outside house foundations. Dust on soil around base of ornamentals. Spot treat nests in lawns.

LAWNS, ORNAMENTAL TURF, GOLF COURSES

Japanese Beetle Grubs, White Grubs, Chafer Beetle Grubs: 1 pound to 700 sq. ft. Water



ORTHO

ORTHO-KLOR[®]10 Chlordane Dust

ACCEPTED
Aug 22 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTENOLIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. _____ SUBJECT
TO ATTACHED COMMENTS

CAUTION: Do not breathe dust or spray mist. Avoid skin contact. In case of contact with skin, wash with soap and water. Wash hands after use.

Avoid contamination of feed and foodstuffs. Do not contaminate drinking water. Do not use on Green or Spring Onions. Do not apply to Beets after the seedling stage if tops are to be used for food. Do not apply to Green or Snap Beans after pods start to form. Do not apply to Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kale, Collards, Cucumbers and Tomatoes after the edible parts start to form. Do not contaminate ornamental fish ponds. Dispose of container promptly when empty.

Toxic to fish and wildlife. Do not contaminate lakes, ponds, streams, woodlands and other non-crop areas. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Note: This product is sold by weight and package is full when packed, but due to fluffy nature of material is inclined to settle.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, Calif. Lafayette, Calif. Des Plaines, Ill. Perth Amboy, N.J. Atlanta, Ga.

Form 4380-L2 Product 2173 Made in U.S.A.

EPA Reg. No. 239-797-AA

HOUSEHOLD

Ants: **OUTSIDE USE**—Dust or spray along runways in cracks in cement walks and patios, around garbage containers under porches, around outside house foundations. Dust on soil around base of ornamentals. Spot treat nests in lawns.

LAWNS, ORNAMENTAL TURF, GOLF COURSES

Japanese Beetle Grubs, White Grubs, Chafer Beetle Grubs: 1 pound to 7 sq ft. Water thoroughly after application. **Lawn Moths, Sod Webworms, Cutworms, Armyworms, Crickets, Mole Crickets, Earwigs:** 1 pound to 700 sq ft. Sprinkle lightly after application. Keep children, pets and domestic animals off treated areas until treated areas are well watered and completely dried. **Note:** Birds feeding in treated areas may be killed.

ORNAMENTALS

Ants, Thrips, Sowbugs, Leaf Miners and Lygus Bugs: Dust or spray upper and lower leaf surfaces in early morning or evening when air is still. Repeat as necessary. **Armyworms, Crickets, Mole Crickets:** 1 pound per 700 sq ft. Apply to soil around base of plants and along rows. **Earwigs:** Use same amount, applying to soil around base of plants and along rows in damp, shady areas, around and under rocks, boards and similar places. **Japanese Beetle Grubs:** 1 pound per 450 sq ft. Work into top 4 inches of soil around base of plants. **Cutworms, Strawberry Root Weevil (Brachyrhinus), Black Vine Weevil:** Dust or spray generously around base of plants, working into top 4 inches of soil. Repeat as necessary. See caution.

VEGETABLES

Soil Application: Preplanting, Planting or Transplanting Time: Beans, Beets, Broccoli, Brussels Sprouts, Cauliflower, Collards, Cucumbers, Kale, Tomatoes. **Ants, Japanese Beetle and June Beetle Grubs, Cutworms, Wireworms.** Apply one pound per 450 square feet at periods indicated only.

Cabbage, Onions: Cabbage Maggot, Onion Maggot—Dust or spray generously around base of plants, work into top 4 inches of soil as a preplanting soil application at time of planting or transplanting only.

CAUTION: Do not breathe dust or spray mist. Avoid skin contact. In case of contact with skin, wash with soap and water. Wash hands after use. Avoid contamination of feed and foodstuffs. Do not contaminate drinking water. Do not use on Green or Spring Onions. Do not apply to Beets after the seedling stage if tops are to be used for food. Do not apply to Green or Snap Beans after pods start to form. Do not apply to Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kale, Collards, Cucumbers and Tomatoes after the edible parts start to form. Do not contaminate ornamental fish ponds. Dispose of container promptly when empty.

Toxic to fish and wildlife. Do not contaminate lakes, ponds, streams, woodlands and other non-crop areas. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Note: This product is sold by weight and package is full when packed, but due to fluffy nature of material is inclined to settle.

Chevron Chemical Company

Orlando, Florida 32817
Rt. 1, Box 100, P.O. Box 100, P.O. Box 100, N. Arlington, N.J. 07016
EPA Reg. No. 239-797-AA

EPA Reg. No. 239-797-AA

Dust

Effective Ant Control.
Also kills Beetle Grubs,
Earwigs, Mole Crickets, Lawn Moths,
Cutworms, Sod Webworms, Armyworms