


# ORTHO<sup>®</sup> Methoxychlor 75-3 Seed Protectant



# ORTHO

ACCEPTED  
FEB 13 1974  
UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC INSECTICIDES REGISTERED UNDER NO. 239-987. SUBJECT TO ATTACHED CONDITIONS.

## (INSECTICIDE-FUNGICIDE)

Active Ingredients	By Wt.
*Captan	75%
**Methoxychlor, Technical	3%
Inert Ingredients	22%
*N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide.	
**Equivalent to 2.64% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and .36% of other isomers and related compounds.	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

STORE IN COOL, DRY PLACE.  
PROTECT FROM EXCESSIVE HEAT.

WASH AND DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

ORTHO<sup>®</sup> Methoxychlor 75-3 Seed Protectant is a special formulation containing ORTHOCIDE (Captan) and Methoxychlor for seed treatment application. It is used in preventing seed rot and damp-off diseases on germinating seeds and seedlings and smut on onion and hulled sorghum seeds. It also protects treated seed against stored grain insects.

USE ISOTOX 75 SEED TREATER in combination with ORTHOCIDE Methoxychlor 75-3 Seed Protectant when treating seed to be planted in soil infested with seed corn maggot and wireworms.

SLURRY TREATMENT: When used in slurry treaters, do not allow mixtures to stand overnight.

GLADIOLUS CORMS: For Prevention of Bulb Rot—Use 1½ ozs. in 1 gal. water (1 lb. to 10 gals.). Soak bulbs in mixture 20 to 30 minutes.

## NET WEIGHT

### Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119  
Richmond, California Fresno, California  
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida  
Form 4675-P EPA Reg. No. 239-987 Product 2575 Made in U.S.A.

ONION SEED PELLETTING: Use 1 pound ORTHOCIDE Methoxychlor 75-3 Seed Protectant per 1 pound of seed.  
THE FOLLOWING TABLE gives amounts of ORTHOCIDE Methoxychlor 75-3 Seed Protectant to use on various seeds.

SEED	Dosage Rate	Lbs. Seed Treated/Lb. of Product
CORN—Field	1¼ oz/100 lbs. ¾ oz/bu	1280
Corn—Sweet	2¼ oz/100 lbs. 1½ oz/bu	600
WHEAT	1¼ oz/100 lbs. ¾ oz/bu	1280
RYE	1¼ oz/100 lbs.	910
OATS	2 oz/100 lbs. ¾ oz/bu	800
BARLEY	1½ oz/100 lbs. ¾ oz/bu	1070
FLAX	2½ oz/100 lbs. 1½ oz/bu	640
BEANS (Except Baby Limas), PEAS	1¼ oz/100 lbs. 1 oz/bu	910
CUCUMBERS CANTALOUPE	2 oz/100 lbs.	800
SQUASH PUMPKIN WATERMELON	1½ oz/100 lbs.	1200

**CAUTION**

**READ THE LABEL**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:** Avoid inhalation of dust. Wear a chemical cartridge respirator approved by the U.S. Bureau of Mines. Avoid contact with skin. Avoid contamination of feed and foodstuffs. Do not use on baby lima beans. Do not use treated seed for human consumption. Treated seed should be labeled as follows: "TREATED SEED, DO NOT USE FOR FOOD, FEED OR OIL." Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is toxic to fish. Keep out of lakes, streams or ponds. Dispose of excess treated seeds by burial. Apply this product only as specified on this label.

SEED	Dosage Rate	Lbs. Seed Treated/Lb. of Product
BEETS, SWISS CHARD	8 oz/100 lbs.	200
SUGAR BEETS	8 oz/100 lbs.	200
CLOVER	5½ oz/100 lbs.	290
ALFALFA	3½ oz/bu	
SOYBEANS	1½ oz/100 lbs. 1 oz/bu	960
COTTON— Acid Delinted	2 oz/100 lbs.	800
COTTON— Machine Delinted Fuzzy (not delinted)	3 oz/100 lbs.	530
BLUE GRASS	5½ oz/100 lbs.	290
CABBAGE CAULIFLOWER BROCCOLI, BRUSSELS SPROUTS	1 oz/100 lbs.	1600
SPINACH	4 oz/100 lbs.	400
SORGHUM hulled (Kernel Smut)	3 oz/100 lbs.	530
PEPPERS	2 oz/100 lbs.	800
COWPEAS	2 oz/100 lbs. 1¼ oz/bu	800
RICE	2¼ oz/100 lbs.	710

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

## TYPE

239-987  
2/13/1974  
Page 121