

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

WHEN CONTAINER IS EMPTY, IMMEDIATELY WASH THOROUGHLY AND DESTROY. NEVER RE-USE.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

Pour recommended amount of this material into partially filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do not allow mixture to stand. Do not combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being mixed has proven them compatible.

USE INDICATED DOSAGE IN 100 GALLONS WATER UNLESS OTHERWISE SPECIFIED. Spray thoroughly.

Not for use or storage in or around the home.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION—Vegetable and Field Crops—30 to 250 gals. Cotton—3 to 20 gals. Deciduous Fruit Crops—60 to 800 gals. Citrus—400 to 3,000 gals. (On Young Fruit Plantings—Use a minimum of 15 gals. water per acre.)

Where Recommended for AIR APPLICATION—Field Crops—3 to 10 gals. Orchard Crops—10 to 20 gals.

ALFALFA, CLOVER: Pea Aphid, Potato Leafhopper, Spotted Alfalfa Aphid, Two-Spotted Mite—1 to 2 pints. Meadow Spittlebug—2 pints. Do not make more than 1 application per cutting. Do not graze or cut for hay within 28 days after application.

APPLE, CRAB APPLE, PEAR, QUINCE: SUMMER SPRAYS: Apple Aphid, Codling Moth, Leafhoppers, Rosy Apple Aphid, Woolly Apple Aphid, European Red Mite, Two-Spotted Mites, Clover (Brown Almond) Mite—1/2 pt. Use at least 2 pts. but not more than 2 1/2 pts. per acre. Apply only after foliage is mature. (30). DORMANT OR DELAYED DORMANT SPRAYS: Forbes Scale, Oystershell Scale, Putnam Scale, Scurfy Scale, San Jose Scale, Pear Psylla and Overwintering Mite and Aphid Eggs—1/2 to 1 pt. plus 1 to 2 gals. of KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray. Do not apply more than 250 gallons diluted spray.

ALMOND: SUMMER: Clover (Brown Almond) Mite, European Red Mite, Two-Spotted Mites—1/2 pint to 100 gals. water. 1 quart per acre in 20 gals. water by air. DORMANT: Olive Scale, European Fruit Lecanium, San Jose Scale, Overwintering Mite and Aphid Eggs—1/2 to 1 pint plus 1 to 2 gallons KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray. Apply a minimum of 1 quart per acre. Do not apply more than 500 gallons diluted spray. If almond hulls are to be used for feed, do not apply within 7 days of harvest.

APRICOT: SUMMER: Codling Moth, Two-Spotted Mites, Clover (Brown Almond) Mite, European Red Mite—1/2 pint. DORMANT: San Jose Scale, Soft Brown Scale, European Fruit Lecanium and Overwintering Mite Eggs—1/2 to 1 pint plus 1 to 2 gallons KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray. Apply a minimum of 1 quart per acre. Do not apply more than 400 gallons diluted spray. Do not apply within 30 days of harvest.

SNAP BEANS, LIMA BEANS: Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two-Spotted Mites—1 pint per acre in water to cover. Do not apply within 7 days of harvest. Do not make more than 2 applications per season. Do not feed treated forage to livestock.

BEANS (dry): Bean Aphid, Mexican Bean Beetle, Potato Leafhopper, Two-Spotted Mites, Serpentine Leaf Miner—1 pint per acre in water to cover. Do not apply more than 2 applications per season. Do not feed treated forage to livestock.

CANTALOUPE, WATERMELONS: Melon Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mites—1 to 2 pts. per acre in water to cover. Do not apply within 5 days of harvest.

CHERRY: SUMMER: Cherry Rust Mite, European Red Mite, Clover (Brown Almond) Mite, Two-Spotted Mites—1/2 pt. DORMANT: European Fruit Lecanium, Forbes Scale, San Jose Scale, Olive Scale, Overwintering Mite and Aphid Eggs—1/2 to 1 pint plus 1 to 2 gallons KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray. Apply a minimum of 1 quart per acre. Do not apply more than 400 gallons diluted spray. Do not apply within 30 days of harvest.

CITRUS: Citrus Red (Purple) Mite, Citrus Rust Mite, Yuma Mite, Six-Spotted Mite, Texas Citrus Mite—1 to 2 pints. Do not apply more than 250 gallons diluted spray. Do not



ORTHO

TRITHION[®]* 4 Emulsive

(INSECTICIDE)

Active Ingredients

Carbophenothion** 45%

Aromatic Petroleum Derivative Solvent 46%

Inert Ingredients 9%

**S-[(p-chlorophenylthio) methyl] O,O-diethyl phosphorodithioate.

Contains 4 pounds Trithion Per Gallon

U. S. Pat. No. 2,793,224



DO NOT TAKE INTERNALLY

DO NOT INHALE

DO NOT GET ON SKIN

KEEP OUT OF REACH OF CHILDREN

WARNING:

MAY BE FATAL IF SWALLOWED. Do not get in eyes, on skin or clothing. In case of contact, wash immediately with soap and water; for eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. For commercial application, wear rubber gloves, protective clothing, rubber overshoes and goggles. Wash contaminated clothing with soap and hot water before re-use. Do not store or transport near feed or food. TRITHION may cause cholinesterase inhibition. ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops about to be harvested. Do not apply oil sprays to trees where soil moisture is low or where soil moisture has recently been low. Do not apply oil sprays when temperatures are in excess of 90° F. or in periods of low humidity caused by dry winds. Do not combine oil sprays with sulfur sprays or apply within 30 days following a sulfur spray. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged. Do not use on or around livestock. Do not use on or around poultry. Do not use on or around bees. Do not use on or around fish. Do not use on or around aquatic life. Do not use on or around wildlife. Do not use on or around domestic animals. Do not use on or around domestic birds. Do not use on or around domestic mammals. Do not use on or around domestic reptiles. Do not use on or around domestic amphibians. Do not use on or around domestic invertebrates. Do not use on or around domestic plants. Do not use on or around domestic fungi. Do not use on or around domestic bacteria. Do not use on or around domestic viruses. Do not use on or around domestic parasites. Do not use on or around domestic diseases. Do not use on or around domestic pests. Do not use on or around domestic weeds. Do not use on or around domestic insects. Do not use on or around domestic mites. Do not use on or around domestic nematodes. Do not use on or around domestic protozoa. Do not use on or around domestic helminths. Do not use on or around domestic arthropods. Do not use on or around domestic mollusks. Do not use on or around domestic annelids. Do not use on or around domestic chordates. Do not use on or around domestic vertebrates. Do not use on or around domestic mammals. Do not use on or around domestic birds. Do not use on or around domestic reptiles. Do not use on or around domestic amphibians. Do not use on or around domestic invertebrates. Do not use on or around domestic plants. Do not use on or around domestic fungi. Do not use on or around domestic bacteria. Do not use on or around domestic viruses. Do not use on or around domestic parasites. Do not use on or around domestic diseases. Do not use on or around domestic pests. Do not use on or around domestic weeds. Do not use on or around domestic insects. Do not use on or around domestic mites. Do not use on or around domestic nematodes. Do not use on or around domestic protozoa. Do not use on or around domestic helminths. Do not use on or around domestic arthropods. Do not use on or around domestic mollusks. Do not use on or around domestic annelids. Do not use on or around domestic chordates. Do not use on or around domestic vertebrates.

ACCEPTED
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER.
ED UNDER NO. 239-019 SUBJECT
TO ATTACHED COMMENTS.

ACCEPTED

Nov 3, 1972

ACCEPTED
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER.
ED UNDER NO. 239-019 SUBJECT
TO ATTACHED COMMENTS.

GROUND APPLICATION—Vegetable and Field Crops—30 to 250 gals. Cotton—3 to 20 gals. Deciduous Fruit Crops—60 to 800 gals. Citrus—400 to 3,000 gals. (On Young Fruit Plantings—Use a minimum of 15 gals. water per acre.)

Where Recommended for AIR APPLICATION—Field Crops—3 to 10 gals. Orchard Crops—10 to 20 gals.

ALFALFA, CLOVER: Pea Aphid, Potato Leafhopper, Spotted Alfalfa Aphid, Two-Spotted Mite—1 to 2 pints. **Meadow Spittlebug**—2 pints. Do not make more than 1 application per cutting. Do not graze or cut for hay within 28 days after application.

APPLE, CRAB APPLE, PEAR, QUINCE: SUMMER SPRAYS: Apple Aphid, Codling Moth, Leafhoppers, Rosy Apple Aphid, Woolly Apple Aphid, European Red Mite, Two-Spotted Mites, Clover (Brown Almond) Mite—1/2 pt. Use at least 2 pts. but not more than 2 1/2 pts. per acre. Apply only after foliage is mature. (30). **DORMANT OR DELAYED DORMANT SPRAYS:** Forbes Scale, Oystershell Scale, Putnam Scale, Scurfy Scale, San Jose Scale, Pear Psylla and Overwintering Mite and Aphid Eggs—1/2 to 1 pt. plus 1 to 2 gals. of KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray. Do not apply more than 250 gallons diluted spray.

ALMOND: SUMMER: Clover (Brown Almond) Mite, European Red Mite, Two-Spotted Mites—1/2 pint to 100 gals. water. 1 quart per acre in 20 gals. water by air. **DORMANT:** Olive Scale, European Fruit Lecanium, San Jose Scale, Overwintering Mite and Aphid Eggs—1/2 to 1 pint plus 1 to 2 gallons KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray. Apply a minimum of 1 quart per acre. Do not apply more than 500 gallons diluted spray. If almond hulls are to be used for feed, do not apply within 7 days of harvest.

APRICOT: SUMMER: Codling Moth, Two-Spotted Mites, Clover (Brown Almond) Mite, European Red Mite—1/2 pint. **DORMANT:** San Jose Scale, Soft Brown Scale, European Fruit Lecanium and Overwintering Mite Eggs—1/2 to 1 pint plus 1 to 2 gallons KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray. Apply a minimum of 1 quart per acre. Do not apply more than 400 gallons diluted spray. Do not apply within 30 days of harvest.

SNAP BEANS, LIMA BEANS: Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two-Spotted Mites—1 pint per acre in water to cover. Do not apply within 7 days of harvest. Do not make more than 2 applications per season. Do not feed treated forage to livestock.

BEANS (dry): Bean Aphid, Mexican Bean Beetle, Potato Leafhopper, Two-Spotted Mites, Serpentine Leaf Miner—1 pint per acre in water to cover. Do not apply more than 2 applications per season. Do not feed treated forage to livestock.

CANTALOUPE, WATERMELONS: Melon Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mites—1 to 2 pts. per acre in water to cover. Do not apply within 5 days of harvest.

CHERRY: SUMMER: Cherry Rust Mite, European Red Mite, Clover (Brown Almond) Mite, Two-Spotted Mites—1/2 pt. **DORMANT:** European Fruit Lecanium, Forbes Scale, San Jose Scale, Olive Scale, Overwintering Mite and Aphid Eggs—1/2 to 1 pint plus 1 to 2 gallons KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray. Apply a minimum of 1 quart per acre. Do not apply more than 400 gallons diluted spray. Do not apply within 30 days of harvest.

CITRUS: Citrus Red (Purple) Mite, Citrus Rust Mite, Yuma Mite, Six-Spotted Mite, Texas Citrus Mite—1/2 to 1 pint. Do not apply more than 750 gallons diluted spray. Do not apply within 14 days of harvest. **California Red Scale, Citricola Scale, Cottony Cushion Scale, Florida Red Scale, Purple Scale, Soft Scale, Yellow Scale**—1 to 1 1/2 pints plus 1 to 2 gallons ORTHOL-K READY-MIX Light Medium or ORTHOL-K Flowable Light Medium. Do not apply more than 650 gallons diluted spray. Do not apply within 30 days of harvest. Do not apply to kumquat or citron. TRITHION may cause injury to grapefruit if used in summer or fall before the fruit is fully colored. Allow 30 days between applications of TRITHION to citrus.

CORN (Field, Sweet): Grass Mite, Pacific Mite, Two-Spotted Mite—1 quart. Do not apply more than 1 quart of TRITHION 4 Emulsive per acre. Do not graze or cut corn forage within 21 days after application. Do not make more than 1 application per season.

COTTON: Cotton Aphid, Cotton Leaf Perforator, Cotton Fleahopper, Cotton Leafworm, Tropical Mite, Two-Spotted Mites, Cinnabar Mite—1 to 2 pints per acre in water to cover. Apply a minimum of 1 quart per acre for control of cotton leaf perforator and aphids. Do not graze dairy or meat animals in treated fields.

CUCUMBERS, SUMMER SQUASH: Melon Aphid, Potato Leafhoppers, Serpentine Leaf Miner, Two-Spotted Mites—1 to 2 pints. Do not apply within 7 days of harvest.

EGGPLANT, PEPPER, PIMENTO: Serpentine Leaf Miner, Two-Spotted Mites, Tropical Mite, Potato Flea Beetle, Potato Leafhopper—1 to 2 pints per acre in water to cover. Do not apply within 7 days of harvest.

FIGS: Two-spotted Mites—1/2 to 1 pint. Do not apply within 7 days of harvest.

Emulsive

(INSECTICIDE)

Active Ingredients

Carbophenothion** 45%

Aromatic Petroleum Derivative Solvent 46%

Inert Ingredients 9%

**S-[(p-chlorophenylthio) methyl] O,O-diethyl phosphorodithioate.

Contains 4 pounds Trithion Per Gallon

U. S. Pat. No. 2,793,224



DO NOT TAKE INTERNALLY

DO NOT INHALE

DO NOT GET ON SKIN

KEEP OUT OF REACH OF CHILDREN

WARNING:

MAY BE FATAL IF SWALLOWED. Do not get in eyes, on skin or clothing. In case of contact, wash immediately with soap and water; for eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. For commercial application, wear rubber gloves, protective clothing, rubber overshoes and goggles. Wash contaminated clothing with soap and hot water before re-use. Do not store or transport near feed or food. TRITHION may cause cholinesterase inhibition. ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops about to be harvested. Do not apply oil sprays to trees where soil moisture is low or where soil moisture has recently been low. Do not apply oil sprays when temperatures are in excess of 90° F. or in periods of low humidity caused by dry winds. Do not combine oil sprays with sulfur sprays or apply within 30 days following a sulfur spray. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not apply to any area not specified on the label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 5110-U Product 2539 Made in U.S.A.

EPA Reg. No. 239-1019-AA

NET CONTENTS 1 GALLON LIQUID

ACCEPTED

Nov 3, 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER.
ED UNDER NO. 239-1019 SUBJECT
IS ATTACHED COMMENTS.



ORTHO

TRITHION[®]* 4 Emulsive

(INSECTICIDE)

Active Ingredients

Carbophenothion**

Aromatic Petroleum Derivative Solvent

Inert Ingredients

By Wt.

45%

46%

9%

**S-[(p-chlorophenylthio) methyl] O,O-diethyl phosphorodithioate.

Contains 4 pounds Trithion Per Gallon

U. S. Pat. No. 2,793,224



DO NOT TAKE INTERNALLY

DO NOT INHALE

DO NOT GET ON SKIN

KEEP OUT OF REACH OF CHILDREN

WARNING:

MAY BE FATAL IF SWALLOWED. Do not get in eyes, on skin or clothing. In case of contact, wash immediately with soap and water; for eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. For commercial application wear rubber gloves, protective clothing, rubber overshoes and goggles. Wash contaminated clothing with soap and hot water before re-use. Do not store or transport near feed or food. TRITHION may cause cholinesterase inhibition. ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops about to be harvested. Do not apply oil sprays to trees where soil moisture is low or where soil moisture has recently been low. Do not apply oil sprays when temperatures are in excess of 90° F, or in periods of low humidity caused by dry winds. Do not combine oil sprays with sulfur sprays or apply within 30 days following a sulfur spray. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

ACCEPTED
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER.
ED UNDER NO. 239-1019 SUBJECT
NO ATTACHED COMMENTS.

ACCEPTED

ACCEPTED

Nov 3, 1972

ACCEPTED
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER.
ED UNDER NO. 239-1019 SUBJECT
NO ATTACHED COMMENTS.

GARDEN BEETS: Bean Aphid, Serpentine Leaf Miner, Two-Spotted Mites, Potato Flea Beetle— $\frac{1}{2}$ to 1 pint per acre in water to cover. Do not apply within 21 days of harvest, if tops are to be used for food or feed.

GRAPES: Grape Leafhopper, Two-Spotted Mites—1 pint per acre in water to cover. Do not apply within 30 days of harvest. Do not apply to sensitive varieties such as Charbono, Gamay, and Muscat.

OLIVE: Black Scale, Oleander Scale, Olive Scale, San Jose Scale— $\frac{1}{2}$ to 1 pint plus 1 to 2 gallons ORTHOL-K Flowable, Light Medium or ORTHOL-K READY-MIX Light Medium. Apply a minimum of 1 quart per acre. Do not apply within 60 days of harvest.

ONIONS From Seeds, Sets or Transplants (For Dry Bulb or Green)—Onion Maggot—Furrow Treatment—Muck Soils— $\frac{1}{2}$ to 2 qts. per acre. Mineral Soil—1 to $\frac{1}{2}$ quarts per acre. Apply at time of seeding or transplanting by spraying in the planting furrow. Mix in 100 gals. water and apply 3 gals. of mixture per 1,000 ft. of furrow. Keep mixture agitated.

ORNAMENTALS: Camellias, Carnations, Gladiolus, Evergreen and Deciduous Shrubs: Two-Spotted Mites, Spruce Spider Mite, Monterey-Pine Aphid, Rose Aphid, Soft Scale Insects— $\frac{1}{2}$ to $\frac{1}{2}$ pints.

PEACHES, NECTARINES: Summer Spray: European Red Mites, Brown Almond Mites, Two-Spotted Mites— $\frac{1}{2}$ to 1 pint per 100 gallons of water but use no more than 2 pounds actual Trithion per acre. 1 quart per acre in 20 gallons of water by air application. Do not apply within 30 days of harvest. **DORMANT:** San Jose Scale, European Fruit Lecanium, Olive Scale— $\frac{1}{2}$ to 1 pint plus 1 to 2 gallons of KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray. Apply a minimum of 1 quart per acre. Do not apply more than 400 gallons diluted spray. Do not apply within 30 days of harvest. **Overwintering Mite Eggs—(Calif. Only)** 1 pint.

PEAS: Pea Aphid, Serpentine Leaf Miner, Two-Spotted Mites—1 to $\frac{1}{2}$ pints per acre in water to cover. Do not apply within 7 days of harvest. Do not graze or feed crop residue to livestock.

PECAN: Two-Spotted Mites— $\frac{1}{2}$ pint. Apply a minimum of 1 quart per acre. Do not apply after husks open.

PLUM, PRUNE: **SUMMER:** Leaf Curl Plum Aphid, Leafhoppers, Mealy Plum Aphid, Two-Spotted Mites, Clover (Brown Almond) Mite, European Red Mite— $\frac{1}{2}$ pint. **DORMANT:** Olive Scale, European Fruit Lecanium, San Jose Scale, Overwintering Mite and Aphid Eggs— $\frac{1}{2}$ to 1 pint plus 1 to 2 gallons KLEENUP READY-MIX Oil Spray or KLEENUP Flowable Emulsion. Apply a minimum of 1 quart per acre. Do not apply more than 400 gallons diluted spray. Do not apply within 30 days of harvest.

POST-HARVEST APPLICATION TO DECIDUOUS FRUIT TREES (Listed on Label): To decrease overwintering populations of Aphids, Leafhoppers, Spider Mites— $\frac{1}{2}$ to 1 pint. Apply a minimum of 1 quart per acre. For control of San Jose Scale, European Fruit Lecanium, Aphid and Mite Eggs— $\frac{1}{2}$ to 1 pint plus 1 to 2 gallons of ORTHOL-K Flowable Light Medium or ORTHOL-K READY-MIX Oil Spray. Cover crop should not be grazed or fed to livestock.

SEED CROPS: Brown Wheat Mite, Bean Aphid, Pea Aphid, Spotted Alfalfa Aphid, Serpentine Leaf Miner, Two-Spotted Mites—1 to 2 pints per acre in water to cover. Do not graze or feed crop residue to livestock. Sugar Beet and crop can be treated up to 14 days of harvest and crop residue can be fed to livestock.

SORGHUM: Two-Spotted Mite—1 quart. Do not apply within 21 days after application. Limit application to 1 per acre.

SOYBEANS: Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two-Spotted Mites—1 to $\frac{1}{2}$ pints per acre in water to cover. Do not apply within 7 days of harvest. Do not feed treated forage to livestock.

SPINACH, BEET TOPS: Serpentine Leaf Miner, Two Spotted Mites, Bean Aphid— $\frac{1}{2}$ pint per acre, in water to cover. Do not apply within 21 days of harvest.

STRAWBERRY: Two-Spotted Mites, Strawberry Aphid—1 pint per acre in water to cover. Do not apply within 3 days of harvest.

SUGAR BEET: Bean Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two Spotted Mites—1 to 2 pints per acre in water to cover. Do not apply within 14 days of harvest.

TOMATO: Serpentine Leaf Miner, Tomato Russet Mite, Two-Spotted Mites, Tropical Mite, Potato Flea Beetle, Potato Leafhopper, Potato Psyllid—1 to 2 pints per acre in water to cover. Do not apply within 7 days of harvest.

WALNUTS: **SUMMER:** Codling Moth, European Red Mite, Two-Spotted Mites, Walnut Aphid and for suppression of European Fruit Lecanium—1 to 2 quarts per acre in

EMULSIVE

(INSECTICIDE)

Active Ingredients

Carbophenothion** 45%

Aromatic Petroleum Derivative Solvent 46%

Inert Ingredients 9%

**S-[(p-chlorophenylthio) methyl] O,O-diethyl phosphorodithioate.

Contains 4 pounds Trithion Per Gallon

U. S. Pat. No. 2,793,224



DO NOT TAKE INTERNALLY

DO NOT INHALE

DO NOT GET ON SKIN

KEEP OUT OF REACH OF CHILDREN

WARNING:

MAY BE FATAL IF SWALLOWED. Do not get in eyes, on skin or clothing. In case of contact, wash immediately with soap and water; for eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. For commercial application, wear rubber gloves, protective clothing, rubber overshoes and goggles. Wash contaminated clothing with soap and hot water before re-use. Do not store or transport near feed or food. TRITHION may cause cholinesterase inhibition. ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops about to be harvested. Do not apply oil sprays to trees where soil moisture is low or where soil moisture has recently been low. Do not apply oil sprays when temperatures are in excess of 90° F. or in periods of low humidity caused by dry winds. Do not combine oil sprays with sulfur sprays or apply within 30 days following a sulfur spray. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not apply to any area not specified on the label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

Chevron Chemical Company

Ortho Division San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 5110-U Product 2539 Made in U.S.A.

EPA Reg. No. 239-1019-AA

NET CONTENTS 1 GALLON LIQUID

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 239-1019 SUBJECT
TO ATTACHED COMMENTS.

By Wt.

45%

46%

9%

Flowable Emulsion or KLEENUP READY-MIX Oil Spray. Apply a minimum of 1 quart per acre. Do not apply more than 400 gallons diluted spray. Do not apply within 30 days of harvest. Overwintering Mite Eggs—(Calif. Only)—1 pint.

PEAS: Pea Aphid, Serpentine Leaf Miner, Two-Spotted Mites 1 to 1½ pints per acre in water to cover. Do not apply within 7 days of harvest. Do not graze or feed crop residue to livestock.

PECAN: Two-Spotted Mites—½ pint. Apply a minimum of 1 quart per acre. Do not apply after husks open.

PLUM, PRUNE: SUMMER: Leaf Curl Plum Aphid, Leafhoppers, Mealy Plum Aphid, Two-Spotted Mites, Clover (Brown Almond) Mite, European Red Mite—½ pint. DORMANT: Olive Scale, European Fruit Lecanium, San Jose Scale, Overwintering Mite and Aphid Eggs—½ to 1 pint plus 1 to 2 gallons KLEENUP READY-MIX Oil Spray or KLEENUP Flowable Emulsion. Apply a minimum of 1 quart per acre. Do not apply more than 400 gallons diluted spray. Do not graze or feed crop residue to livestock.

POST-HARVEST APPLICATION TO DECIDUOUS FRUIT TREES Listed on Label: To decrease overwintering populations of Aphids, Leafhoppers, Spider Mites ½ to 1 pint. Apply a minimum of 1 quart per acre. For control of San Jose Scale, European Fruit Lecanium, Aphid and Mite Eggs ½ to 1 pint plus 1 to 2 gallons of ORTHOLK Flowable Light Medium or ORTHOLK READY-MIX Spray. Cider crops should not be grazed or fed to livestock.

SEED CROPS: Brown Wheat Mite, Bean Aphid, Pea Aphid, Spotted Alfalfa Aphid, Serpentine Leaf Miner, Two-Spotted Mites—1 to 2 pints per acre in water to cover. Do not graze or feed crop residue to livestock. Cereal Beet seed crops can be treated up to 14 days of harvest and crop residue can be fed to livestock.

SORGHUM: Two-Spotted Mite 1 quart. Do not apply more than once within 21 days after application. Limit application to 1 per season.

SOYBEANS: Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two-Spotted Mites 1 to 1½ pints per acre in water to cover. Do not apply within 7 days of harvest. Do not feed treated forage to livestock.

SPINACH, BEET TOPS: Serpentine Leaf Miner, Two-Spotted Mites, Bean Aphid ½ to 1 pint per acre, in water to cover. Do not apply within 21 days of harvest.

STRAWBERRY: Two-Spotted Mites, Strawberry Aphid—1 pint per acre in water to cover. Do not apply within 3 days of harvest.

SUGAR BEET: Bean Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mites—1 to 2 pints per acre in water to cover. Do not apply within 14 days of harvest.

TOMATO: Serpentine Leaf Miner, Tomato Russet Mite, Two-Spotted Mites, Tropical Mite, Potato Flea Beetle, Potato Leafhopper, Potato Psyllid 1 to 2 pints per acre in water to cover. Do not apply within 7 days of harvest.

WALNUTS: SUMMER: Codling Moth, European Red Mite, Two-Spotted Mites, Walnut Aphid and for suppression of European Fruit Lecanium 1 to 2 quarts per acre in water to cover. Best results for scale insect suppression are obtained if application is made after eggs have hatched. DORMANT: Calico Scale, European Fruit Lecanium, Frosted Scale, Walnut Aphid, Overwintering Mite Eggs—1 to 2 quarts per acre plus 2 gallons KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray in water to cover. Do not apply after husks open. Do not apply more than 2 gals. of oil per acre.

CONDITIONS OF SALE: 1. Chevron Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty, or FITNESS FOR A PARTICULAR PURPOSE, guarantee or representation, expressed or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemical products, including, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants and crops. Such risks occur even though the product is reasonably fit for the use intended hereon, and even though label directions are followed. Buyer and user acknowledge that as to all risks and liability (except those assumed by Chevron in connection with the storage, handling, storage, and use of this material).

*T.M. Reg. Stauffer Chemical Company, Stauffer Chemical Co.