

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Causes eye irritation. Harmful if inhaled. May cause allergic skin reaction. Avoid contact with eyes, skin and clothing. Avoid breathing dust or spray mist.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, immediately flush eyes with fresh water for 15 minutes. If irritation continues, see a doctor. Wash skin thoroughly with soap and water. If a skin rash develops, see a doctor. If respiratory irritation or any signs or symptoms occur from inhalation, move the person to fresh air. If these effects continue, see a doctor.

Note to Physicians: Emergency Information - call (415) 233-7337.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated.

DIRECTIONS FOR USE

Read and follow all directions on this product in a manner consistent with its labeling. Read entire label. Use strictly in accordance with PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Protect from excessive heat. Do not store or transport near food or feed.

PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

ORTHO

PHALTAN® 50 Wettable

Folpet FUNGICIDE

Active Ingredient: Folpet 50% By Wt.
Inert Ingredients: 50%
Net Weight: 40 LBS.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

BEST DOCUMENT AVAILABLE

NET WEIGHT 40 LBS.
ACCEPTED

SEP 15 1982

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 227-1162

USE DESIGNATED AMOUNT IN 100 GALS. WATER UNLESS OTHERWISE SPECIFIED.

APPLES, CRABAPPLES: Bitter Rot, Botryospheria, Alternaria Spot, Brooks Fruit Spot, Sooty Blotch, Black Rot, Fly Speck, Scab—1½ to 2 lbs. Apply up to 800 gals. per acre. Apply at delayed dormant and repeat at 7 to 14 day intervals until harvest. Caution—East—Use during the critical period (just before bloom until 30 days after petal fall) may cause russeting to sensitive varieties such as Golden Delicious, Red Delicious and Stayman Winesap. West—Do not use between bloom and July 1 to avoid injury.

AVOCADOS: Scab—3 lbs. Apply up to 1,000 gals. per acre. Begin applications when the bloom buds begin to swell in the very susceptible varieties and during late bloom when some fruit has set in the slight and moderately susceptible varieties. Continue to apply at 2 to 3 week intervals as needed.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES: Fruit Rot, Gray Mold, Spur Blight—2 lbs. Apply 250 gals. per acre. Apply when berries have set and repeat at 7 day intervals until harvest.

BLUEBERRIES, HUCKLEBERRIES: Gray Mold, Powdery Mildew, Leaf Spot Gloeosporium—2 lbs. Apply 250 gals. per acre. Apply before first infection. Repeat at 7 day intervals until harvest.

CELERY: Early Blight, Late Blight, Cercospora Leaf Spot—2 to 4 lbs. per acre in water to cover. Apply after plants are set in field and established. Repeat at 7 day intervals. Do not apply within 7 days of harvest.

CHERRIES (RED TART): Leaf Spot, Brown Rot—1 to 2 lbs. (1 lb. plus ½ pt. CHEVRON Spray Sticker in New York and other states where recommended). Apply up to 500 gals. per acre. Apply at prebloom and popcorn stage. Repeat in bloom and at 7 to 14 day intervals making last application to foliage postharvest. Add ½ pt. CHEVRON Spray Sticker per 100 gals. water to the sprays. Do not use on sweet cherries as injury may occur.

CITRUS (Oranges, Grapefruit, Lemons, Limes, Tangelos, Tangerines): Melonose, Scab, Greasy Spot—2 lbs. Apply up to 1,000 gals. per acre. Apply dormant, ½ petal fall, 2 weeks later and on the fall flush in August and September.

CRANBERRIES: Storage Rot—9 lbs. per acre. Apply during bloom and repeat at 10 to 14 day intervals. Do not apply within 30 days of harvest.

CUCUMBERS: Fruit Rots (Rhizoctonia & Pythium)—6 to 8 lbs. per acre applied in a minimum of 200 gals. of water as a directed spray at the soil in the row. Apply when plants are beginning to "run." Downy Mildew, Powdery Mildew, Anthracnose—2 to 4 lbs. per acre. Add ½ pt. CHEVRON Spray Sticker per 100 gals. water to the sprays. Begin at first true leaf stage. Repeat every 5 to 10 days depending upon disease pressure until harvest.

GRAPES: Downy Mildew, Powdery Mildew, Black Rot, Dead Arm—2 lbs. Apply 250 to 300 gals. per acre. Dead Arm—East—Apply when new growth is 1 to 2 inches long, repeat when growth is 4 to 6 inches long. West—first application bud break to 1 inch shoot length. Second application 2 weeks later or 4 to 8 inch shoot length. Black Rot, Downy Mildew—Apply just before bloom, repeat immediately after bloom. Depending upon the susceptibility of the grape variety from 1 to 3 more sprays may be needed at 7 to 10 day intervals. Powdery Mildew—East only—Apply 2 to 3 weeks after bloom and repeat at 2 week intervals as needed. Note: During extended periods of hot dry weather, foliage injury to several varieties of grapes may occur.

GOOSEBERRIES, CURRANTS: Leaf Spot—4 lbs. Apply 150 gals. per acre. Apply just before bloom period. Repeat at 7 to 10 day intervals in the postbloom period.

LETTUCE: Downy Mildew—2 lbs. Apply up to 200 gals. per acre. Apply when plants are established in field. Repeat at 7 day intervals until harvest.

MELONS: SUMMER: Mildew, depending and repeat ONIONS: dew—Use infection. before harvest ORNAMENTAL: Septoria AND IRIS: naria Leaf MARIGOLD: DRAGON: at 7 to 11 Thoroughly set (Gr and use 1 in 10 do to soak to not use w. TOMATO: Apply do until harvest.

Do not apply to water. COMPAT most of the exc time, with table. Fl with or limited COND: warrants on the Chevron tive to ABILITY, ing this 2. Crit. prevent of cher plants plete c crops. for the followe liability ing from

Ch On Sa Pa Fo EF

BEST D

ORTHIO

PHALTAN® 50 Wettable

Folpet FUNGICIDE

Active Ingredients: Folpet 50%
Inert Ingredients: 50%
Net Weight: 40 lbs.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

BEST DOCUMENT AVAILABLE

NET WEIGHT 40 LBS.
ACCEPTED

SEP 15 1982

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 239-1762

DIRECTIONS

USE DESIGNATED AMOUNT IN 100 GALS. WATER UNLESS OTHERWISE SPECIFIED.

APPLES, CRABAPPLES: Bitter Rot, Botryospheria, Alternaria Spot, Brooks Fruit Spot, Sooty Blotch, Black Rot, Fly Speck, Scab—1½ to 2 lbs. Apply up to 800 gals. per acre. Apply at delayed dormant and repeat at 7 to 14 day intervals until harvest. Caution—East—Use during the critical period (just before bloom until 30 days after petal fall) may cause russeting to sensitive varieties such as Golden Delicious, Red Delicious and Stayman Winesap. West—Do not use between bloom and July 1 to avoid injury.

AVOCADOS: Scab—3 lbs. Apply up to 1,000 gals. per acre. Begin applications when the bloom buds begin to swell in the very susceptible varieties and during late bloom when some fruit has set in the slight and moderately susceptible varieties. Continue to apply at 2 to 3 week intervals as needed.

RASPBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, BLACKBERRIES: Fruit Rot, Gray Mold, Spur Blight—2 lbs. Apply 250 gals. per acre. Apply when berries have set and repeat at 7 day intervals until harvest.

BLUEBERRIES, HUCKLEBERRIES: Gray Mold, Powdery Mildew, Leaf Spot Gloeosporium—2 lbs. Apply 250 gals. per acre. Apply before first infection. Repeat at 7 day intervals until harvest.

CELERY: Early Blight, Late Blight, Cercospora Leaf Spot—2 to 4 lbs. per acre in water to cover. Apply after plants are set in field and established. Repeat at 7 day intervals. Do not apply within 7 days of harvest.

CHERRIES (RED TART): Leaf Spot, Brown Rot—1 to 2 lbs. (1 lb. plus ½ pt. CHEVRON Spray Sticker in New York and other states where recommended). Apply up to 500 gals. per acre. Apply at prebloom and popcorn stage. Repeat in bloom and at 7 to 14 day intervals making last application to foliage postharvest. Add ½ pt. CHEVRON Spray Sticker per 100 gals. water to the sprays.

Do not use on sweet cherries as injury may occur.

CITRUS (Oranges, Grapefruit, Lemons, Limes, Tangelos, Tangerines): Melanose, Scab, Greasy Spot—2 lbs. Apply up to 1,000 gals. per acre. Apply dormant, 35 petal fall, 2 weeks later and on the fall flush in August and September.

CRANBERRIES: Storage Rot—9 lbs. per acre. Apply during bloom and repeat at 10 to 14 day intervals. Do not apply within 30 days of harvest.

CUCUMBERS: Fruit Rot (Rhizoctonia & Pythium)—6 to 8 lbs. per acre applied in a minimum of 200 gals. of water as a directed spray at the soil in the row. Apply when plants are beginning to "run." Downy Mildew, Powdery Mildew, Anthracnose—2 to 4 lbs. per acre. Add ½ pt. CHEVRON Spray Sticker per 100 gals. water to the sprays. Begin at first true leaf stage. Repeat every 5 to 10 days depending upon disease pressure until harvest.

GRAPES: Downy Mildew, Powdery Mildew, Black Rot, Dead Arm—2 lbs. Apply 250 to 300 gals. per acre. Dead Arm—East—Apply when new growth is 1 to 2 inches long, repeat when growth is 4 to 6 inches long. West—First application bud break to 1 inch shoot length. Second application 2 weeks later or 4 to 8 inch shoot length. Black Rot, Downy Mildew—Apply just before bloom, repeat immediately after bloom. Depending upon the susceptibility of the grape variety from 1 to 3 more sprays may be needed at 7 to 10 day intervals. Powdery Mildew—East only—Apply 2 to 3 weeks after bloom and repeat at 2 week intervals as needed. Note: During extended periods of hot dry weather, foliage injury to several varieties of grapes may occur.

GOOSEBERRIES, CURRANTS: Leaf Spot—4 lbs. Apply 150 gals. per acre. Apply just before bloom period. Repeat at 7 to 10 day intervals in the postbloom period.

LETTUCE: Downy Mildew—2 lbs. Apply up to 200 gals. per acre. Apply when plants are established in field. Repeat at 7 day intervals until harvest.

MELONS (Cantaloupe, Watermelon, Honeydew), PUMPKINS, SUMMER AND WINTER SQUASH: Downy Mildew, Powdery Mildew, Anthracnose—2 to 4 lbs. Apply up to 200 gals. per acre depending on size of vines. Apply when first true leaves appear and repeat at weekly intervals until harvest.

ONIONS, GARLIC, LEEKS, SHALLOTS: Purple Blotch, Downy Mildew—Use 4 lbs. Apply 200 gals. per acre. Apply before first infection. Repeat at weekly intervals until tops begin to dry before harvest.

ORNAMENTALS: Powdery Mildew and Black Spot on ROSES, Septoria Leafspot, Didymella Leafspot on CHRYSANTHEMUMS AND IPIS; Rust and Alternaria Leafspot on CARNATIONS; Alternaria Leafspot, Powdery Mildew and Anthracnose on ZINNIA; MARIGOLDS; CHRYSANTHEMUMS, ASTERS, PHLOX AND SNAP DRAGONS—2 lbs. Apply when young leaves first appear. Repeat at 7 to 10 day intervals or as necessary throughout the season. Thorough coverage is required. Pythium and Root Rot on Poinsettia (Greenhouse)—2 lbs. per 100 gals. water plus a spreader and use 1 pt. per 8 inch pan. Apply at time of potting. Repeat in 10 days. Damp off on AZALEA Cuttings—1½ to 2 lbs. Allow to soak for 15 to 30 minutes before placing in rooting media. Do not use wetting agents on roses as injury may result.

TOMATOES: Anthracnose—4 lbs. Use up to 200 gals. per acre. Apply during first bloom and repeat at 7 to 10 day intervals until harvest.

Do not apply under conditions involving possible drift to food crops or other plants that might be damaged or the crops thereof rendered unfit for sale or consumption.

COMPATIBILITY: PHALTAN 50 Wettable may be combined with most commonly used insecticides, adjuvants and fungicides with the exception of strongly alkaline material such as hydrated lime, which reduces the fungicidal activity of PHALTAN 50 Wettable. PHALTAN 50 Wettable should not be used in combination with or closely following oil sprays. Observe all cautions and limitations on labeling of all products used in mixtures.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage and use of this material.

Chevron Chemical Company

Ortho Agricultural Chemicals Division
San Francisco CA 94119 Richmond CA 94801

Product 2632 Made in U.S.A.

Form 9716 A

EPA Est. 239-FJ-1

EPA Reg. No. 239-1102-AA

BEST DOCUMENT AVAILABLE