

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
DANGER
 Causes irreparable eye damage. May cause allergic skin reactions. Harmful if swallowed or inhaled. Do not get in eyes. Wear goggles or face shield when mixing. Avoid contact with skin and clothing. Avoid breathing of dust or vapor. Wash hands after use. Remove and launder clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT
 In case of eye contact, immediately flush eyes with fresh water for 15 minutes and see a doctor. If swallowed, give a large amount of water to drink, make person vomit, and call a doctor. For skin contact, wash with plenty of soap and water. See a doctor if irritation occurs or rash develops.
Note to Physicians: Emergency Information - call (415) 233-3737.

ENVIRONMENTAL HAZARDS
 This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

STORAGE AND DISPOSAL
PROHIBITIONS
 Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.
 Open dumping is prohibited.

STORAGE
 Keep pesticide in original container.
 Store in cool, dry place.
 Protect from excessive heat.
 Do not store or transport near feed or food.

PESTICIDE DISPOSAL
 Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

CONTAINER DISPOSAL
 Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

DIRECTIONS
 This product is formulated for application by ground or air equipment.

Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality.

Chevron

ORTHO

ORTHOCIDE®

Sulfur 10-25

Dust

ACCEPTED
 MAY 4 1982

(FUNGICIDE)

Active Ingredients	by weight
*Captan	5.0%
Related Derivatives	0.5%
Sulfur	25.0%
Inert Ingredients	65.0%
*N-[trichloromethyl.thio]-4-cyclohexene-1,2-dicarboximide.	

KEEP OUT OF REACH OF CHILDREN

DANGER

FOR AGRICULTURAL USE ONLY. DO NOT STORE IN AREAS ACCESSIBLE TO CHILDREN.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Chevron Chemical Company
 Ortho Agricultural Chemicals Division
 San Francisco CA 94120 7144 Richmond CA 94804 0036
 Product 2500 Made in USA
 Form 5218 V EPA Est. 239-CA-5
 EPA Reg. No. 239 1116 AA

NET WEIGHT 50 POUNDS

CELERY: Pink Rot, Damp-Off (Field)—20 lbs. per acre. (Seed and soil)—40 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

GRAPES: Botrytis Rot, Bunch Rot, Powdery Mildew, Silver Mite, Storage Rot—4 to 60 lbs. per acre. Make 2 applications before harvest and 1 application after bloom. Repeat period between applications as necessary, before harvest close.

PEACHES: Brown Rot, Storage Rot, Powdery Mildew, Silver Mite, Peach Scab—5 to 60 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at intervals of 3 to 4 covers as necessary. Make post-harvest application where necessary.

POTATOES: Early and Late Blight—Begin applications when plants are 2 to 4 inches high and repeat at 7 to 10 day intervals as long as diseases threaten. Apply 25 to 30 lbs. per acre.

ROSES: Field: Rust, Powdery Mildew—35 to 40 lbs. per acre. Apply to give even coverage of upper and lower leaf surfaces to protect from disease. Repeat at regular intervals as necessary to maintain control.

SOIL TREATMENT: Preplanting for Pink Rot, Damp-Off on Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Celery, Eggplant, Lettuce, Peas, Peppers, Spinach, Sweet Corn, Tomatoes—5 to 60 lbs. per acre. Dust or broadcast, then disk into first few inches of soil surface.

SPINACH: (Seedling stage only): Downy Mildew, Powdery Mildew—35 to 40 lbs. per acre. Apply when all seeds have germinated. Repeat at 7 to 10 day intervals as long as diseases are active.

STRAWBERRIES: Botrytis Rot, Powdery Mildew—30 lbs. per acre. Apply after picking. Repeat applications after picking as necessary.

TOMATOES: (Established Plants): Early Blight, Late Blight, Anthracnose, Russet Mite—On young plants, apply 25 to 30 lbs. per acre. Increase to 35 to 40 lbs. per acre on mature plants. Apply at first sign of insects and diseases. Repeat at 5 to 7 day intervals as necessary for control. Because of the sulfur content of this dust, first obtain permission from local health or fire department before using it on tomatoes intended for sale to such persons as children.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

COMPATIBILITY: Do not use this product on foliage closely adjacent to other crops.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed thereon. 2. This warranty does not authorize any agent or representative of Chevron to make any other warranty of FITNESS or MERCHANTABILITY, guarantee, or representation, express or implied, concerning this material.

3. Certain inherent and foreseeable factors beyond Chevron's control may result in some minor risks in connection with the use of this material. Such risks include, but are not limited to, damage to plants and crops to which the material is applied; lack of complete control; and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably well used in accordance with the label directions. 4. The user is advised to read and understand all risks involved in the use of this material and to assume all risks involved in the use of this material.