

# TRITHION<sup>®</sup> 3 Dust



# ORTHO

ACCEPTED

FEB 8 1974

## INSECTICIDE)

Active Ingredient: Carbophenothion  
Inert Ingredients: 97%  
S-methylchlorophenylthio methyl O,O-diethyl phosphorodithioate

By Wt.  
3%  
97%

U.S. Pat. No. 2,793,224

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH INSTRUCTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP OUT OF REACH OF CHILDREN. See back of container for complete directions.

KEEP BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

Do not use in or near areas where people or animals are present. Do not use in or near areas where people or animals are present.

**APPLE, CRABAPPLE, PEAR, QUINCE:** Apple Aphid, European Red Mite, Rosy Apple Aphid, Two-Spotted Mites, Woolly Apple Aphid, Red-Backed Mite. 30 to 40 pounds per acre. Apply at pre-pink stage. Do not apply within 14 days of harvest.

**ALMOND, APRICOT, CHERRY, NECTARINE, PEACH, PECAN, PLUM, PRUNE, WALNUT:** Aphids, Orchard Mites. 20 to 30 pounds per acre. Do not apply within 14 days of harvest. Do not apply within 14 days of harvest.

**BEAN, LIMA BEAN, SOYBEAN (in pod):** Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two-Spotted Mites. 20 to 30 pounds per acre. Do not apply within 7 days of harvest. Do not apply within 7 days of harvest.

**CANTALOUPE, WATERMELON:** Melon Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mites. 20 to 30 pounds per acre. Do not apply within 5 days of harvest.

**CITRUS (Grapefruit, Lemons, Limes, Oranges, Tangelos, Tangerines):** Citrus Red (Purple) Mite, Citrus Rust Mite, Six-Spotted Mite, Texas Citrus Mite, Yuma Mite. 35 to 60 lbs per acre. Allow 14 days between applications. Do not apply within 14 days of harvest. Do not apply to kumquats or citrons.

**COTTON:** Cotton Aphid, Cotton Leaf Perforator, Cotton Flea Hopper, Cotton Leafworm, Cinnabar Mite, Tropical Mite, Two-Spotted Mites. 10 to 30 lbs per acre. Apply 30 lbs for control of cotton leaf perforator and aphids. Do not graze dairy or meat animals in treated fields.

**CUCURBIT (SUMMER SQUASH):** Melon Aphid, Potato Leafhopper, Serpentine Leaf Miner, Two-Spotted Mites. 20 to 35 lbs per acre. Do not apply within 7 days of harvest.

**EGGPLANT:** Serpentine Leaf Miner, Potato Leafhopper, Potato Flea Beetle, Tropical Mite, Two-Spotted Mite. 20 to 35 pounds per acre. Do not apply within 7 days of harvest.

**FIGS:** Two-Spotted Mites. 30 to 40 lbs per acre. Do not apply within 7 days of harvest.

**GARDEN BEETS:** Bean Aphid, Leafhoppers, Serpentine Leaf Miner, Two-Spotted Mites, Potato Flea Beetle. 15 to 25 pounds per acre.

Do not apply within 21 days of harvest if tops are to be used for feed or feed.

**GRAPES:** Grape Leafhopper, Two-Spotted Mites. 20 to 25 pounds per acre. Do not apply within 20 days of harvest.

**ONIONS:** Onion Maggot. Bulb: Onion Maggot. 33 to 66 lbs per acre depending on soil types. Apply at the time of seeding in the planting furrow, using a combination air/fertilizer spreader, or other suitable application equipment.

**ORNAMENTALS:** Camellias, Carnations, Gladiolus, Violets, Evergreens, Deciduous Shrubs, Spider Mites, Aphids. 15 to 25 lbs per acre.

**PEA:** Pea Aphid, Serpentine Leaf Miner, Two-Spotted Mites. 20 to 25 lbs per acre. Do not apply within 7 days of harvest. Do not use in or near areas where people or animals are present.

**PEPPER, PIMENTO:** Serpentine Leaf Miner, Potato Flea Beetle, Potato Leafhopper, Tropical Mite, Two-Spotted Mites. 20 to 30 lbs per acre. Do not apply within 7 days of harvest.

**SEED CROPS:** Alfalfa, Clover, Beans, Peas, Sugar Beets. Brown Wheat Mite, Pea Aphid, Bean Aphid, Spotted Alfalfa Aphid, Serpentine Leaf Miner, Two-Spotted Mites. 20 to 30 pounds per acre. Do not feed crop residue to livestock. Cover crops should not be grazed or fed to livestock.

**SPINACH, BEET TOPS:** Serpentine Leaf Miner, Two-Spotted Mites, Bean Aphid. 15 to 25 pounds per acre. Do not apply within 21 days of harvest.

**STRAWBERRY:** Two-Spotted Mites, Strawberry Aphid. 20 to 30 pounds per acre. Do not apply within 3 days of harvest.

**SUGAR BEET:** Bean Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mites. 20 to 30 pounds per acre. Do not apply within 14 days of harvest.

**TOMATOES:** Serpentine Leaf Miner, Potato Psyllid, Potato Flea Beetle, Potato Leafhopper, Tomato Russet Mite, Two-Spotted Mites, Tropical Mite. 20 to 30 pounds per acre. Do not apply within 7 days of harvest.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

TRITHION<sup>®</sup> registered trademark of Chevron Chemical Company. For more information, contact your local dealer.



**CAUTION**

**READ THE LABEL**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:** HAZARDOUS IF SWALLOWED. Avoid breathing dust. Wear a respirator and goggles. Avoid contact with eyes. Wash hands, arms and face thoroughly with soap and water after handling or smoking. Wash all contaminated clothing with soap and hot water before re-use. Do not store or use near food or feed. TRITHION may cause cholinesterase inhibition. Atropine is antidote. 2-PAM is also antidote and may be administered in conjunction with Atropine. Do not apply, or allow to drift to areas occupied by unprotected humans, beneficial animals, or onto growing food and forage crops about to be harvested. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption. This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply to any area not specified on the label. This product is toxic to bees and should not be applied when bees are present on the plants being treated.

NET WEIGHT 50 LBS.

Chevron Chemical Company

San Francisco Division/San Francisco, California 94119

Richmond, California Fresno, California

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 5244-E2 Product 2584 Made in U.S.A.

PA Reg. No. 239-1187-AA