



DIBROM[®] 4 Dust

NET WEIGHT 50 LBS.
Naled Insecticide

Active Ingredient	By Wt.
*Naled	4%
Inert Ingredients	96%
*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate.	
U. S. Pat. No. 2,971,882	

Use within 90 days of manufacture only, as this product decomposes.

Date of Manufacture

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. KEEP PESTICIDE IN ORIGINAL CONTAINER.

**STORE IN COOL, DRY PLACE,
OUT OF DIRECT SUNLIGHT.**

BOTTLES CONTAIN 10 LBS. NET WEIGHT. SEE BOTTOM OF BOTTLE FOR DIRECTIONS AND REGULATIONS.

ORTHO DIBROM 4 Dust is an insecticide formulated with an organic phosphate with excellent insecticidal activity. Use 40 to 50 pounds per acre unless otherwise specified. This product is formulated for application by ground or air equipment. For Airplane Application, use heavier dosages. Begin application at first sign of insects. Repeat as necessary, apply from start and through coverage to obtain control of insects.

ADULT MOSQUITO CONTROL: Crop and Residential Areas, Tidal Marshes, Swamps, Woodlands, Feed Lots and Pastures. 8 to 10 lbs. per acre. Avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat at 7 to 10 day intervals. Do not pasture animals on treated areas within 4 days after application. Do not apply to crop areas within 4 days of harvest of any crop or as indicated. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources.

BEANS (Dry or Field): Aphids, Mites, Lygus Bugs, Thrips, Leafhoppers, Leaf Miners, Loopers. Do not apply within 4 days of harvest.

BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, BRUSSELS SPROUTS, KALE: Cabbage Looper, Imported Cab-



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after using. Do not store or transport near feed or food. Do not use on cherries as injury may result. May corrode metal dust equipment, clean equipment after use.

This product will kill fish and wildlife. Keep out of lakes, streams and ponds. This product is highly toxic to bees exposed to direct treatment or residues on crops. Birds feeding on treated areas may be killed. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not contaminate water by cleaning of equipment, or disposal of wastes. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not apply when weather conditions favor drift of dust to non-treated areas.

bageworm, Diamond-Back Caterpillar, Aphids. May be used 1 day before harvest on Broccoli, Cabbage, Cauliflower and Brussels Sprouts. Do not apply to Collards and Kale within 4 days of harvest.

CITRUS (Oranges, Lemons, Grapefruit): Aphids, Red Spider Mites, Citrus Cutworms, Thrips. Do not apply within 7 days of harvest. For California and Arizona—Do not apply within 1 day of harvest.

COTTON: Lygus, Cotton Leaf Perforator, Armyworms, Looper, Fleahoppers. 30 to 40 lbs. per acre. Apply at 7 to 10-day intervals, which corresponds to 20 to 30 days of handpicking.

CUCUMBERS SUMMER SQUASH: Cucumber Beetles, Aphids, Leaf Miners, Pickleworms, Armyworms, Loopers, Leafhoppers—40 to 50 lbs. per acre. May be used up to harvest.

GRAPES: Leafhoppers 40 to 50 lbs. per acre, Thrips—40 to 25 lbs. per acre, Drosophila 40 lbs. per acre. Apply pre-harvest at first sign of Drosophila or bunch rot. Repeat

at weekly intervals as necessary to maintain control. Do not apply within 4 days of harvest.

HOPS: Hop Looper—20 lbs. per acre. Do not apply within 4 days of harvest.

LETTUCE, SPINACH, CELERY, CHARD, TURNIP GREENS: Loopers, Aphids—37 lbs. per acre. May be used 1 day before harvest on Lettuce, Spinach, Turnip Greens and Chard. Do not apply to Celery within 4 days of harvest.

MELONS (Cantaloupes, Honeydews, Muskmelons, Watermelons), PUMPKINS, WINTER SQUASH: Loopers, Armyworms, Leafhoppers, Aphids, Lygus Bugs. May be used 1 day before harvest.

PEACHES: Western States Only — Aphids, Fruit Flies (Drosophila), Red Spider Mites, Peach Twig Borer, Lygus Bugs, Thrips—50 lbs. per acre. Do not apply within 4 days of harvest.

ROSES (Field Grown): Aphids, Mites, Tent Caterpillars, Thrips, Leaf Rollers—25 lbs. per acre.

STRAWBERRIES: Spider Mites, Aphids. Do not apply within 4 days of harvest.

TOMATOES (field): Fruit Flies (Drosophila sp.), Two-Spotted Mites, Aphids, Leafminers, Tomato Fruitworms, Hornworms—25 lbs. per acre. Make first application at 5 to 7 days before first picking and continue at 5 to 7-day intervals. May be used 1 day before harvest.

CONDITIONS OF SALE: 1. Chevron Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed thereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Certain and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the use stated hereon and even though the label directions are followed. Buyer assumes all risks and liability for such damage, and hereby releases Chevron and its agents from all liability for such damage and loss of the material.

CHEVRON CHEMICAL COMPANY

FCPM 5566 12

FFA Reg. No. 239-1280

ORTHO DIVISION — San Francisco, California 94119

Product 1758

Made in U. S. A.

RICHMOND, CALIFORNIA

HONOLULU, HAWAII

FRESNO, CALIFORNIA



DES MOINES, IOWA

PERTH AMBOY, N. J.

ORLANDO, FLA.