

239-1281

8-3-74

1 of 4

PPA

PPA

Copy A: causes eye and skin

Copy B: fatal

Copy C: Do not breathe

Copy D: ,birds and other

Copy E: Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on the label.

Copy F: DESTROY CONTAINER WHEN EMPTY.



ORTHO

DIBROM[®] 8

Emulsive

Naled Insecticide



READ THE LABEL
DO NOT TAKE INTERNALLY
DO NOT GET IN EYES
DO NOT GET ON SKIN
DANGER: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL DANGER STATEMENTS
NET CONTENTS 1 GALLON LIQUID

Best Available copy

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:
Ground Application—Vegetable and Field Crops—30 to 250 gals. Cotton—3 to 20 gals. Deciduous Fruit Crops—50 to 800 gals. Citrus—400 to 3,000 gals. [On Young Fruit Plantings—Use a minimum of 15 gals. water per acre.]

A short residual contact insecticide for kill of insect and mite adults and larvae present at the time of application. Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly alkaline materials such as lime or bordeaux.

[illegible]

COTTON: Types—4 to 1 pt. per acre by air or ground. Leaf Perforator, Fleashoppers, Armyworms, Bollworms, Loopers and For Suppression of Moths—1 pt. per acre by air or ground. Apply at 7 to 10 day intervals while pests persist. **FOR HEAVY POPULATION OF LISTED INSECTS:** Use $\frac{1}{2}$ pt. per acre by air or ground in control programs with other insecticides registered for use on cotton.

WATER: Ammonia— $\frac{1}{8}$ to 1 pt. to 100 gals. water. Use up to 10 days of harvest.
GRAPES: Dithionite, leafhoppers— $\frac{1}{2}$ to $\frac{3}{4}$ pt. per 100 gals. water. (100 to 130 gals. diluted spray per acre depending on size of vines.) Use only $\frac{1}{2}$ pt. per 100 gals. water on Muscat varieties. Do not use on Italia varieties as fruit injury may occur. Do not enter vineyards for 24 hours following treatment.

1200 g/kg. (dried spore per kg). Do not apply when temperature is over 90°F.

STRAWBERRIES: Leaf Miner, Spider Mites, Aphids, Omomyous Leaf Tier, Spittlebugs (Northwest) - 1 pt. per acre. May be used 1 day before harvest.

SUGAR BEETS: Spider Mites, Leafhoppers - 1 pt. per acre by air; water per acre. Do not apply within 5 days of harvest.

SUGAR BEETS: Spider Mites, Leafhoppers - 1 pt. per acre by air; water per acre. Do not apply within 5 days of harvest.

TOBACCO: Homomom - 1 pt. to 100 gals. water; use up to 200 gals. diluted spray per acre. Flys Beeties - 1 pt. per acre in water to cover. Apply at first signs of insects. Repeat as needed.

Spray plants thoroughly for head-to-toe (up to 200) gals diluted spray per acre. Do not apply when temperature is over 90° F. May be used 1 day before harvest!

FOREST AND SHADE TREES.—CONIFERS (Abies, Douglas Fir, Fir, Hemlock, Juniper, Pine, Spruce). Suppression of Red Spider Mites, Aphids, Spittlebugs, Zimmerman Moths, Juniper Webworms, Tussock Moths, Spruce Budworms, Adult Sierra Fir and Western Hemlock Bark Borer.

[illegible]

Chemical description on the label and in this Material Safety Data Sheet (MSDS) are for informational purposes only. This product is highly toxic, so users exposed to direct treatment of animals should wear protective clothing. NEVER RE-USE WHEN CONTAINER IS EMPTY. IMMEDIATELY WASH THOROUGHLY AND SECTION. NEVER RE-USE. CONDITIONS OF SALE: Chemicon Chemical Company (Chemicon) warrants that the material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chemicon does not warrant the purity or merchantability, quantities or representation, express or implied, concerning this product. Collectively, the material safety data sheet, container, and other materials of FITNESS FOR USE AS DIRECTED HEREON.

The users stated herein and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chernus under 1 above) resulting from handling, storage, and use of this material.

Richmond, Calif. Fresno, Calif. Orlando, Fla. Des Moines, Iowa Perth Amboy, N.J.
Form 5536-G3 Product 2738 Made in U.S.A.

MM-1071-SC7 .0M .fau HJ3

239-1281

4 of 4

239-1281

ACCEPTED
239-1281
AUG 3 1974
UNDER THE FEDERAL FUNGICIDE
FUNGICIDE AND ROBOTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 1281 SUBJECT
TO ATTACHED COMMENTS