



Naled Insecticide

Contains 8 pounds technical noled per gallon equivalent to 7.2 pounds noled

DO NOT TAKE INTERNALLY. DO NOT GET IN EYES. DO NOT GET ON SKIN. SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NET CONTENTS 1 GALLON

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION—Vegetable and Fruit Crops—Deciduous Fruit Crops—60 to 80⁰ Fruit Plantings—Use a minimum of 15 gals. (On Young Cotton—3 to 20 Citrus—400 to 3,000 gals (On Young Water per acre)

Where Recommended for AIR APPLICATION—field Crops—3 to 10 gals
A short residual contact insecticide for kill of insect and mite adults and larvae present at the time of application

Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly alkaline materials such as lime or bordeaux.

Do not spray on Nectarines Ornamental Cherries Liquidambar or Chrysanthemums as injury may occur.

MIXING DIRECTIONS: Add to spray tank when $\frac{1}{4}$ to $\frac{1}{2}$ full with agitator running and maintain agitation until spraying is completed. Make new dilution for each use. Wash metal equipment thoroughly after use to avoid corrosion.

ALMONDS: Peach Twig Borer—Apply only once during the dormant or delayed dormant period. Apply 1 pt. in combination with 1½ to 2 gals. supreme type oil or 2 to 3 gals. dormant oil spray per 100 gals. water in high gallonage spray equipment. Thorough coverage is required. Observe oil spray precautions.

BEANS, LIMA BEANS (Dry Form): SPINACH: Aphids, Spider Mites, Leafhoppers, Beet Beetles, Leafminers, Flea Beetles, Cutworms, Armyworms, Cucumber Beetles, Colorado Potato Beetles, Cabbageworms, Looper, Aphids, Spider Mites, Lygus, Leaf Miner; **GREEN BEANS:** Aphids, Spider Mites, Loopers, Nitidulids, Leafhoppers. 1 pt per acre in water to cover 100 gal's per acre is preferable. **Large Loopers, Stink Bugs.** 1 1/2 pts per acre in water to cover. Do not apply with n 1 day of harvest! Do not feed treated vines.

BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, KALE, COLLARDS. Imported Cabbageworm, Diamond-Back Caterpillars, Aphids—1 pt per acre. Looper—1 pt per acre. Do not apply within 1 day of harvest.

CELERY: Aphids, Looper, Armyworms, Salt Marsh Caterpillars, Leaf Miners. Use ground application only with fine mist spray. 1 pt. in 50 to 75 gals. water per acre on plants less than 1/4 grown. On larger plants use 1 1/2 pts. in 100 to 150 gals. water per acre. Do not apply within 1 day of harvest.

COTTON: Lygus - 1/2 to 1 pt. per acre by air or ground. Leaf Perforator, Fleashoppers, Armyworms, Bollworms, Loopers and For Suppression of Mites - 1 pt. per acre by air or ground. Apply at 7 to 14 day intervals while pests persist. **FOR HEAVY POPULATION OF LISTED INSECTS** - Use 1/2 pt. per acre by air or ground in control programs with other insecticides registered for use on cotton. Do not apply after first bolls open. Do not graze livestock in treated fields.

GRIZE LIVESTOCK in treated areas
CUCUMBERS, SUMMER SQUASH: Cucumber Beetles, Aphids, Spider Mites, Loopers, Thrips, Armyworms 1 qt. per acre in water to cover. Pickleworms, Leaf Miners 1 1/2 pts. per acre in water to cover. **By Air**—Use minimum of 10 pts. water per acre. **May** be used up to harvest.

EGGPLANT, PEPPERS: Aphids, Spider Mites, blister Beetles, Flea Beetles, L-of Miners, Whitefly - 1 pt per acre. May be used up to 1 day before harvest

GRAPES: Drosophila, Leafhoppers 1/2 to 2 1/2 pt. per 100 gal. water (100 to 130 gals. diluted spray per acre depending on size of vines) Use only 1/2 pt. per 100 gal. water on Muscat varieties. **CAUTION:** Post bloom applications can cause fruit russetting. Do not use on Italian varieties as fruit injury may occur. Do not enter vineyards for 24 hours.

GENERAL
Consult federal, state or local disposal authorities for approved alternative procedures.

Ortho Agricultural Chemical's Division, San Francisco, CA 94113, Richmond, CA 94804
Product 2778 Made in U.S.A.

Product 2738 Made in USA
R Form 9219 H

EPA Reg. No. 239-1281-AA

DIRECTIONS FOR USE

diluted spray per acre. Do not apply when temperature is over 90 °F. May be used 1 day before harvest.

WALNUTS: Aphids—1/2 to 1 qt to 100 gal's water. Do not apply within 10 days of harvest. Do not graze livestock in treated orchards.

GREENHOUSE, VAPOR TREATMENT OF ROSES, OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS: Spider Mites, Adult Whiteflies, Aphids, Leafrollers, Mealybugs
- 1 fl oz. per 10,000 cu. ft. when plants are dry. Apply undiluted on heat pipes for 17

160 f. In steam heated houses, close vents for at least 1 hour after pipes are hot.

With hot water systems, close house for at least 3 hours. Houses may remain closed overnight following treatment. When application is made to hot pipes, wear goggles and approved respirator. Ventilate house before workers re-enter. Wear respirator when

and approved respirator. Wear eye goggles when bees enter. Wear respirator when entering treated houses to ventilate. Protect hives of bees in cucumber and other houses by covering with plastic or rubber sheet during treatment and until house is ventilated. Do not apply to tomatoes or cucumbers less than 1 day before harvest.

Spider Mites (Resistant and Non-resistant) Make 3 to 4 treatments at 3 or 4 day intervals. Repeat as necessary. **Adult Whiteflies** Make 4 applications at 3 day intervals. Repeat as necessary. **Aphids** Apply when aphids appear. Repeat as necessary.

WARNING: Avoid overwatering and direct application to plants as injury may result.
White Butterfly and Golden Rapture Rose: Pink Champagne Chrysanthemum, Green
Wanderer, Jew Poinsettias and Begonias: Pink may be treated with DIBROM vapor.

FOREST AND SHADE TREES: CONIFERS (Arborvitae, Douglas Fir, Fir, Hemlock, Juniper, Pine, Spruce): Suppression of Two-Spot and Red Spider Mites, Aphids, Spittlebugs, Zimmerman Moths, Juniper Webworms, Tussock Moths, Spruce Budworms, Adult Sierra

21. Zimmerman moths, Juniper webworms, Tussock moths, Sp. Ltr. sawworms, Adult Shear Fir and Western Hemlock Bark Borers. **BROADLEAF TREES** (Ash, Birch, Black Walnut, Box Elder, Dogwood, Elm, Evergreen Pear, Flowering Plum, and Quince, Lvs. 4, Hag nolia, Maple, Oak, Sycamore, Willow) Suppression of Two Soot and Red Soot. White,

Aphids: Leaf Miners, Thrips, Elm Leaf Beetles, California Oakworms (California city), Spring and Fall Cankerworms, Fall Webworm, Oak W-burner, and Furze Moths !
Use when plastication is done. One to two per tree.

FRUIT FLIES (*Drosophila* sp.) in and around Food Processing Plants, Loading Docks, Cull Piles and Refuse Areas. Cider Mills. Also found in homes, with a propensity to infest areas close to refuse, cull piles, dumpsites, dairies, wineries, etc. and in the bins of more

These piles are processed by means of a continuous machine which separates the piles into three piles: one containing the good paper, one containing the paper which is not recycled, and one containing the paper which is recycled. The paper which is recycled is then used to produce new paper. The paper which is not recycled is used to produce new paper. The paper which is good is used to produce new paper.

GNAT, STABLE FLY (Dox Flies), AND HOUSE FLY CONTROL—RESIDENTIAL AREAS.

CROPS LISTED ON LABEL. SWAMP Island. Consult your State fish and Game Agency before applying this product. It is not necessary to avoid farm buildings. Make applications during peak of infestation and repeat as necessary. See crop chart. **USE FLY CONTROL—RESIDENTIAL AREAS.**

Make applications during peak of infestation and repeat as necessary. See crop recommendations for use limitations near harvest. This rate of application will kill shrimp. Do not apply in tidal or marsh waters which are important shrimp producing areas. **AIRCRAFT:** Rate: Apply 0.8 to 4.0 fl. oz. per acre diluted with water. Apply 2

Apply 0.8 to 4 fl. oz. per acre diluted with water. Apply 2 to 9 qts. of diluted spray per acre. West---Apply 1.6 to 4 fl. oz. per acre diluted with

