

239-1349

4-21-1971

1/2

Best Available COPY

NET WEIGHT
4 POUNDS


ORTHO®



SEVIN*

dust

[insecticide]



FOR LIVESTOCK AND GARDEN USE

Active Ingredient	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	5%
Inert Ingredients	95%
U. S. Pat. Nos. 2,993,478 - 3,009,855	

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS ON BACK

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1. above) resulting from handling, storage, and use of this material.

ACCEPTED

APR 21 1971

UNDER THE FEDERAL INSECTICIDE
HERBICIDE AND RODENTICIDE ACT
FOR EPONOMIC POISON REGISTERED
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

239-1349

2/2

7 ✓
Protective Beef
Cattle House & Sheep
Hops deleted



SEVIN 5 dust

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BEANS: Mexican Bean Beetle— $\frac{1}{4}$ lb. per 1,000 sq. ft. Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Velvetbean Caterpillar, Flea Beetles, Japanese Beetle— $\frac{1}{2}$ lb. per 1,000 sq. ft. Corn Earworm, Armyworms, Tarnished Plant Bug, Stink Bugs— $\frac{1}{2}$ to $\frac{3}{4}$ lb. per 1,000 sq. ft. May be applied up to harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI: Flea Beetles, Harlequin Bug— $\frac{1}{4}$ to $\frac{1}{2}$ lb. per 1,000 sq. ft. Armyworms, Imported Cabbageworm, Young Cabbage Loopers, Salt-Marsh Caterpillars, Corn Earworm— $\frac{1}{2}$ to 1 lb. per 1,000 sq. ft. Do not apply within 3 days of harvest.

CORN: Corn Earworm (Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations). Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetle, Leafhoppers, Flea Beetles— $\frac{3}{4}$ to 1 lb. per 1,000 sq. ft. For insects attacking silks and ears apply at 2 to 3-day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. May be applied up to harvest.

CUCUMBER, MELONS, PUMPKIN, SQUASH: Pickleworm, Melonworm— $\frac{1}{4}$ to $\frac{1}{2}$ lb. per 1,000 sq. ft. Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs— $\frac{1}{2}$ lb. per 1,000 sq. ft. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use SEVIN on watermelons in Florida. May be used up to harvest.

LETTUCE, CARROTS: Flea Beetles, Harlequin Bug, Leafhoppers— $\frac{1}{4}$ to $\frac{1}{2}$ lb. per 1,000 sq. ft. Six-Spotted Leafhopper— $\frac{1}{2}$ to $\frac{3}{4}$ lb. in a 5 to 7-day schedule. Armyworms, Cabbage Caterpillars, Corn Earworm, Spittlebugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs— $\frac{1}{2}$ to 1 lb. per 1,000 sq. ft. For best results in the control of Corn Earworm after lettuce heads begin to form use 1 lb. in a 5 to 7-day schedule. May be used up to harvest on carrots. Do not use on head lettuce within 3 days of harvest and to leaf lettuce within 14 days.

EGGPLANT, PEPPERS: Colorado Potato Beetle, Flea Beetle— $\frac{1}{4}$ to $\frac{1}{2}$ lb. per 1,000 sq. ft. Tomato Fruitworm, Tomato Hornworm, European Corn Borer, Lace Bugs, $\frac{1}{2}$ to 1 lb. per 1,000 sq. ft.

STRAWBERRIES: Meadow Spittlebug, Strawberry Leaf Roller, Strawberry Weevil— $\frac{1}{2}$ to 1 lb. per 1,000 sq. ft. Do not apply within 1 day of harvest.

TOMATO: Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer—1 to $\frac{1}{2}$ lbs. per 1,000 sq. ft.

POULTRY (Chickens, Partridges, Pheasants, Pigeons): Northern Fowl Mite, Lice and as a supplement to pre-use treatments for Chicken Mite, Fleas—1 lb. per 100 birds. Apply thoroughly by means of a shaker can or hand duster. Repeat in 4 weeks if necessary. Do not use on Poultry or game birds within 7 days of slaughter.

IN OR AROUND POULTRY HOUSES: Chicken Mite, Fleas, Bed Bugs—1 lb. per 40 sq. ft. of litter or floor space, and to roosts and adjacent areas by means of a shaker can, hand duster or power duster. Force dust into cracks and crevices. Repeat as necessary. Do not treat nest litter. Avoid direct contamination of eggs. Do not use in premise treatments within 7 days of slaughter.

DOGS, CATS: Brown Dog Ticks, Fleas—Dust liberally over the animal and rub thoroughly into the skin. Apply also to the feet between the toes and on the legs. Apply at weekly intervals if needed. Avoid treatment of kittens less than 4 weeks of age.

IN OR AROUND DOG AND CAT QUARTERS: Ticks, Fleas—Dust liberally in and around sleeping quarters, kennels and dog houses by means of a shaker can, hand or power duster. Force dust into cracks and crevices. Repeat as necessary, but not more often than once weekly. For other uses, see larger size bags.

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long-Sleeved Work Clothing. Change to Clean Clothing Daily. Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidote. AVOID CONTAMINATION OF FOOD, FEEDING TROUGHES AND WATERING RECEPTACLES.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.



CHEVRON CHEMICAL COMPANY

ORTHO DIVISION — San Francisco, Calif. 94120

RICHMOND, CALIF.

PORTLAND, ORE.

FRESNO, CALIF.

DES MOINES, IOWA

Form 6391-H

HONOLULU, HAWAII

CHERRY HILL, N. J.

ORLANDO, FLA.

USDA Reg. No. 239-1349

Product 2845

Made in U.S.A.