

TO OPEN-CUT HERE
[ICOM] OUTDOOR USE ONLY

ORTHO® SEVIN® 5 Dust
[ABN: ORTHO® SEVIN® brand carbaryl insecticide 5 Dust]

KILLS GARDEN PESTS IN THE GARDEN: Armyworms, Cutworms, Squash Bugs, Tomato Hornworms, Certain Other Beetles And Other Garden Pests As Listed.

KILLS FLEAS & TICKS ON DOGS AND CATS AND IN DOGHOUSES: Fleas and Ticks on Animals and other outdoor living areas.

Kills: Beetles, Bugs, Leafhoppers and Worms as Listed On Vegetables.
May be Used up to Harvest Unless Otherwise Noted.


Active Ingredient
Carbaryl (1-naphthyl N-methylcarbamate).....5%
Inert Ingredients95%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for additional precautionary statements on back of bag.

Specially formulated for residential use.
For home use only

NET WT 4 LBS / 1.81 kg



REC'D G.A./P/DPD1
96 DEC 26 P1:17

TO OPEN-CUT HERE

READ ENTIRE LABEL BEFORE EACH USE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

~~READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.~~

VEGETABLES & FRUITS

PLANTS	INSECTS/PESTS Amount To Use	WHEN TO HARVEST
BEANS	Mexican Bean Beetles 1/4 lb per 1,000 sq ft Leafhoppers, Bean Leaf Beetles, Cucumber Beetles, Western Bean Cutworms, Velvet Bean Caterpillars, Flea Beetles, Japanese Beetles 1/2 lb per 1,000 sq ft Corn Earworms, Armyworms, Tarnished Plant Bugs, Stink Bugs 3/4 lb per 1,000 sq ft	May be applied up to day of harvest.
CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER KOHLRABI	Flea Beetles, Harlequin Bugs 1/2 lb per 1,000 sq ft Armyworms, Imported Cabbageworms, Young Cabbage Loopers, Salt Marsh Caterpillars, Corn Earworms 1 lb per 1,000 sq ft	Do not apply within 3 days of harvest.
CORN	Corn Earworms, Corn Rootworm Adults, European Corn Borers, Fall Armyworms, Sap Beetles, Japanese Beetles, Leafhoppers, Flea Beetles 1 lb per 1,000 sq ft	May be applied up to day of harvest.
CUCUMBERS MELONS, PUMPKINS, SQUASH Do not use this product on watermelons in Florida.	Pickleworms, Melonworms 1/2 lb per 1,000 sq ft Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs 1/2 lb per 1,000 sq ft	May be applied up to day of harvest.
LETTUCE, CARROTS	Flea Beetles, Harlequin Bugs, Leafhoppers 1/2 lb per 1,000 sq ft Sixspotted Leafhoppers 3/4 lb per 1,000 sq ft Armyworms, Cabbage Caterpillars, Corn Earworms, Spittlebugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs 1 lb per 1,000 sq ft	May be applied up to day of harvest to carrots. Do not apply to head lettuce within 3 days of harvest or to leaf lettuce within 14 days of harvest.

POTATOES, EGGPLANT, PEPPERS	Colorado Potato Beetles, Flea Beetles, Leafhoppers 1/2 lb per 1,000 sq ft Tomato Fruitworms, Tomato Hornworms, European Corn Borers, Lace Bugs 1 lb per 1,000 sq ft	May be applied up to day of harvest.
STRAWBERRIES	Meadow Spittlebugs, Strawberry Leafrollers, Strawberry Weevils 1 lb per 1,000 sq ft	Do not apply within 1 day of harvest.
TOMATOES	Flea Beetles, Leafhoppers, Tomato Fruitworms, Fall Armyworms, Tomato Hornworms, European Corn Borers 1 1/2 lb per 1,000 sq ft.	May be applied up to day of harvest.

HOW TO APPLY

- Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces.
- Apply when air is calm to avoid drift and contact with eyes and skin.
- Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.
- For plants above chest height, consider using a liquid carbaryl product.

WHEN TO APPLY

- Apply dust at the first signs of insects/pests.
- For best results, Dust in early morning or late evening when temperature is low and air is still.
- Repeat/Reapply as necessary, but not more often than once weekly.
- May be applied up to harvest unless otherwise noted.
- **CORN:** Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations. For insects attacking silks and ears apply when first silks appear and continue until silks begin to dry. Three or more applications may be required depending on severity of infestation.
- **CUCUMBERS, MELONS, PUMPKINS, SQUASH:** Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

BEANS: Mexican Bean Beetle 1/4 lb per 1,000 sq ft. Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Velvetbean Caterpillar, Flea Beetles, Japanese Beetle 1/2 lb. per 1,000 sq ft. Corn Earworm, Armyworms, Tarnished Plant Bug, Stink Bugs 3/4 lb per 1,000 sq ft. May be applied up to harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI: Flea Beetles, Harlequin Bug 1/2 lb per 1,000 sq ft. Armyworms, Imported Cabbageworm, Young Cabbage Loopers, Salt Marsh Caterpillars, Corn Earworm 1 lb per 1,000 sq ft. Do not apply within 3 days of harvest.

CORN: Corn Earworm (Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations). Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetle, Leafhoppers, Flea Beetles 1 lb per 1,000 sq ft. For insect attacking silks and ears apply at 2 to 3 day intervals starting when first silks appear and continuing until silks began to dry. Three or more applications may be required depending on severity of infestation. May be applied up to harvest.

CUCUMBER, MELONS, PUMPKIN, SQUASH: Pickleworm, Melonworm 1/2 lbs per 1,000 sq ft. Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs 1/2 lb per 1,000 sq ft. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use this product on watermelons in Florida. May be used up to harvest.

LETTUCE, CARROTS: Flea Beetles, Harlequin Bug, Leafhoppers 1/2 lb per 1,000 sq ft. Sixspotted Leafhopper 3/4 lb in a 5 to 7 day schedule. Armyworms, Cabbage Caterpillars, Corn Earworm, Spittlebugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs 1 lb per 1,000 sq ft. For best results in the control of Corn Earworm after lettuce heads begin to form use 1 lb in a 5 to 7 day schedule. May be used up to harvest on carrots. Do not use on head lettuce within 3 days of harvest and on leaf lettuce within 14 days.

EGGPLANT, PEPPERS: Colorado Potato Beetles, Flea Beetles, Leafhoppers 1/2 lb per 1,000 sq ft. Tomato Fruitworm, Tomato Hornworm, European Corn Borer, Lace Bugs 1 lb per 1,000 sq ft.

STRAWBERRIES: Meadow Spittlebug, Strawberry Leafroller, Strawberry Weevil 1 lb per 1,000 sq ft. Do not apply within 1 day of harvest.

TOMATO: Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer 1 1/2 lb per 1,000 sq ft.

NON-COMMERCIAL POULTRY

For Chickens, Partridges, Pheasants, Pigeons: To control Northern Fowl Mite, Lice and as a supplement to premise treatments for Chicken Mites and Fleas: Use 1 lb per 100 birds. Apply thoroughly by means of a shaker can or hand duster. Repeat/Reapply in 4 weeks if necessary. Do not use on Poultry or game birds within 7 days of slaughter.

IN OR AROUND POULTRY HOUSES

To control Chicken Mite, Fleas, Bed Bugs: Use 1 lb per 40 sq ft of litter or floor space and to roosts and adjacent areas by means of a shaker can or hand duster or power duster. Force dust into cracks and crevices. Repeat/Reapply as necessary. Do not treat nest litter. Avoid direct contamination of eggs. Do not use in premise treatment within 7 days of slaughter.

DOGS AND CATS

To control Ticks and Fleas: Use product only outside or in a well ventilated area. Wearing household latex or rubber gloves, lightly dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. This product should not be applied by children. Dust lightly over the animal and rub thoroughly into the skin. Apply also to the feet between the toes and on the legs. Repeat/Reapply as necessary, but not more often than once weekly. Apply at weekly intervals if needed. **DO NOT USE ON PREGNANT DOGS.** Avoid treatment of Do not use on kittens and or puppies under 12 4 weeks of age. **NOTE:** Do not allow treated pets back on carpets or rugs in the house as this product may stain or discolor rugs and wall-to-wall carpets. Dust bedding and doghouse lightly in and around kennels and hog houses by means of a using a shaker can, hand or other dust applicator power duster. Force dust into cracks and crevices. Repeat/Reapply as necessary, but not more often than once weekly. **DO NOT USE IN THE HOME: DO NOT APPLY TO RUGS AND CARPETS AS STAINING MAY OCCUR.**

STORAGE AND DISPOSAL

STORAGE: Store in original container in a safe place. Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. **DISPOSAL:** Do not reuse container. Securely wrap partially filled or empty container in newspaper and discard in trash. **PRODUCT:** Partially filled bag may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER:** Do not reuse bag. Discard bag in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly after handling.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash thoroughly immediately after handling and before eating, smoking or using the toilet.

(reentry icon) Do not allow children or pets to contact treated area until dust has settled.

FIRST AID: IF SWALLOWED: Immediately telephone a poison control center, emergency treatment center, or a physician for advice. DO NOT make person vomit unless directed to do so by medical personnel. Give water or milk to drink and telephone for medical advice. Consult medical personnel before inducing vomiting. If medical advice cannot be obtained, then immediately take person and product container to the nearest medical emergency treatment center or hospital. **IF IN EYES:** Flush eyes with plenty of water in case of eye contact wash eyes immediately with fresh water for at least 15 minutes. Call a physician. Get medical attention if irritation persists. **For skin, wash with plenty of soap and water.** If inhaled, remove person from exposure area. **Note to Physicians:** Emergency Information call 1-800-454-2333. Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwater by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming plants, crops or weeds. Do not apply this product or allow it to drift to blooming plants, crops or weeds if bees are visiting the treatment area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

☐ Medical Information call 1-800-454-2333

Product Information call 1-800-225-2883

ORTHO®, SOLARIS® Registered Trademarks of Monsanto Company
SEVIN® is the registered trademark of Rhone-Poulenc for carbaryl insecticide.

©Monsanto Company _____

6 7 6

Manufactured for
The SOLARIS Group of
Monsanto Company
P.O. Box 5008
San Ramon CA 94583-0808
Form _____
~~Product 2845~~
EPA Reg. No. 239-1349
EPA Est 58996-MO-1^A, 769-GA-1^G
Superscript is first letter of lot number
Made in USA