ORTHO®

[MITICIDE]

Jul 12, 1971

NET CONTENTS_____
LIQUID

Active Ingredients	By Wt.
*Tetradifon	12.3%
Petroleum hydrocarbons	35.0%
Inert Ingredients	52.7%

*p-chlorophenyl 2,4,5-trichlorophenyl sulfone
Contains 1 pound Tedion per gallon
U. S. Pat. No. 2,812,281

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

[illegible] ORIGINAL CONTAINER. DO NOT PUT INTO [illegible] INTO FOOD OR DRINK CONTAINERS.

WASH AND DESTROY CONTAINER WHEN EMPTY.
NEVER RE-USE.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:
GROUND APPLICATION—Vegetable and Field Crops—[illegible] 250 gals.
Where Recommended for AIR APPLICATION—Field Crops—[illegible] gals.

APPLES, CRABAPPLES, PEARS, QUINCES: European Red Mite, Two-Spotted Spider Mite, McDaniel Mite, Canadensis Mite, Schoenii Mite, Carpini Mite—WEST COAST AREAS: 1 quart per 100 gals. water or 1¼ to 2½ gals. per acre in water to cover. **EASTERN AREAS:** [illegible] quart per 100 gals. water. (No more than [illegible] quarts diluted spray per acre.) For both Eastern and Western Areas, applications may be made up to day of harvest but make no more than 3 applications after petal fall. Do not feed pomace from treated apples to livestock.

CAUTION

KEEP OUT OF
REACH OF CHILDREN

CAUTION: [illegible] Avoid breathing vapor. Avoid contact with skin. Do [illegible] OPEN FLAME. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.
[illegible] toxic to shrimp and crabs. Keep out of [illegible] waters.

APRICOTS, CHERRIES, NECTARINES, PEACHES: European Red Mite, Two-Spotted Spider Mite, McDaniel Mite—WEST COAST AREAS: [illegible] **EASTERN AREAS:** [illegible] No more than 800 [illegible] Western [illegible]

CITRUS (Citron, Grapefruit, Limes, Oranges, Tangelos, Tangerines): Citrus Red Mite (Florida Purple Mite), Six-Spotted Mite, Texas Citrus Mite—Apply when mites first appear. **CALIFORNIA CITRUS AREAS:** [illegible] 2½ gals. per acre in water to thoroughly cover fruit and foliage. **FLORIDA CITRUS AREAS:** Use 1 to 2 [illegible] gallons. Apply sufficient gallonage to thoroughly [illegible] and foliage. **OTHER CITRUS AREAS**—Use 1 quart per [illegible] Apply sufficient gallonage to thoroughly cover [illegible] and foliage. Do not apply more than 1 application to the [illegible] per season.

CUCUMBERS: Two-Spotted Mites—Greenhouse—1 quart per 100 [illegible] **Field**—2 quarts per [illegible] water to cover. Spray [illegible] Apply when mites first appear and repeat as [illegible] apply more than 2 applications during growing [illegible]

GRAPES: Pacific Mite, Two-Spotted Spider Mite, Willamette Mite [illegible] Make no more than 3 appli[illegible] May be applied up to the [illegible]

HOPS: Two-Spotted Mite—[illegible] per acre in water to cover. [illegible] when mites first appear. [illegible] make more than 2 appli[illegible] petal fall. Do not apply within 14 days of harvest. Do not feed straw to livestock.

MELONS, PUMPKINS, WINTER SQUASH: Two-Spotted Spider Mite—2 quarts per acre in water to cover when mites first appear. Repeat as necessary. May be applied up to the day of harvest.

ORNAMENTALS (Greenhouse and Out-of-doors): Two-Spotted Spider Mite, Citrus Red Mite—Use 2 to 2½ pints per 100 gallons of water. Spray upper and lower leaf surfaces thoroughly when mites first appear and repeat as required. For Standard Rose Plants in the greenhouse, apply 30 to 35 gals. spray per [illegible] square feet of bench space, depending upon the size of the plants. Applications of this material may cause leaf drop on White Butterfly and Cinderella varieties of roses.

PEPPERMINT, SPEARMINT: Two-Spotted Mite—1¼ quarts per acre [illegible] water to cover. Apply when mites first appear. Repeat as re[illegible] make more than 2 applications per season. Do [illegible] days of cutting. Do not feed mint hay or spent mint hay to livestock.

PLUMS, PRUNES: European Red Mite, Two-Spotted Spider Mite, McDaniel Mite—WEST COAST AREAS—1 quart per 100 gals. water or 1¼ to 2½ gals. per acre in water to cover. **EASTERN AREAS**—½ [illegible] per 100 gals. water. For both Eastern and Western Areas, applications may be made up to day of harvest, [illegible] more than 3 applications during fruiting season.

STRAWBERRIES: Two-Spotted Spider Mites—1 gal. per acre in water to cover when mites first appear. During fruiting period, [illegible] interval less than 35 days. Do not [illegible] berries.

TOMATOES: Tropical Mite, Two-Spotted Spider Mite—2 quarts per acre in water to cover when mites first appear. Repeat as necessary [illegible] apply more than 3 applications during the [illegible] period. May be applied up to the day of harvest.

NOTE: [illegible] effective against hatching mite eggs and young nymphs. Kill of adult mites is not rapid and several days may elapse before full effect is noted. However, this material has [illegible] residual activity.

CONDITIONS OF SALE: 1. [illegible]