

VOLCK Supreme-Ethion Combination

ACCEPTED
 MAR 31 1976
 Under the Federal Insecticide, Fungicide, and Rodenticide Act
 as amended by the pesticide
 laws of 1972
 EPA Reg. No. 239-1444



ORTHO

(INSECTICIDE-MITICIDE)

Active Ingredients	By Wt.
Petroleum Oil	93.0
*Ethion	3.4
Inert Ingredients	3.6
*O,O,O',O' Tetraethyl S,S' methylene bisphosphorodithioate	

Contains 0.25 pound Ethion per gallon
 Petroleum Oil Classification: Unclassified
 Petroleum Oil Min. Unsulfonated Residue: 92%
 U.S. Pat. No. 2,873,228

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DIRECTED INTO OTHER CONTAINERS

DO NOT STORE AT TEMPERATURES BELOW 50°F

WHEN POURING IN FAMILY TRAPLE RINSE FOR EQUIVALENT AND OFFER FOR RECYCLING. DO NOT REUSE FOR OTHER PURPOSES. APPROVED BY EPA FOR RECYCLING IN A SAFE PLACE

NET CONTENTS LIQUID

Chevron Chemical Company
 Ortho Division, San Francisco, CA 94119
 Pat. No. 2,873,228 Made in U.S.A.
 Form 5042 P
 EPA Reg. No. 239-1444 AA

VOLCK Supreme-Ethion Combination is a contact and contact-feeding insecticide and miticide. It is effective against many insects and mites. Use in accordance with the label directions. Do not apply to food crops. Do not use on ornamental plants.

DIRECTIONS

ALMOND, NECTARINE, PEACH, PLUM, PRUNE San Jose Scale, Soft Scales, Overwintering Eggs of European Red Mite and Brown Almond Mite, Parlatoria (Olive) Scale. Use 1 to 1.5 gals. per 100 gals. water during the dormant period. Do not spray or flock in late dormant stages.

APPLES San Jose Scale, Soft Scales, Overwintering eggs of Rosy Apple Aphid and Green Apple Aphid, European Red Mite or Brown Almond Mite. Use 1 to 1.5 gals. per 100 gals. water in the dormant period. **Blister Mites** Use 2 gals. per 100 gals. water in the late dormant period. Do not use in areas where the most recent harvest was in the fall. Do not use on young trees.

STOP
 DO NOT TAKE INTERNALLY
 DO NOT INHALE
 DO NOT GET ON SKIN

KEEP OUT OF REACH OF CHILDREN

WARNING:

Caution: This product contains a highly flammable solvent. Do not use in enclosed areas. Do not use near open flames or other sources of ignition. Do not use in areas where smoking is prohibited. Do not use in areas where children are present. Do not use in areas where pets are present. Do not use in areas where food is stored. Do not use in areas where food is prepared. Do not use in areas where food is served. Do not use in areas where food is consumed. Do not use in areas where food is stored, prepared, or served. Do not use in areas where food is consumed. Do not use in areas where food is stored, prepared, or served. Do not use in areas where food is consumed.

PEAR San Jose Scale, Soft Scales, Overwintering Eggs of European Red Mite and Brown Almond Mite. Use 1 to 1.5 gals. per 100 gals. water in dormant period. **Pear Psylla** Use 1.5 gals. per 100 gals. water in dormant period. **Blister Mites** Use 2 gals. per 100 gals. water in dormant period. Do not use in areas where food is stored, prepared, or served.

PLUM, PRUNE, APRICOT, CHERRY Overwintering eggs of Mealy Plum Aphid, Leaf Curl Plum Aphid, European Red Mite and Brown Almond Mite. Use 1 to 1.5 gals. per 100 gals. water in the late dormant period.

MIXING DIRECTIONS With enough water to make to cover the plants and with a spray nozzle, add desired amount of the concentrate to the tank. Pump this mixture through the overflow and back into tank for about two minutes to insure proper emulsification. Fill tank with water, adding materials mentioned below, if desired, and mix well.

NOTE This product may be used in combination with ORTHO Copper 53 Fungicide. It may also be used in combination with ORTHO Pledge 1 Fungicide, ORTHO Fungicide, or other fungicides. Do not use in areas where food is stored, prepared, or served. Do not use in areas where food is consumed. Do not use in areas where food is stored, prepared, or served. Do not use in areas where food is consumed.

CONDITIONS OF SALE This product is sold as a commodity (Chevron) and is not intended for use as a pesticide. It is sold as a commodity and is not intended for use as a pesticide. It is sold as a commodity and is not intended for use as a pesticide. It is sold as a commodity and is not intended for use as a pesticide.

VOLCK[®] Supreme- Ethion Combination

ACC

MA

Under the
Fungicide,
as amended
registered
EPA Reg.

(INSECTICIDE-MITICIDE)

Active Ingredients	By Wt.
Petroleum Oil	93.0%
*Ethion	3.4%
Inert Ingredients	3.6%
*O,O,O',O'-Tetraethyl S,S'-methylene bisphosphorodithioate	

Contains 0.25 pound Ethion per gallon
Petroleum Oil Classification—Unclassified
Petroleum Oil Min. Unsulfonated Residue—92%
U. S. Pat. No. 2,873,228

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DO NOT STORE AT TEMPERATURES BELOW 32° F.

WHEN DRUM IS EMPTY, TRIPLE RINSE (OR EQUIVALENT) AND OFFER FOR RECYCLING, RECONDITIONING OR DISPOSAL IN APPROVED LANDFILL OR BURY IN A SAFE PLACE.

NET CONTENTS _____ LIQUID

Chevron Chemical Company

Ortho Division/San Francisco CA 94119 Richmond CA 94804
Product 2991 Made in U.S.A.
Form 5942-P
EPA Reg. No. 239-1444-AA

VOLCK Supreme-Ethion Combination is for use in the Pacific Coast dormant and late dormant periods or edible parts begin to form. Do not use in undiluted form.

DIRECTIONS

ALMOND, NECTARINE, PEACH: Soft Scales, Overwintering Eggs of Almond Mite, Parlatoria (Olive) Scale—Use 1 to 1½ gals. water during the dormant period in treated almond groves.

APPLES: San Jose Scale, Soft Scale, Apple Aphid and Green Apple Aphid—Use 1 to 1½ gals. water during dormant period. Blister Mites—Use 1 to 1½ gals. water during late dormant period. Do not use on Pratt, Melba, Wealthy, Williams Red and other varieties. Injury may result.



DO NOT T
DO NOT I
DO NOT C

KEEP OUT OF REA

WARNING:

May be absorbed through skin. Handle concentrate in a well-ventilated area. Do not breathe vapor or spray mist. Avoid contact with clothing. Wash thoroughly after handling. Do not use near feed or food.

Note to Physicians: Emergency treatment with Atropine is antidotal. 2 PAM is administered in conjunction with atropine. KEEP AWAY FROM HEAT OR OPEN FLAME.

This product is toxic to fish and wildlife in streams or ponds. Do not apply where runoff may occur. Do not apply when weather conditions favor runoff. Do not contaminate water by cleaning of equipment and containers and disposal of washwaters. Apply this product only to crop lands away from water supplies.

VOLCK[®] Dormant & Summer Oil



ORTHO

DORMANT AND GROWING SEASON INSECT SPRAY.

Controls Scale Insects, Mealybugs, White Flies, Red Spider Mites

Active Ingredient	By Wt.
Petroleum Oil	97%
Inert Ingredients	3%
(Pet. Oil—Min. U.R. 91%—Class Light-Med.)	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DIRECTIONS

Shake well before using. Use a forceful spray, wetting pests and all parts of plants thoroughly. Do not store diluted spray. Make new dilution for each use.

NET CONTENTS

LIQUID

Chevron Chemical Company

Ortho Division San Francisco, Calif. 94119

Richmond, Calif.


Lafayette, Calif.

Des Plaines, Ill.

Perth Amboy, N.J.

Atlanta, Ga.

Form 8354 EPA Reg. No. 239 2399 Product 710 Made in U.S.A.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

CAUTION: Harmful if swallowed. Avoid inhalation of vapors or spray mist. Wash hands after use. Do not use VOLCK Dormant & Summer Oil on foliage within 30 days before or after using sulfur. Avoid spraying when temperature is excessively high or when plants are suffering from drought. Oil sprays should not be applied when freezing or near-freezing temperatures may be expected. Do not use on Maidenhair Ferns, Cocos Palms, Sugar and Japanese Maples, Beech, Birch, Hickory, Walnut, Butternut or non-woody plants unless specified. VOLCK Dormant & Summer Oil may cause discoloration to Blue Spruce. Do not use in combination with or closely following treatment with ORTHO-CIDE Garden Fungicide.

Do not re-use container. Destroy when empty. Do not contaminate water by cleaning of equipment or disposal of wastes

FOR DORMANT USE—To Control Scale Insects, Red Spider Mites, Mites, and Mealybugs on Dormant Shrubs, Evergreens (note caution). Woody Plants, Fruit Trees (Apples, Plums, Pears, Apricots, Peaches, Cherries) and Shade Trees: 1/3 pint (5 1/3 ozs.) per gal. water.

GROWING SEASON—To Control Scale Insects, Red Spider Mites, Mites, Mealybugs and Whitefly Larvae on Citrus (Oranges, Lemons, Grapefruit, Tangerines, Limes), Fruit Trees (Apples, Plums, Pears, Apricots, Peaches, Cherries), Shade Trees, Evergreens, Camellias, Azaleas, Hardy Shrubs: 5 Tablespoonfuls (2 1/2 ozs.) per gal. water. This application is also effective in the removal of sooty mold fungus through the control of whitefly larvae.

COMBINATION WITH ORTHO Malathion 50 Insect Spray on Citrus, Avocado, Mangoes, Gardenias and Ixora for Control of Scale Insects, Red Spider Mites, Mealybugs, Whiteflies, Aphids, Thrips and for the removal of Sooty Mold Fungus through control of Whitefly Larvae—Use 1 Tablespoonful VOLCK Dormant & Summer Oil and 1 Tablespoonful ORTHO Malathion 50 Insect Spray for each gallon of diluted spray. Do not use on citrus during full bloom. Do not use on citrus or avocados within 7 days of harvest. Do not use on mangoes within 2 days of harvest.

OVICIDE: Helps to destroy Red Spider Mite Eggs, Oak Moth Eggs and Codling Moth Eggs.

AS A PLANT POLISH—To brighten leaves of hardy house plants: Use 3 Tablespoonfuls (1 1/2 ozs.) per gal. water.

GREENHOUSES: To Control Scale Insects, Red Spider Mites, Mites, Whitefly, Mealybugs on Hardy Shrubs, Gardenias, Camellias and Azaleas—2 Tablespoonfuls (1 oz.) per gal. water. Allow at least 60 days between oil sprays and limiting to 3 applications per year.

PALM TREES: Scale Insects, Mealybugs, Red Spider Mites—3 Tablespoonfuls (1 1/2 ozs.) per gal. water. Note caution.

AZALEAS, CAMELIAS: Scale Insects, Red Spider Mites, Mealybugs, Whitefly Larvae, Lace Bug Larvae—5 Tablespoonfuls (2 1/2 ozs.) per gal. water. Apply in spring after blooming and in fall prior to blooming.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions