

Chevron

# ORTHO Fly Killer D

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE FEDERAL AND STATE REGULATIONS**

**KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.**

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

**STORE IN COOL, DRY PLACE.**

Do not store diluted spray. Make new dilution for each use.

Do not use this product on livestock or domestic animals other than indicated on the label.

**AS A SPACE SPRAY for Control of Resistant and Nonresistant House Flies, Little House Flies, Mosquitoes, Gnats and Fruit Flies (Drosophila species) in and around Dairy Barns, Livestock Barns, Pig Pens, Poultry Houses, Feed Lots, Cattle Pens, Garbage Dumps, Outside Meat Packing Establishments (including Federally Inspected) pens, docks, ramps, disposal areas and Cider Mills:** Use 5 teaspoonfuls of ORTHO Fly Killer D in 1 gal. water (2 oz. in 2 1/2 gal. or 2 pts. to 40 gals.) and direct the spray throughout the fly infested area such as dairy and stock barns, spraying around and above animals but not directly at animals. Repeat as necessary. Do not use inside dwellings or in milk processing rooms. **For increased residual control,** concentrate spray application to walls, beams, rafters and around windows, doorways and other fly resting areas. Add 1/2 lb. sugar or 1/2 pt. Karo syrup when 40 gals. water is used for best results. **Fruit Flies (Drosophila sp.) in and around Food Processing Plants, Loading Docks, Cull Piles and Refuse Areas and Cider Mills:** Use 4 oz. in 2 1/2 gal. water (10 pts. to 100 gals.). Apply as a coarse spray to walls, floors, doorways, windows, refuse and cull piles. Apply every 5 to 7 days as long as necessary to maintain control. Do not apply to cull fruit or refuse piles to be fed to livestock. Avoid contamination of feeds, foodstuffs and food processing machinery. Do not apply when plants are in operation or when foods are present or exposed. Do not spray surfaces which will come into contact with foods. Cover food containers during spraying periods.

**ADULT MOSQUITO, GNAT AND HOUSEFLY CONTROL—RESIDENTIAL AREAS, FEED LOTS AND PASTURES INCLUDING DAIRY CATTLE, WOODLANDS, SWAMPS:** It is not necessary to avoid farm buildings and dairy barns. Time application for peak infestation and repeat as necessary. Do not apply to pastures within 400 yds. of any crop. **AIRCRAFT APPLICATIONS:** 1.6 to 2.0 oz. per acre (0.5 to .25 lb. technical Naled per acre). Apply in water to give 1 to 2 oz. diluted spray per acre. **MIST BLOWER APPLICATION:** 1 1/2 to 2 1/2 oz. in 100 gals. water. Calibrate equipment rate of travel and output to apply 1 to 2 1/2 lb. technical Naled per acre. Treat directly on vegetation where mosquitoes may be a problem, and vegetation around stagnant pools, marshes, swamps, and other areas where mosquitoes may be a problem. Do not apply to water.

**PENS, HOUSES, CAGES AND COOPS containing Poultry, Chickens, Turkeys and Pheasants: Mites, Ticks, Flies, Mosquitoes and Gnats—**1 pt. to 20 gals. water. Apply at rate of 1 gal. diluted spray per 500 sq. ft. Spray roosts, perches, walls of houses, nests and other areas for pest control. Birds may be present while spraying.

**AS A BAIT for Control of Resistant or Nonresistant House Flies in and around Dairy Barns, Livestock Barns, Stables, Pig Pens, Poultry Houses, Outside Meat Packing Establishments (including Federally Inspected) pens, docks, ramps, disposal areas, Animal Hospitals, Dog Kennels, Open Air Theatres, Food Processing Plants, Restaurants, Drive-Ins, and Where Flies Breed and Congregate:** **As a Dry Bait—**Put 1 lb. granulated sugar into a 1-gal. jar or a 1-gal. can. Add 1/2 oz. ORTHO Fly Killer D. Tighten lid and shake vigorously with rotating motion for two minutes. Scatter resulting bait on floors, window sills and loading docks in barns, kennels, animal hospitals, refuse pits, patios and where flies congregate. In food processing areas, use only when the facility is not in operation and food is not exposed. Use only in dust boxes where dust can be collected for removal and not for bait boxes prior to receiving food processing. Repeat as necessary. Unused bait will remain effective for two or three weeks. Keep in original container. **As a Wet Bait—**Use 1 lb. of sugar or 1/2 pt. Karo syrup or 1/2 pt. molasses in 2 1/2 gals. water in a sprinkling can. Add 2 Tbsp. of ORTHO Fly Killer D. Sprinkle on entry ways, windows, loading docks and walkways in stock barns, food processing plants, kennels, poultry houses, and refuse areas where flies congregate and feed as necessary. Prepare new bait each time.

**ROACHES (All Species of Resistant and Nonresistant Cockroaches, including German, Brownbanded, Oriental and American Roaches of Houses, Apartments, Hotels, Motels, Restaurants, Institutions and Other Buildings: FOR USE BY PEST CONTROL OPERATOR)** 1 gal. water. Apply as a fine stream or coarse spray into cracks, crevices, and where insects can be contacted.

**POULTRY, CHICKENS, TURKEYS, PHEASANTS: Poultry Lice (Body Louse, Fluff Louse, Shaft Louse), Chicken Mites—**1 pt. to 20 gal. except head with light mist. Use at approximate rate of 1 gal. diluted spray per 100 birds. Concentrate application under wings and around tail area. Do not use on age and turkeys under 3 months. Repeat applications as necessary to maintain control.

**BROWN DOG TICK on DOGS—**Prepare a dip using 1 oz. to 2 gals. water. Water dips before dipping to prevent dripping of dip. Do not allow dip to get in eyes. Dip for 10 to 15 minutes. Wipe off excess dip. Wash dog's face and ears. Do not use on cats. **In Kennels—**Do not use this dilution on animals. Apply as a directed insect wet spray, being especially careful to hit cracks and crevices in floor, walls and ceiling. Also spray out doorways. Repeat if tick activity is observed.

**DANGER:** May be fatal if swallowed, inhaled or absorbed through the skin. Corrosive—causes eye damage and skin irritation. Do not get in eyes or on skin. Vapors or spray mist. Wear goggles or face shield and rubber gloves when handling. In case of skin contact, immediately remove contaminated clothing and wash with plenty of water and get medical attention.

**Note to Physicians:** Emergency Information—call (415) 233-5737. Atropine is antidotal. 2 PAM is also antidotal and may be administered in conjunction with atropine. Avoid contamination of food, foodstuffs and drinking water. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** May corrode metal. Wash thoroughly after use. When container is empty, immediately wash thoroughly or destroy. Never reuse.

This product is toxic to fish, shrimp, crab, birds and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crab are recommended on this label. Do not apply where these organisms are reared. Do not apply where run-off is likely to occur. Do not apply when weather or treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and directed hereon. Chevron neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guaranty or implied, concerning this material. 2. Certain and/or foreseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include damage from application, lack of complete control, and damage caused by theft. Such risks occur even though the product is representative of the user's label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under applicable laws) arising from this material.

## Chevron Chemical Company

Ortho Division, San Francisco CA 94119 Richmond CA 94804

Product 3021 Made in U.S.A.

Form 60/22 82

EPA Reg. No. 239-1466-ZA

BA

[239-1466]

Chevron



F

5000  
Cond  
Dros

**ROACHES (All Species of Resistant and Nonresistant Cockroaches, including German, Brownbanded, Oriental and American Roaches) Inside and Outside of Houses, Apartments, Hotels, Motels, Restaurants, Institutions and Other Buildings: FOR USE BY PEST CONTROL OPERATORS ONLY**—Use 3 fl. oz. in 1 gal. water. Apply as a fine stream or coarse spray into cracks, crevices, and where insects can be contacted.

**POULTRY, CHICKENS, TURKEYS, PHEASANTS: Poultry Lice (Body Louse, Fluff Louse, Shaft Louse), Chicken Mites**—1 pt. to 20 gals. water. Spray entire bird except head with light mist. Use at approximate rate of 1 gal. diluted spray per 100 birds. Concentrate application under wings and around tail area. Do not treat chickens under 5 weeks of age and turkeys under 3 months. Repeat applications as necessary to maintain control.

**BROWN DOG TICK on DOGS**—Prepare a dip using 1 oz. to 2 gals. water. Water dogs before dipping to prevent dripping of dip. Do not allow dip to contact eyes of dog. One drop of glycerine in each eye will prevent irritation from splashing. Wear rubber gloves and work dip well into hair of dog. Use glycerine to wash inside of ears with dip. Do not allow dog to remain in dip for more than 30 seconds. Dry immediately out of draft. Do not use on cats. **In Kennels—Do not use this dilution on animals**—Use 5 oz. to 2 gals. water. Apply as a directed coarse wet spray, being especially careful to hit all cracks and crevices in floor, walls and ceiling. Avoid contact with humans. Repeat treatment in 7 days if control tick activity is observed.

**DANGER:** May be fatal if swallowed, inhaled or absorbed through the skin. Corrosive, causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Avoid breathing vapors or spray mist. Wear goggles or face shield and rubber gloves when handling. In case of skin contact, immediately remove contaminated clothing and flush skin with plenty of water. For eyes, immediately flush with plenty of water and get medical attention.

**Note to Physicians:** Emergency Information—call (415) 233-8737. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Avoid contamination of feed, feedstuffs and drinking water. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. May corrode metal spray equipment. Wash equipment thoroughly after use. When container is empty, immediately wash thoroughly and destroy. Never reuse.

This product is toxic to fish, shrimp, crab, birds and other wildlife. Keep out of lakes, streams, ponds, sea marshes and streams. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where there are important riparian areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.  
2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage from application, lack of complete control, and damage caused by drift. Such risks do not even though the product is reasonably fit for the use stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability, except those assumed by Chevron under 1. above, arising from handling, storage, and use of this material.

### Chevron Chemical Company

Ortho Division, San Francisco CA 94119 Richmond CA 94804  
Product 3021 Made in U.S.A.  
Form 6022 B2

EPA Reg. No. 239-1466-ZA

BA [239-1466]

Chevron



# ORTHO

## Fly Killer D

ACCEPTED

AUG 15 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 239-1466

... and American Roaches). Inside and Outside  
CONTROL OPERATORS ONLY--Use 3 fl. oz. in

... Mites--1 pt. to 20 gals. water. Spray entire bird  
... around tail area. Do not treat chickens under 6 weeks

... Do not allow dip to contact eyes of dog. One drop  
... to wash inside of ears with dip. Do not allow dog to  
... dilution on animals--Use 3 oz. to 2 gals. water  
... outside always. Repeat treatment in 7 days if continual

... do not get in eyes, on skin, or on clothing. Avoid breathing  
... contaminated clothing and flush skin with plenty of water.

... administered in conjunction with atropine.  
... May corrode metal spray equipment. Wash equipment

... Shrimp and crab may be killed at application rates  
... not apply when weather conditions favor drift from areas

... description on the label and is reasonably fit for use as  
... MERCHANTABILITY, guarantee or representation, express

... of chemicals. Such risks include, but are not limited to,  
... reasonably fit for the uses stated herein and even though  
... per 1. However, risks may vary from handling, storage, and use of

... contains DIBROMO Naled Insecticide  
Controls Flies, Mosquitoes, Gnats, Brown Dog Ticks,  
Drosophila, Poultry Lice and Mites