


ORTHO[®] Sulfur 10-50 Dust



ORTHO



(FUNGICIDE)

Active Ingredients	By Wt.
Captan*	10%
Sulfur	50%
Inert Ingredients	40%

* N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL, DRY PLACE

PROTECT FROM EXCESSIVE HEAT

BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

This product is formulated for application by ground or air equipment.

CELERY: Pink Rot, Damp-Off: (Field)—20 lbs. per acre. (Seed Bed)—35 to 40 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

GRAPES: Botrytis Rot (Bunch Rot), Powdery Mildew, Silver Mite, Storage Rots—40 to 50 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically making up to 3 cover applications before bunches close.

PEACHES: Brown Rot, Storage Rots, Powdery Mildew, Silver Mite, Peach Scab—50 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, California 94119

Richmond, California Fresno, California

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 4294-B2 Product 2130 Made in U.S.A.

EPA Reg. No. 239-1470-AA

PLUMS: Brown Rot, Storage Rots, Powdery Mildew, Silver Mite—45 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

POTATOES: Early and Late Blight—Begin applications when plants are 2 to 4 inches high and repeat at 7 to 10-day intervals as long as disease threatens. Apply 25 to 30 lbs. per acre.

ROSES (Field): Rust, Powdery Mildew—35 to 40 lbs. per acre. Apply to give even coverage of upper and lower leaf surfaces at first sign of disease. Repeat at regular intervals as necessary to maintain control.

SOIL TREATMENT: Pre-planting for Pink Rot, Damp-Off on Vegetable Field Crops—33 lbs. per acre. Dust or broadcast, then dig in first few inches of soil surface.

SPINACH (Seedling only): Downy Mildew, Powdery Mildew

—35 to 40 lbs. per acre. Apply when all seeds have germinated. Repeat at 7 to 10-day intervals as long as diseases are active.

SQUASH (California Only): Powdery Mildew, Downy Mildew, Angular Leaf Spot, Anthracnose—Apply 30 lbs. per acre on young plants increasing to 35 lbs. per acre on mature plants. Apply at first sign of disease. Repeat at 5 to 10 day intervals until danger of infection is past. Do not apply to squash varieties susceptible to sulfur injury.

STRAWBERRIES: Botrytis Rot, Powdery Mildew—30 lbs. per acre immediately after picking. Repeat applications after picking as necessary.

TOMATOES (Established Plants): Early Blight, Late Blight, Anthracnose, Russet Mite—On young plants, apply 25 to 30 lbs. per acre, increasing to 35 to 40 lbs. per acre on mature plants. Apply at first sign of insects and diseases. Repeat at 5 to 7-day intervals as necessary for control. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

COMPATIBILITY: Do not use this product on foliage closely following oil sprays.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Avoid inhalation of dust. Avoid contact with skin. Do not store or transport near feed or food. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. **Do not apply under conditions involving possible drift to food, forage, or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.