

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly after handling. ~~Avoid contamination of food, feed, feeding troughs and water receptacles.~~

STATEMENT OF PRACTICAL TREATMENT

If swallowed, DO NOT make person vomit unless directed to do so by a doctor. Immediately take person and container of product to, or if not possible, call a doctor or an emergency hospital. In case of eye contact, immediately flush eyes with fresh water for 15 minutes. Get medical attention if irritation persists. For skin, wash with plenty of soap and water. **Note to Physicians:** Emergency Information - call (415) 233-3737. Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water ~~except for rice uses. Discharge from the fields may kill aquatic and estuarine invertebrates.~~ Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

REENTRY STATEMENT

Keep all unprotected persons out of operating areas or vicinity where there may be drift. Do not enter treated areas without protective clothing until dusts have settled. Certain states may require more restrictive reentry intervals. Consult your State Department of Agriculture for further information.

Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas about to be treated.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE

Keep pesticide in original container. Do not store or transport near feed or food. For help with any spill, leak, fire or exposure involving this material, call day or night (415) 233-3737.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Do not reuse container. Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

May be applied up to harvest unless otherwise noted. This product is formulated for application by ground or air equipment.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

ALFALFA, CLOVERS, COWPEAS, PEANUTS, SOYBEANS: Blister Beetles, Mexican Bean Beetle 5 to 10 lbs. per acre. Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetles, Green Cloverworm, Japanese Beetle, Leafhoppers, Threecornered Alfalfa Hopper, Thrips, Velvetbean Caterpillar—10 lbs. per acre. Armyworms, Corn Earworm, Stink Bugs, Webworms—10 to 15 lbs. per acre. Some leaf injury may occur if applications are made when tender foliage is present. Do not apply to Alfalfa and Cowpeas within 3 days of harvest.

BEANS: Mexican Bean Beetle—5 lbs. per acre. Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Velvet Bean Caterpillar, Flea Beetles, Japanese Beetle—10 lbs. per acre. Corn Earworm, Armyworms, Tarnished Plant Bug, Stink Bugs—10 to 15 lbs. per acre. In California for Corn Earworm, Lima Bean Pod Borer, Lygus and Stink Bugs—20 lbs. per acre.

BLUEBERRIES: Blueberry Maggot, Cherry and Cranberry Fruitworms—15 to 20 lbs. per acre. Apply 3 weeks before harvest and repeat 10 days later or as necessary.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: Flea Beetles, Harlequin Bug—5 to 10 lbs. per acre. Armyworms, Corn Earworm, Imported Cabbageworm, Young Cabbage Looper—10 to 20 lbs. per acre. Where Cabbage Looper is the principal problem, use an alternate pesticide that is specifically recommended. Do not apply within 3 days of harvest.

CORN: Corn Earworm (Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations). Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetle, Leafhoppers, Flea Beetles—15 to 20 lbs. per acre. For insects attacking silks and ears apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. May be applied up to harvest.

ACCEPTED

17 JAN 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 239-1513



ORTHO

SEVIN[®] 10
Dust

(INSECTICIDE)

Kills: Beetles, Bugs, Leafhoppers and Worms listed on vegetables. May be used up to harvest unless otherwise noted.

Kills Fleas & Ticks on dogs and cats.

On Vegetables

On Dogs and Cats

Stink Bugs, Tomato Hornworms, Brown Dog Tick
Flea Beetle, Bean Leafhopper, Fleas
Corn Earworm

Active Ingredient

Carbaryl (1-naphthyl N-methylcarbamate)

Inert Ingredients

SEVIN[®]—Reg. TM of Union Carbide Corp. for Carbaryl Insecticide.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.

Chevron Chemical Company ©1983

Ortho Consumer Products Division

San Francisco CA 94128-7144

Richmond CA 94804-0036

Product 2903

Made in U.S.A.

R Form 10345-C

EPA Est. 239-CA-9

EPA Reg. No. 239-1513-AA

NET WEIGHT 50 POUNDS

BEST COPY AVAILABLE

ORTHO SEVIN[®] 10 Dust (INSECTICIDE)

Beetles, Bugs, Leafhoppers and Worms as
red on vegetables. May be used up to harvest
less otherwise noted.

Beetles & Ticks on dogs and cats.

On Vegetables	On Dogs and Cats
Stink Bugs, Tomato Hornworms, Flea Beetle, Bean Leafhopper, Corn Earworm	Brown Dog Ticks, Fleas

Active Ingredient
Carbaryl (1-naphthyl N-methylcarbamate) 10%
Inert Ingredients 90%
EPA Reg. No. 239-1513-AA

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.

Chevron Chemical Company © 1962
The Consumer Products Division
San Francisco, CA 94120-7144
Product 2903
Form 10345-C
EPA Reg. No. 239-1513-AA

NET WEIGHT 50 POUNDS

COTTON: Early Season—Thrips, Cotton Fleahopper—8 to 15 lbs. per acre as soon as insects appear. Mid and Late Season—Boll Weevil, Bollworm, Pink Bollworm, Cotton Leafworm, Lygus Bugs, Cotton Leaf Perforator—20 lbs. per acre. Apply when insects appear and repeat at 5 to 7 day intervals. Aphid populations will usually be suppressed when this insecticide is used on a regular schedule of treatments. Do not apply within 7 days of harvest.

CUCUMBERS, MELONS, PUMPKIN, SQUASH: Pickleworm, Melonworm—5 to 10 lbs. per acre. Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs—10 lbs. per acre. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use SEVIN 10 Dust on watermelons in Florida.

PEACHES: Filbert Aphid, Filbert Leafroller—40 lbs. per acre. Apply when Leafroller eggs are hatching. Filbertworm—40 lbs. per acre. Apply when adult moths first appear and repeat 3 to 4 weeks later. Do not apply within 60 days of harvest.

GRAPES: Grape Leaf Folder, Grape Leafhopper, Grape Leaf Skeletonizer—10 to 20 lbs. per acre. Apply just before first broad leaf folder larvae emerge from rolls and as needed for leafhoppers. Grape Berry Moth, Japanese Beetle, June Beetles, Redbanded Leafroller—20 lbs. per acre.

GRASSHOPPERS: Alfalfa, Clovers, Cowpeas, Peanuts, Soybeans, Rice, Forage Grasses, Pasture, Milo and Grain Sorghum, Sugar Beets—5 to 15 lbs. per acre. Apply 5 to 10 lbs. for Nymphs on small plants or sparse vegetation in wasteland, rangeland, ditch banks and borders. Use 10 to 15 lbs. when grasshoppers are mature or material is applied to crops requiring greater coverage. Do not apply to Rice or Sugar Beets within 14 days of harvest, and to Sorghum Grain within 21 days of harvest.

LETTUCE, CARROTS: Flea Beetles, Harlequin Bug, Leafhoppers—5 to 10 lbs. per acre. Six-spotted Leafhopper—10 to 15 lbs. in a 5 to 7 day schedule. Armyworms, Imported Cabbage worm, Corn Earworm, Spittlebugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs—10 to 20 lbs. per acre. For best results in the control of Corn Earworm after lettuce heads begin to form use 20 lbs. in a 5 to 7 day schedule. May be used up to harvest on carrots. Do not use on Head Lettuce within 3 days of harvest and to Leaf Lettuce within 14 days.

PEACHES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Catfacing insects, Periodical Cicada, Codling Moth—35 to 40 lbs. per acre. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. May be applied the day before harvest.

PLUMS, PRUNES: Fruit Tree Leafroller, Peach Twig Borer, Plum Curculio, Redbanded Leafroller, Codling Moth, Eastern Tent Caterpillar, Orange Tortrix, Western Tussock Moth—40 to 50 lbs. per acre. Apply when insects appear and repeat as necessary. Make full coverage application. Do not apply within 1 day of harvest.

POTATO, EGGPLANT, PEPPER: Colorado Potato Beetle, Flea Beetles, Leafhoppers—5 to 10 lbs. per acre. Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus Bugs, Stink Bugs, Lace Bugs, Tarnished Plant Bugs—10 to 20 lbs. per acre.

RICE, FORAGE GRASSES, PASTURE: Armyworms, Stink Bugs, Thrips—10 to 15 lbs. per acre. For optimum thrip control in grasses grown for seed, direct dust into boot. Do not apply to Rice within 14 days of harvest. May be applied up to harvest on Pastures and Forage Grasses. Do not use on rice fields in which crayfish and/or catfish farming is included in the cultural practice.

SORGHUMS (Including Milo Grain Sorghum): Armyworms, Corn Earworm, Stink Bugs, Webworms—10 to 20 lbs. per acre. Sorghum Midge—15 lbs. per acre. Dust into forming heads. Treatment for sorghum midge control should be made 3 to 4 days after heads have emerged from boot. Do not apply to Sorghum Grain within 21 days of harvest.

STRAWBERRIES: Meadow Spittlebug, Strawberry Leafroller, Strawberry Weevil—10 to 20 lbs. per acre. Do not apply within 1 day of harvest.

TOBACCO: Hand Dusting for Budworms and Tobacco Flea Beetles—4 to 6 lbs. per acre when hand dust applications are made on individual plants. Heavier application may cause plant injury. Rotary dusters are recommended. Shaker cans and cloth bags usually result in excessive applications. Broadcast Application for Hornworms, Budworms, Japanese Beetles, Grasshoppers, Flea Beetles—15 to 25 lbs. per acre depending on size of plants. Use highest rate only on large plants. Full coverage is essential. Begin treatment when worms are small and repeat as needed. Heavier applications may cause plant injury. If late applications are necessary, treat immediately after pruning. Do not apply within 5 days of pruning or 15 days of cutting. Avoid getting wet with plant juices when cutting the crop.

TOMATO: Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer—20 to 40 lbs. per acre.

DOGS, CATS: Ticks, Fleas—Dust lightly over the animal and rub thoroughly into the skin. Apply also to the feet, between the toes and on the legs. Apply at weekly intervals if needed. Avoid treatment of kittens and puppies less than 4 weeks of age. NOTE: Do not allow treated pets back on carpets or rugs in the house as SEVIN 10 Dust may stain or discolor rugs and wall-to-wall carpets. Dust lightly in and around kennels and dog houses by means of a shaker can, hand or power duster. Force dust into cracks and crevices. Repeat as necessary, but not more often than once weekly. DO NOT USE IN THE HOME: DO NOT APPLY TO RUGS AND CARPETS AS STAINING MAY OCCUR. DO NOT USE ON PREGNANT DOGS.

Note: Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

BEST COPY AVAILABLE

ORTHO SEVIN 10 Dust
EPA Reg. No. 239-1513

COPY A

Apply dust at first signs of insects. Repeat as necessary. For best results, dust in early morning or late evening when temperature is low and air is still. Apply dust so that all parts of plants are contacted especially upper and lower leaf surfaces. May be applied up to harvest unless otherwise noted. ✓

COPY B

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

DISPOSAL: **PRODUCT**-Partially filled bag may be disposed of by securely wrapping original container in several layers of newspaper ✓ and discard in trash. **CONTAINER**-Do not reuse bag. Discard bag in trash.

COPY C

FOR HOME USE ONLY ✓

COPY D

Manufactured for

The SOLARIS Group
A Division of Monsanto Company
P. O. Box 5008
San Ramon CA 94583-0808 ✓
Product No. 2903
Form RS10345-D
EPA Reg. No. 239-1513
EPA Est. 62499-MO-1

NET WEIGHT 4 POUNDS