

PM19

239-1513

1/7

TO OPEN-CUT HERE
[COM] OUTDOOR USE ONLY

ORTHO® SEVIN® 10 DUST
[ABN: ORTHO® SEVIN® brand carbaryl insecticide 10 Dust]

Kills: Beetles, Bugs, Leafhoppers and Worms as Listed On Vegetables.

May be Used up to Harvest Unless Otherwise Noted.

KILLS FLEAS & TICKS ON DOGS AND CATS ~~AND IN DOGHOUSES~~

~~KILLS GARDEN PESTS: Armyworms, Cutworms, Squash Bugs, Tomato Hornworms, Certain Beetles And Other Garden Pests As Listed~~

Active Ingredient

Carbaryl (1-naphthyl N-methylcarbamate).....10%

Inert Ingredients.....90%

KEEP OUT OF REACH OF CHILDREN


CAUTION

See back panel for additional precautionary statements ~~on back of bag.~~

~~Specially formulated for residential use.~~

~~For home use only~~

NET WT 4 LBS / 1.81 kg



96 DEC 31 AM 1:10

RECD ... 270PD1

2/7

TO OPEN CUT HERE

READ ENTIRE LABEL BEFORE EACH USE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

VEGETABLES & FRUITS

PLANTS	INSECTS/PESTS	WHEN TO HARVEST
BEANS	Mexican Bean Beetles, Leafhoppers, Bean Leaf Beetles, Cucumber Beetles, Western Bean Cutworms, Velvet Bean Caterpillars, Flea Beetles, Japanese Beetles, Corn Earworms, Armyworms, Tarnished Plant Bugs, Stink Bugs	May be applied up to day of harvest.
BLUEBERRIES	Blueberry Maggots, Cherry and Cranberry Fruitworms	May be applied up to day of harvest.
CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER KOHLRABI	Flea Beetles, Harlequin Bugs, Armyworms, Imported Cabbageworms, Corn Earworms	Do not apply within 3 days of harvest.
CORN	Corn Earworms, Corn Rootworm Adults, European Corn Borers, Fall Armyworms, Sap Beetles, Japanese Beetles, Leafhoppers, Flea Beetles	May be applied up to day of harvest.
CUCUMBERS, MELONS, PUMPKINS, SQUASH	Pickleworms, Melonworms, Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs	May be applied up to day of harvest.
Do not use this product on watermelons in Florida.		
LETTUCE, CARROTS	Flea Beetles, Harlequin Bugs, Leafhoppers, Sixspotted Leafhoppers, Armyworms, Imported Cabbageworms, Corn Earworms, Spittlebugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs	May be applied up to day of harvest to carrots. Do not apply to head lettuce within 3 days of harvest or to leaf lettuce within 14 days of harvest.
PEACHES	Oriental Fruit Moths, Peach Twig Borers, Plum Curculios, Catfacing Insects, Periodical Cicadas, Codling Moths	Do not apply within 1 day of harvest.

PLUMS, PRUNES	Fruit Tree Leafrollers, Peach Twig Borers, Plum Curculius, Red-Banded Leafrollers, Codling Moths, Eastern Tent Caterpillars, Orange Tortrix, Western Tussock Moths	Do not apply within 1 day of harvest.
POTATOES, EGGPLANT, PEPPERS	Colorado Potato Beetles, Flea Beetles, Leafhoppers, Tomato Fruitworms, Fall Armyworms, Tomato Hornworms, European Corn Borers, Lygus Bugs, Stink Bugs, Lace Bugs, Tarnished Plant Bugs	May be applied up to day of harvest.
STRAWBERRIES	Meadow Spittlebugs, Strawberry Leafrollers, Strawberry Weevils	Do not apply within 1 day of harvest.
TOMATOES	Flea Beetles, Leafhoppers, Tomato Fruitworms, Fall Armyworms, Tomato Hornworms, European Corn Borers	May be applied up to day of harvest.

HOW TO APPLY

- Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces.
- Apply when air is calm to avoid drift and contact with eyes and skin.
- Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.
- For plants above chest height, consider using a liquid carbaryl product.

WHEN TO APPLY

- Apply dust at first signs of insects/pests.
- For best results, Dust in early morning or late evening when temperature is low and air is still.
- Repeat/Reapply as necessary, but not more often than once weekly.
- May be applied up to harvest unless otherwise noted.
- **BLUEBERRIES:** Apply 3 weeks before harvest and repeat/reapply 10 days later or as necessary.
- **CORN:** For insects attacking silks and ears apply when first silks appear and continue until silks begin to dry. Three or more applications may be required depending on severity of infestation.
- **CUCUMBERS, MELONS, PUMPKINS, SQUASH:** Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.
- **PEACHES:** Apply at petal fall and every 10 to 14 days thereafter until control is achieved.

BEANS: Mexican Bean Beetle, Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Velvet Bean Caterpillar, Flea Beetles, Japanese Beetle, Corn Earworm, Armyworms, Tarnished Plant Bug, Stink Bugs.

BLUEBERRIES: Blueberry Maggot, Cherry and Cranberry Fruitworms Apply 3 weeks before harvest and repeat 10 days later or as necessary.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: Flea Beetles, Harlequin Bug, Armyworms, Corn Earworm, Imported Cabbageworm Do not apply within 3 days of harvest.

CORN: Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetle, Leaf hoppers, Flea Beetles For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. May be applied up to harvest.

CUCUMBERS, MELONS, PUMPKIN, SQUASH: Pickleworm, Melonworm, Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use this product on watermelons in Florida.

LETTUCE, CARROTS: Flea Beetles, Harlequin Bug, Leafhoppers, Six spotted Leafhopper, Armyworms, Imported Cabbageworm, Corn Earworm, Spittlebugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs. For best results in the control of Corn Earworm after lettuce heads begin to form use in a 5 to 7 day schedule. May be used up to harvest on carrots. Do not use on Head Lettuce within 3 days of harvest and on Leaf Lettuce within 14 days.

PEACHES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Catfacing Insects, Periodical Cicada, Codling Moth. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. May be applied the day before harvest.

PLUMS, PRUNES: Fruit Tree Leafroller, Peach Twig Borer, Plum Curculio, Red Banded Leafroller, Codling Moth, Eastern Tent Caterpillar, Orange Tortrix, Western Tussock Moth. Apply when insects appear and repeat as necessary. Make full coverage application. Do not apply within 1 day of harvest.

POTATO, EGGPLANT, PEPPER: Colorado Potato Beetle, Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus Bugs, Stink bugs, Lace Bugs, Tarnished Plant Bugs.

STRAWBERRIES: Meadow Spittlebug, Strawberry Leafroller, Strawberry Weevil. Do not apply within 1 day of harvest.

TOMATO: Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer.

DOGS AND CATS

To control Ticks and Fleas: Use product only outside or in a well ventilated area. Wearing household latex or rubber gloves, lightly dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. This product should not be applied by children. Dust lightly over the animal and rub thoroughly into the skin. Apply also to the feet between the toes and on the legs. Repeat/Reapply as necessary, but not more often than once weekly. Apply at weekly intervals if needed. **DO NOT USE ON PREGNANT DOGS.** Avoid treatment of Do not use on kittens and on puppies under 12 4-weeks of age. **NOTE:** Do not allow treated pets back on carpets or rugs in the house as this product may stain or discolor rugs and wall-to-wall carpets. Dust bedding and doghouse lightly in and around kennels and hog-houses by means of a using a shaker can or other dust applicator hand duster. Force dust into cracks and crevices. Repeat/Reapply as necessary, but not more often than once weekly. **DO NOT USE IN THE HOME: DO NOT APPLY TO RUGS AND CARPETS AS STAINING MAY OCCUR.**

STORAGE AND DISPOSAL

STORAGE: Store in original container in a safe place. Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of food and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. **DISPOSAL:** Do not reuse container. Securely wrap partially filled or empty container in newspaper and discard in trash. **PRODUCT:** Partially filled bag may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER:** Do not reuse bag. Discard bag in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly after handling.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash thoroughly immediately after handling and before eating, smoking or using the toilet.

(reentry icon) Do not allow children or pets to contact treated area until dust has settled.

FIRST AID: IF SWALLOWED: Immediately telephone a poison control center, emergency treatment center or a physician for advice. DO NOT make person vomit unless directed to do so by medical personnel or a doctor. If medical advice cannot be obtained, then immediately take person and product container to an emergency treatment center or hospital or product to, or if not possible, call a doctor or an emergency hospital. **IF IN EYES:** flush eyes with plenty of water. In case of eye contact, immediately flush eyes with fresh water for 15 minutes. Call a physician. Get medical attention if irritation persists. For skin, wash with plenty of soap and water. **IF INHALED:** remove person from exposure area. **Note to Physicians:** Emergency Information call 1-800-454-2333. Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to ~~any body of water~~. Do not contaminate water ~~when disposing of equipment washwaters by cleaning of equipment or disposal of wastes~~. This product is highly toxic to bees exposed to direct treatment or residues on blooming ~~plants crops or weeds~~. Do not apply this product or allow it to drift to blooming ~~plants crops or weeds~~ if bees are visiting the treatment area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Medical Information call 1-800-454-2333

Product Information call 1-800-225-2883

ORTHO® ~~SOLARIS®~~ Registered-Trademarks of Monsanto Company
SEVIN® is the registered trademark of Rhone-Poulenc for carbaryl insecticide.

©Monsanto Company _____

Manufactured for

The SOLARIS Group of

Monsanto Company

P.O. Box 5008

San Ramon CA 94583-0808

Form _____

Product 2003

EPA Reg. No. 239-1513

EPA Est. 58996-MO-1^A, 769-GA-1^G

Superscript is first letter of lot number

Made in USA