



CHEVRON[®]
LAWN DISEASE
control

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

CHEVRON Lawn Disease Control is a wettable powder formulation to be used as a spray through the ORTHO Lawn Sprayer or Sprinkling Can and is designed to control common grass diseases such as Brown Patch (*Helminthosporium solani*) and certain other diseases including Leaf Spot (*Helminthosporium*), Cottony Blight (*Pythium*), Dollar Spot (*Sclerotinia*), Fading Out (*Cervularia*), and Gray Leaf Spot (*Cercospora*), particularly on St. Augustine, Bermuda, Bluegrass, Zoysia, Rye and Dichondra lawns. Do not store diluted spray. Make new dilution for each use.

DIRECTIONS

To obtain maximum results, it is important that the lawn be properly managed before and after treatment with CHEVRON Lawn Disease Control. Use the following Management Schedule as guide.

MANAGEMENT SCHEDULE:

1. One day before treatment: Mow lawn. Irrigate if necessary.
2. Day of treatment: After morning dew is off lawn, apply CHEVRON Lawn Disease Control as directed (see below).
3. First and second day following treatment: Do not mow lawn. Do not irrigate.
4. Repeat this schedule prior to retreatment according to preventive or curative programs outlined below.

APPLICATION RATE: Use ORTHO Lawn Sprayer or Sprinkling Can to apply treatments. Add the required amount of CHEVRON Lawn Disease Control to the Sprayer jar and add a small amount of water. Mix until a thin paste is formed; add water to bring to desired level and stir. Apply spray evenly. Fifteen gallons treats 1000 square feet. **AGITATE JAR WHILE SPRAYING.**


Amount of CHEVRON Lawn Disease Control to Use	Add Water to Make	To Treat
PREVENTIVE		
5 level tablespoons (1 2 oz.)	3 gallons	200 sq. ft.
25 level tablespoons (6 oz.)	15 gallons	1000 sq. ft.
CURATIVE		
5 level tablespoons (1 2 oz.)	3 gallons	100 sq. ft.
25 level tablespoons (6 oz.)	15 gallons	500 sq. ft.

PREVENTIVE PROGRAM: Apply in early fall. Successive applications may be necessary if warm weather persists during fall and winter. If necessary, treat every 30 to 35 days through February, depending upon temperature and environmental conditions. Application in late February may be necessary if ideal conditions for Brown Patch persist. After first spring cutting, repeat for control of such diseases as Gray Leaf Spot, Melting Out, Fading Out. Additional applications may be necessary in spring and summer if diseases persist.

CURATIVE PROGRAM: Use 6 ounces of CHEVRON Lawn Disease Control in ORTHO Lawn Sprayer and fill to 15 gallon mark and apply this to 500 sq. ft. of lawn area. Follow the Management Schedule and continue until disease is halted. Re-sod and top dress with soil areas destroyed by disease. Continue treatments using Preventive Program.

SPECIAL COMBINATION SPRAY: To control lawn insects such as white grubs, wireworms, sod webworms and ants, use ORTHO KLOR 74 Chlordane Spray at 1 Tablespoonful (1 2 oz.) per 1 gal. of water to cover 200 sq. ft. of lawn area with CHEVRON Lawn Disease Control at recommended rate.

COMMERCIAL APPLICATION: For use in tank type sprayers, use 4 lbs. of CHEVRON Lawn Disease Control in each 50 gallons of water to spray 10,000 square feet.



CHEVRON[®]
LAWN
DISEASE
control



CHEVRON[®]
LAWN DISEASE
control



CHEVRON[®]
LAWN
DISEASE
control

controls **BROWN PATCH**



GRAY LEAF SPOT
DOLLAR SPOT
FADING OUT
COTTONY BLIGHT
LEAF SPOT

Also
Controls
DICHONDRA
Diseases



CHEVRON[®]
LAWN DISEASE
control

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

CHEVRON Lawn Disease Control is a wettable powder formulation to be used as a spray through the ORTHO Lawn Sprayer or Sprinkling Can and is designed to control common grass diseases such as Brown Patch (*Blizzaria*), Dollar Spot and certain other diseases including Leaf Spot (*Blizzaria*), Cottony Blight (*Pythium*), Dollar Spot (*Blizzaria*), Fading Out (*Curvularia*), and Gray Leaf Spot (*Exserohilum*), particularly on St. Augustine, Bermuda, Bluegrass, Zoysia, Rye and Dichondra lawns. Do not store diluted spray. Make new solution for each use.

DIRECTIONS

To obtain maximum results it is important that the lawn be properly managed before and after treatment with CHEVRON Lawn Disease Control. Use the following Management Schedule as guide.

MANAGEMENT SCHEDULE:

1. One day before treatment — Mow lawn. Irrigate if necessary.
2. Day of treatment — After morning dew is off lawn, apply CHEVRON Lawn Disease Control as directed (see below).
3. First and second day following treatment — Do not mow lawn. Do not irrigate.
4. Repeat this schedule prior to retreatment according to preventive or curative programs outlined below.

APPLICATION RATE: Use ORTHO Lawn Sprayer or Sprinkling Can to apply treatments. Add the required amount of CHEVRON Lawn Disease Control to the Sprayer jar and add a small amount of water. Mix until a thin paste is formed, add water to bring to desired level and stir. Apply spray evenly. Fifteen gallons treats 1000 square feet. AGITATE JAR WHILE SPRAYING.


PREVENTIVE		
5 level tablespoons (1.2 oz.)	3 gallons	200 sq. ft.
25 level tablespoons (6 oz.)	15 gallons	1000 sq. ft.
CURATIVE		
5 level tablespoons (1.2 oz.)	3 gallons	100 sq. ft.
25 level tablespoons (6 oz.)	15 gallons	500 sq. ft.

PREVENTIVE PROGRAM: Apply in early fall. Successive applications may be necessary if warm weather persists during fall and winter. If necessary, treat every 30 to 35 days through February depending upon temperature and environmental conditions. Application in late February may be necessary if ideal conditions for Brown Patch persist. After first spring cutting repeat for control of such diseases as Gray Leaf Spot, Melting Out, Fading Out. Additional applications may be necessary in spring and summer if diseases persist.


CURATIVE PROGRAM: Use 6 ounces of CHEVRON Lawn Disease Control in ORTHO Lawn Sprayer and fill to 15 gallon mark and apply this to 500 sq. ft. of lawn area. Follow the Management Schedule and continue until disease is halted. Re-sod and top dress with soil, areas destroyed by disease. Continue treatments using Preventive Program.

SPECIAL COMBINATION SPRAY: To control lawn insects such as white grubs, wireworms, sod webworms and ants, use ORTHO-KLOR 74 Chlordane Spray at 1 Tablespoonful (1.2 oz.) per 1 gal. of water to cover 200 sq. ft. of lawn area with CHEVRON Lawn Disease Control at recommended rate.

COMMERCIAL APPLICATION: For use in tank type sprayers, use 4 lbs. of CHEVRON Lawn Disease Control in each 50 gallons of water to spray 10,000 square feet.



CHEVRON[®]
LAWN DISEASE
control



CHEVRON[®]
LAWN DISEASE
control

controls BROWN PATCH



GRAY LEAF SPOT
DOLLAR SPOT
FADING OUT
COTTONY BLIGHT
LEAF SPOT



CHEVRON
LAWN DISEASE
control

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

For use on lawns and turf areas to control the following diseases:
Rhizoctonia, Pythium, Sclerotinia, Fusarium, and other fungi.
Also controls the following insects:
Grasshoppers, crickets, beet beetles, and other insects.

MANAGEMENT:

APPLICATION:

Apply to lawn areas.

PREVENTION:


CONTROL:

RE-ENTRY:

RE-ENTRY:



CHEVRON
LAWN DISEASE
control



CHEVRON
LAWN DISEASE
control



CHEVRON
LAWN DISEASE
control

