

VOLCK[®] 70 Supreme Ethion Combination



ORTHO

(INSECTICIDE-MITICIDE)

[239-1585]

Active Ingredients	By Wt.
Petroleum Oil	94%
Ethion*	2%
Inert Ingredients	4%
*O,O,O',O'-Tetraethyl S,S'-methylene bisphosphorodithioate	
Contains 0.15 pound Ethion per gallon	
Petroleum Oil Min. Unsulfonated Residue — 92%	

A 70 SECOND SUPERIOR OIL
Gravity, API-34 Typical
U. S. Pat. No. 2,873,228

Do not reuse empty containers. Destroy by perforating and burning in a safe place.

DO NOT STORE AT TEMPERATURES BELOW 0° F.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

VOLCK 70 Supreme Ethion Combination is for use in the dormant and delayed dormant periods for control of scale insects, aphids and mite eggs. THOROUGH COVERAGE IS NECESSARY. The product is for use during dormant and late dormant periods only. Do not apply after foliage or edible parts begin to form. Do not apply to forage crops. Do not use in undiluted form.

NET CONTENTS
LIQUID

Chevron Chemical Company
Ortho Division/San Francisco, California 94119
Richmond, California Fresno California Honolulu Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 6278 E EPA Reg. No. 239-1585 Product 3133 Made in USA

DIRECTIONS

APPLES: European Red Mite, Apple Aphid, Rosy Apple Aphid, San Jose Scale—2 gals. per 100 gals. water. Apply during dormant or green tip period. Do not use on Astrachan, Duchess, Maria or Pratt, Meba, Wealthy, Williams Red (Red Williams), Yellow Transparent, or any variety maturing before Early McIntosh, as injury may result.

PEACHES, PEARS, PLUMS, PRUNES: San Jose Scale (Apply during dormant period), Forbes Scale, Putnam Scale (Apply during delayed dormant period)—2 gals. per 100 gals. water.

PEACHES, PLUMS: Parlatoria (Olive) Scale—3 gals. per 100 gals. water. Apply during dormant period.

PLUMS, PRUNES: Mealy Plum Aphid, Leaf Curl Plum Aphid—2 gals. per 100 gals. water. To control overwintering eggs, apply during late dormant period.

PEACHES, PEARS, PLUMS: Brown Almond Mite, European Red Mite—2 gals. per 100 gals. water. To control overwintering eggs, apply during late dormant period.

CHERRIES: San Jose Scale, Black Cherry Aphid—2 gals. per 100 gals. water. Apply during dormant period.

ORNAMENTALS (Taxus, Arborvitae, Magnolia, Tulip Tree, Maple, Elm, Catalpa, Boxwood, Privet, Oak, Ash, Poplar, Birch, Willow, Dogwood, Evonymus, Pine, Juniper): Soft Scales (Lecanium, Magnolia, Tulip Tree, Cottony Maple, European Elm, European Fruit), Hard or Armored Scales (Oystershell, Scurfy, San Jose, Pine

Needle, Juniper, Evonymus): Use 2 to 3 gals. per 100 gals. water on deciduous ornamentals and 2 gals. per 100 gals. water on evergreens. Apply as dormant spray after most danger of freezing nights has passed, but before buds have begun to open. Spray on clear mid day and do not spray in late afternoon. Be cautious with oil sprays. Do not use on easily injured plants such as sugar and Japanese Maple, Beech, Hickory, Walnut and Butternut. Oil will injure Douglas fir flower buds and remove blue color from Colorado blue Spruce.

MIXING DIRECTIONS: With agitator running start filling spray tank with water. Add the dry spray materials. The VOLCK 70 Supreme Ethion Combination should be added when tank is about one-half full and while there is sufficient agitation of the spray to insure complete emulsification.

NOTE: Do not use this material in conjunction with Sulfur before using this material. Do not combine Sulfur sprays with this material and do not follow a previous application of Sulfur until sufficient time has elapsed for safety, usually 12 to 14 days. Do not use this product on plants previously treated with Sulfur, using 1 quart of this material to 25 gallons of water. If sufficient time has not elapsed harm will show in several days. Do not use this material when temperatures are above 90° F. When temperatures drop below 32° F. (usually nighttime lows), do not spray tree and vine crops until daytime temperatures have warmed to a point above 35° F. and are free from moisture due to melting frost or heavy dews. On varieties which are sensitive to spray injury, use freshly Hydrated Lime at 2 pounds or more per 100 gallons when using Lead Arsenate.

DO NOT TAKE INTERNALLY
DO NOT INHALE
DO NOT GET ON SKIN
KEEP OUT OF REACH OF CHILDREN

WARNING: May be fatal if swallowed. Absorbed through skin. Handle concentrate in a well-ventilated area. Do not breathe vapor or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Do not store or transport near feed or food.

The product is toxic to fish and wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment and disposal of wastes. For other product only, as specified on this label.

CONDITIONS OF SALE: The Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied concerning this material.

2. Certain and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of this material. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks of liability, except those assumed by Chevron under 1. (above) resulting from handling, storage and use of this material.

VOLCK® 70 Supreme Ethion Combination

(INSECTICIDE-MITICIDE)

[239-1585]

Active Ingredients	By Wt.
Petroleum Oil	94%
Ethion*	2%
Inert Ingredients	4%
*O,O,O',O'-Tetraethyl S,S'-methylene bisphosphorodithioate	
Contains 0.15 pound Ethion per gallon	
Petroleum Oil Min. Unsulfonated Residue — 92%	

A 70 SECOND SUPERIOR OIL
Gravity, API-34 Typical
U. S. Pat. No. 2,873,228

DO NOT STORE AT TEMPERATURES BELOW 0° F.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP THIS AND ALL ORIGINAL CONTAINERS FROM CHILDREN. CONTAINERS SHOULD BE KEPT UPRIGHT AND UNOPENED.

Use this combination Ethion and oil spray during the dormant and late dormant periods of the life cycle of insects. Its application should be thorough. REFRIGERATION IS NECESSARY. This product is for use on dormant and late dormant periods only. Do not apply to foliage or other parts of plants. Do not apply to foliage or other parts of plants. Do not apply to foliage or other parts of plants.

NET CONTENTS
LIQUID

Chevron Chemical Company

Oil Division, San Francisco, California 94119

Manufactured by Chevron Chemical Company, Houston, Texas

© 1968 Chevron Chemical Company. All rights reserved.

DIRECTIONS

APPLES: European Red Mite, San Jose Scale—2 gals. per 100 or green tip period. Do not use on Bartlett, Melba, Wealthy, Will Transparent, or any variety of apple, etc.

PEACHES, PEARS, PLUMS, PRUNES: dormant period, Forbes Scale (over dormant period) 2 gals.

PEACHES, PLUMS: Parlatoria (water). Apply during dormant period.

PLUMS, PRUNES: Mealy Plum (water) 100 gals. water. To use during dormant period.

PEACHES, PEARS, PLUMS: Brown Mite 2 gals. per 100 gals. water. Apply during late dormant period.

CHERRIES: San Jose Scale, Black Scale 2 gals. water. Apply during dormant period.

ORNAMENTALS (Taxus, Arborvitae, Elm, Catalpa, Boxwood, Privet, Dogwood, Evonymus, Pine, Juniper, Nolia, Tulip Tree, Cottony Maple, Hard or Armored Scales (Coccinellids, etc.))

DO NOT
DO NOT
DO NOT

STOP

KEEP OUT OF REACH OF CHILDREN

WARNING: May be harmful if swallowed. Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly with soap and water if contact occurs. Do not breathe vapors. Do not get on clothing. Do not use near food or feedstuffs.