


[239-1624]

DIBROM[®] SEVIN[®] 2-5 Dust



ORTHO



(INSECTICIDE)

Active Ingredients	By Wt.
**Naled	2%
Carbaryl (1-naphthyl N-methylcarbamate)	5%
Inert Ingredients	93%
**1,2-dibromo-2,2-dichloroethyl dimethyl phosphate U. S. Pat. Nos. 2,971,882—2,903,478—3,009,855	

Use within _____ days of manufacture only, as this product decomposes

Date of Manufacture _____

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER

STORE IN COOL, DRY PLACE. OUT OF DIRECT SUNLIGHT.

DO NOT USE NEAR OPEN FLAMES. STAY OUT OF SMOKE.

DIRECTIONS

This product is formulated for application by ground or air equipment.

NET WEIGHT 50 LBS.

Chevron Chemical Company

1000 West 10th Street
P.O. Box 1000
Chevron, Texas 75509

BEANS: Lygus Bugs, Stink Bugs, Mexican Bean Beetle—40 lbs. per acre. Apply thoroughly when insects appear and repeat every 7 to 10 days as necessary. Do not apply within 4 days of harvest.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH THE SKIN. AVOID BREATHING DUST. Do Not Take Internally. May cause irritation to eyes, nose and throat. **Skin Contact May Be Harmful.** Avoid contact with skin and eyes. Wash Hands and Face Before Eating, Take Shower or Bath After Work. Wear Regular Long-sleeved Work Clothing. Change to Clean Clothing Daily. **Note for Physician:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not administer antiparkinsonian drugs. **AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.** Do not use on chemicals as injury may result. May corrode metal dust equipment, clean equipment thoroughly. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

COTTON: Salt-Marsh Caterpillar, Bollworms, Lygus, Flea-noppers, Aphids, Cotton Leaf Perforator—50 lbs. per acre. Apply in early morning or evening at 7 to 10 day intervals and repeat until insects are well controlled.

GRAPES: Leafhoppers, Leaf Rollers—50 lbs. per acre. Apply in early morning or evening at 7 to 10 day intervals and repeat until insects are well controlled.

STRAWBERRIES: Spider Mites, Aphids, Meadow Spittlebug, Strawberry Leaf Roller, Strawberry Weevil—40 lbs. per acre. Apply in early morning or evening at 7 to 10 day intervals and repeat until insects are well controlled.

TOMATOES: Tomato Fruitworm, Fall Armyworm, Flea Beetles, Tomato Hornworm, European Corn Borer, Stink Bugs, Leafminers, Aphids, Drosophila—50 lbs. per acre. Apply in early morning or evening at 7 to 10 day intervals and repeat until insects are well controlled.

CONDITIONS OF SALE: Chevron Chemical Company warrants that the material conforms to the chemical description on the label and is reasonably fit for use as described. It does not warrant, make any other warranty, or accept any representation to make any other warranty, except the CHEVRON GUARANTEE, guarantee or representation made by Chevron Chemical Company.

It is the policy of Chevron Chemical Company to sell its products to the general public through its authorized distributors. It is not the policy of Chevron Chemical Company to sell its products directly to the general public. It is the policy of Chevron Chemical Company to sell its products to the general public through its authorized distributors. It is not the policy of Chevron Chemical Company to sell its products directly to the general public.