

# ORTHO<sup>®</sup> Sulfur 15-40 Dust



# ORTHO

## (FUNGICIDE)

| Active Ingredients | By Wt. |
|--------------------|--------|
| Captan*            | 15%    |
| Sulfur             | 40%    |
| Inert Ingredients  | 45%    |

\*N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL, DRY PLACE  
PROTECT FROM EXCESSIVE HEAT

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKES.

### DIRECTIONS

This product is formulated for application by ground or air equipment.

**CELERY:** Pink Rot, Damp-Off—(Field)—13 lbs. per acre. (Seed Bed)—25 to 30 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

## NET WEIGHT 50 LBS.

### Chevron Chemical Company

Ortho Division, San Francisco, Calif. 94119  
Richmond, California      Fresno, California  
Des Moines, Iowa      Perth Amboy, New Jersey  
Portland, Oregon      St. Louis, Missouri  
Tulsa, Oklahoma      Wichita, Kansas

**GRAPES:** Botrytis Rot (Bunch Rot), Powdery Mildew, Silver Mite, Storage Rots—33 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically, making up to 3 cover applications before harvest.


**PEACHES:** Brown Rot, Storage Rots, Powdery Mildew, Silver Mite, Peach Scab—33 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.



## READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Avoid inhalation of dust or contact with skin. Do not store or transport near food. Food, seeds, fruits, and other plants are susceptible to injury from certain conditions. The use of Ortho Sulfur 15-40 Dust under certain conditions may be hazardous to the health of any person. Local authorities should be consulted for information on the use of this product. Do not use under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.



**PLUMS:** Brown Rot, Storage Rots, Powdery Mildew, Silver Mite—33 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

**ROSES (Field):** Rust, Powdery Mildew—35 to 40 lbs. per acre. Apply to give even coverage of upper and lower leaf surfaces at first sign of disease. Repeat at regular intervals as necessary to maintain control.

**STRAWBERRIES:** Botrytis Rot, Powdery Mildew—20 lbs. per acre immediately after picking. Repeat applications after picking as necessary.

**COMPATIBILITY:** Do not mix this product with fungicides or oil sprays.

**CONDITIONS OF SALE:** The Chevron Chemical Company warrants that the material conforms to the chemical description on the label and is reasonably fit for use as directed thereon. It does not warrant, make, or authorize any agent or representative to make any other warranty of FULFILLMENT, GUARANTEE, or similar representation, express or implied, in connection with this product.

The use of this product is subject to the provisions of the Federal Insecticide, Fungicide, and Rodenticide Act, and the regulations thereunder. The use of this product is also subject to the provisions of the Federal Food, Drug, and Cosmetic Act, and the regulations thereunder. The use of this product is also subject to the provisions of the Federal Environmental Pesticide Control Act, and the regulations thereunder. The use of this product is also subject to the provisions of the Federal Insecticide, Fungicide, and Rodenticide Act, and the regulations thereunder.