



DIFOLATAN[®] 80 Wettable

NET WEIGHT 50 LBS.

SULFENIMIDE FUNGICIDE

Active Ingredient	By Wt.
cis-N-[(1,1,2,2-tetrachloroethyl)thio]-4-cyclohexene-1,2-dicarboximide*	80%
Inert Ingredients	20%

*U. S. Pat. No. 3,178,447


READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

STORE IN A COOL, DRY PLACE.

PROTECT FROM EXCESSIVE HEAT.

POTATOES: Early Blight, Late Blight—1 to 2 pounds per acre in water to cover. Begin treatment when plants are 6" high and continue at 7 to 10-day intervals. Under severe disease conditions, use higher listed dosage or shorter spray interval.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN.

CAUTION: May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin and clothing. Avoid breathing dust or spray mist. In case of contact, flush skin and eyes with plenty of water; for any irritation get prompt medical attention. Avoid contamination of feed or foodstuffs. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is toxic to fish. Do not contaminate reservoirs, lakes, ponds or streams.

COMPATIBILITY: ORTHO DIFOLATAN 80 Wettable may be combined with most commonly used insecticides, adjuvants and fungicides with the exception of strongly alkaline material such as hydrated lime, caustic, etc., which reduces the fungicidal activity of ORTHO DIFOLATAN 80 Wettable.

CONDITIONS OF SALE: 1. Chevron Chemical Company (manufacturer) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Manufacturer neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond the manufacturer's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by the manufacturer under 1 above) resulting from handling, storage, and use of this material.

239-1716

ACCEPTED

JUN 23 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 239-1716

CHEVRON CHEMICAL COMPANY

FORM 7094-D

USDA Reg. No. 239-1716

ORTHO DIVISION — San Francisco, California

Product No. 3848

Made in U. S. A.

RICHMOND, CALIFORNIA

PORTLAND, OREGON

FRESNO, CALIFORNIA

DES MOINES, IOWA

CHERRY HILL, NEW JERSEY

DALLAS, TEXAS

WHITTIER, CALIFORNIA

HONOLULU, HAWAII

ATLANTA, GEORGIA

FORT MADISON, IOWA

ORLANDO, FLORIDA



PEACHES, APRICOTS: Scab, Brown Rot, Brown Rot Blossom Blight, Phiggy Rot - 1 to 1 1/4 lbs. per 100 gals., using sufficient spray per acre, depending on tree size, to provide thorough coverage. Apply at prebloom, full bloom, petal fall and shoot split and repeat at 10 to 14-day intervals through harvest. Do not apply after shoot split in Calif., Oregon, Washington or if fruit is to be dried.

PEACHES, NECTARINES, APRICOTS (Dormant Spray): Cankering Blight (Shot Hole), Peach Leaf Curl - 2 lbs. per 100 gals. (or 8 lbs. per acre in water to cover), and apply in fall before rains begin and repeat in late winter prior to bud swell; or 4 lbs. per 100 gal. (or 16 lbs. per acre in water to cover) and apply as a single fall application prior to cell rains.

CHERRIES (sour): Cherry Leaf Spot - 1/2 to 1 lb. per 100 gals. spray, using sufficient spray per acre, depending on tree size, to provide thorough coverage. Begin sprays at petal fall and repeat at approximately 10-day intervals for 3 to 4 applications, making the last application to the foliage after harvest.

CHERRIES, PINES, FRUIT: Brown Rot Blossom Blight (California and Oregon Only) - 1 to 2 lbs. per 100 gals. using sufficient spray per acre, depending on tree size, to provide thorough coverage. Apply at prebloom, bloom and petal fall.

TOMATOES: Anthracnose, Early Blight, Late Blight, Green Leaf Spot (Stemphylium), Yellow Spot, Fruit Rot - 1 1/2 to 3 lbs. per acre in 100 to 150 gals. water. Apply when fruit of first clusters are well formed and repeat at 7 to 10-day intervals. When weather conditions favor disease development, use the higher rate and begin applications when plants are set in fields.

MELONS (Cantaloupe, Honeydew, Muskmelon & Others): Alternaria Leaf Blight, Downy Mildew, Anthracnose, Gummy Stem Blight - 1 1/2 to 3 lbs. per acre in 100 to 150 gals. spray as required for thorough coverage, depending on size of plants. Apply when first true leaves appear and repeat at weekly intervals. If powdery mildew is a problem, use a mildew fungicide as needed.

WATERMELONS: Gummy Stem Blight, Anthracnose, Downy Mildew - 1 1/2 to 3 lbs. per acre in 50 to 150 gals. spray as required for thorough coverage, depending on size of plants. Begin applications when plants start to vine and repeat at 5 to 10-day intervals as needed.

CUCUMBERS: Downy Mildew, Scab, Gummy Stem Blight, Anthracnose, Alternaria Leaf Blight - 1 1/2 to 2 lbs. per acre in 50 to 150 gals. of spray as required for thorough coverage, depending on size of plants. Apply when first true leaves appear and continue at weekly intervals. If powdery mildew is a problem, use a mildew fungicide as needed.