

DIBROM[®] Toxaphene 3-10 Dust



ORTHO

(INSECTICIDE)

[239-1835]

Active Ingredients	By Wt.
*Naled	3%
**Toxaphene	10%
Inert Ingredients	87%
*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate	
**Technical Chlorinated Camphene	
(Chlorine Content — 67% to 69%)	
U. S. Pat. No. 2,971,882	

Use within 3 days of manufacture only, as this product decomposes.

Date of Manufacture

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL, DRY PLACE,
OUT OF DIRECT SUNLIGHT.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, California Fresno, California

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 6592-G Product 3330 Made in U.S.A.

EPA Reg. No. 239-1835-AA

DIRECTIONS

This product is formulated for application by ground or air equipment.

BEANS: Lygus Bugs, Aphids, Red Spider Mites—40 to 50 lbs. per acre. Do not apply to green or snap beans within 7 days of harvest. Do not feed treated vines to dairy animals or animals being finished for slaughter.

COTTON: Lygus Bugs, Fleahoppers, Armyworms, Bollworm, Stinkbugs, Thrips, Loopers—35 lbs. per acre. Apply when pests appear and repeat at 7 to 10-day intervals while pests persist. Do not feed treated cotton trash to dairy animals or animals being finished for slaughter. Do not use on cotton within 4 days of picking.

LETTUCE: Loopers, Cutworms, Aphids—40 to 50 lbs. per acre. Do not apply after seedling stage for leaf lettuce. Do not apply to head lettuce after heads start to form.

ORNAMENTALS (Roses, Dahlias, Chrysanthemums, Canterbury Bells, Arborvitae, Pittosporum, Snow Ball, Chinese Magnolia, Aucuba, Zinnias, Stocks, Azaleas, Willow, Privet, Holly, Birch): Lygus Bug, Thrips, Aphid, Mites—40 to 50 lbs. per acre.

PEPPERS: Hornworms, Climbing Cutworms, Loopers, Beet Armyworms, Aphids, Flea Beetles, Leaf Miners—30 lbs. per acre. Do not apply within 5 days of harvest.

TOMATOES: Leaf Miner, Thrips, Armyworms, Tomato Fruitworms, Hornworms, Aphids—20 lbs. per acre. Do not apply within 2 days of harvest.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful If Swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after using. Do not store or transport near feed or food. **Note to Physicians:** Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not use on cherries as injury may result. May corrode metal dust equipment, clean equipment after use. Do not use on cucurbits. Do not use on young plants. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product will kill fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, and ponds. Apply this product only as specified. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

