

[239-1888]



ORTHO

ACCEPTED

DEC 29 1972

UNCLASSIFIED HAS DISSEMINATION
FUNCTION AND POTENTIAL SIDE EFFECTS
ECONOMIC POLICY DECISIONS
AND UNDER NO CIRCUMSTANCES SUBJECT
TO ATTACHED COMMENTS.

POISON

If Swallowed. Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY.** **Note to Physicians:** Atropine is Antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine. **If On Skin.** Remove contaminated clothing and wash skin immediately with soap and water. **CALL A PHYSICIAN IMMEDIATELY!**

DANGER:

KEEP OUT OF REACH OF CHILDREN

DO NOT TAKE INTERNALLY

DO NOT INHALE

DO NOT GET ON SKIN

DANGER:

4.1.5. Effect of Sediment Inhaled

As a result of the "Shed" War and a lack of respirator of
any type, the U.S. Bureau of Mines for physical
protection against biological "Shed" Diseases
has been the subject of the "Shed" War and the

Do not apply this herbicide to crops or plantings that are of any kind which are in contact with soil and water. Work only on bare ground, including with soil and water, and on the surface of standing water, except for the following: (1) water in contact with a surface of any kind, including a body of water, and (2) any other body of water where there may be danger of drift. Aerial spraying is prohibited on water in contact with any surface, including a body of water, and on any other body of water. Do not use this herbicide near food or feed. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

NET WEIGHT 50 LBS.

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific requirements of the task.

LETTUCE, SPINACH: Aphids—35 to 50 lbs. per acre. Corn Earworm, Cutworms (climbing), Dipterous Leafminer (adult), Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Leafhoppers, Mites, Lygus Bugs, Salt-Marsh Caterpillar, False Chinch Bug, Thrips—50 lbs. per acre. Do not apply to lettuce within 4 days of harvest and to spinach within 7 days of harvest.

CONDITIONS OF SALE: 1. Chevron Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed herein. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, written or implied, concerning this material.

2. The user must understand and take probable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of this material. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the material is reasonably fit for the uses stated hereon, and when the label directions are followed. These and other risks are excluded, and are considered risks and liabilities of the user. Chevron disclaims liability above.

3. The user must understand and use this material in accordance with the label directions and use of this material

It is possible that the effect of the *in vitro* treatment on the *in vivo* response is not linear. For example, the effect of the *in vitro* treatment on the *in vivo* response may be more pronounced at lower concentrations of the treatment. This is a possibility that should be investigated in future studies.