


DIBROM[®] Sulfur 4-40 Dust

[239-1984]



ORTHO



Apr 30 1974

239-1984

(INSECTICIDE-FUNGICIDE)

Active Ingredients	By Wt.
*Naled	4%
Sulfur	40%
Inert Ingredients	56%
*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate U. S. Pat. No. 2,971,882	

Use within _____ days of manufacture only, as this product decomposes.

Date of Manufacture _____

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER

STORE IN COOL, DRY PLACE. OUT OF DIRECT SUNLIGHT.

DO NOT MIX OR EMPLOY IN ANY MANNER WITH WASTE OIL, FURNACE OIL, OR OTHER LIQUID OR FUMES.

DIRECTIONS

Use 40 to 60 pounds per acre unless otherwise specified. This product is formulated for application by ground or air equipment. Begin applications at first sign of insects or disease. Repeat as necessary. Apply for contact and thorough coverage to control insects and diseases.

BEANS: Aphids, Mites, Lygus Bugs, Thrips, Leafhoppers, Leafminers, Loopers—40 lbs. per acre by ground or air. Do not apply within 4 days of harvest.

CUCULI, CABBAGE, CAULIFLOWER, COLLARDS, BRUSSELS SPROUTS, KALE: Cabbage Looper, Imported Cabbageworm, Diamond-Back Caterpillar, Aphids. May be used 1 day before harvest on Brussels Sprouts, Cabbage, Cauliflower and Brussels Sprouts. Do not apply to Collards and Kale within 4 days of harvest.

CITRUS (Oranges, Lemons, Grapefruit): Aphids, Red Spider Mites, Citrus Cutworms, Thrips—Do not apply within 4 days of harvest. California and Arizona do not harvest within 1 day of harvest.

COTTON: Lygus, Cotton Leaf Perforator, Armyworms, Loopers, Fleahoppers—20 lbs. per acre. Apply at 7 to 10 day intervals while pests persist. Do not apply within 4 days of hand picking.

GRAPE: Leafhoppers, Powdery Mildew—40 to 60 lbs. per acre. Thrips—20 to 25 lbs. per acre. Do not apply within 4 days of harvest.

LETTUCE, SPINACH, CELERY, CHARD, TURNIP GREENS: Loopers, Aphids—37 lbs. per acre. May be used 1 day before harvest on Lettuce, Spinach, Turnip Greens and Chard. Do not apply to Celery within 4 days of harvest.

PEACHES: Western States Only—Fruit Flies (Drosophila), Red Spider Mites, Peach Twig Borer, Lygus Bugs, Thrips (Calif. Only), Powdery Mildew—50 lbs. per acre. Do not apply within 4 days of harvest.

STRAWBERRIES: Spider Mites, Aphids, Powdery Mildew—Do not apply within 4 days of harvest.

TOMATOES (Field): Two-Spotted Mites, Aphids, Leaf Miners, Tomato Fruitworms, Hornworms, Tomato Russet Mites—20 to 30 lbs. per acre. Make first application at 5 to 7 days before first picking and continue at 5 to 7-day intervals until harvest is complete. Do not apply within 1 day of harvest because of toxic fur content of this dust, first obtain approval from your canner or processor before using.

CONDITIONS OF SALE: 1. Chevron Chemical Company warrants that this material conforms to the description on the label and is reasonably fit for the uses stated herein. Chevron neither makes nor authorizes any agent or representative to make, any other warranty, representation, express or implied, concerning this material. 2. Chevron and its authorized factors, buyers, Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated herein and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability for the use of this material as stated by Chevron under 1 above.

3. The user is responsible for handling, storage, and use of this material.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after use. Do not eat, drink, or smoke while using.

Note to Physicians: In case of emergency, call your physician or poison control center. Do not induce vomiting unless advised to do so by a physician.

This product is highly toxic to bees exposed to direct treatment. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply when runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is highly toxic to bees exposed to direct treatment. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco California 94119
Richmond, California Fresno California
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida
Form 6345-T Product 3971 Made in USA
EPA Reg. No. 239-1984-AA