


DIBROM ZINC COPOSIL 4-35 Dust

[239-2148]



ORTHO

(INSECTICIDE - FUNGICIDE)

Active Ingredients	By Wt
*Naled	4.0%
Copper in basic cupric zinc sulfate complex expressed as elemental	6.6
Inert Ingredients	89.4
Contains 6.6% zinc as elemental derived from basic cupric zinc sulfate complex.	
*1,2-dibromo 2,2-dichloroethyl dimethyl phosphate U.S. Pat. No. 2,971,882	

Use within 3 days of manufacture, as this product decomposes.

Date of Manufacture _____

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. STORE IN COOL DRY PLACE, OUT OF DIRECT SUNLIGHT.

DIRECTIONS

Use 40 to 50 pounds per acre unless otherwise specified. This product is formulated for application by ground or air equipment. For Airplane Application, use heavier dosages. Begin applications at first sign of insects or diseases. Repeat as necessary unless otherwise specified. Apply for contact and thorough coverage to obtain control of insects.

BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS: Cabbage Looper, Imported Cabbageworm, Diamond-back Caterpillar, Aphids. May be used 1 day before harvest.

NET WEIGHT 50 LBS.

Chevron Chemical Company
 Ortho Division San Francisco, Calif. 94119
 Richmond, California Fresno, California Des Moines, Iowa
 Perth Amboy, New Jersey Orlando, Florida
 Form 5889 I Product 2968 Made in U.S.A.
 EPA Reg. No. 239-214-AA

CUCUMBERS: Looper Caterpillars, Aphids, Diabrotica Beetles, Lygus Bugs, Powdery and Downy Mildew—Use 30 to 35 lbs. per acre. Apply when diseases appear. Repeat at 7 to 10 day intervals. May be used up to harvest.

GRAPES: Leafhoppers. Do not apply within 4 days of harvest.

LETTUCE, SPINACH, CELERY: 37 lbs. per acre.

Looper Caterpillars, Aphids, Downy Mildew, Early and Late Blight of Celery. May be used up to 1 day before harvest on Lettuce and Spinach. Do not use on Celery within 4 days of harvest.

MELONS: Aphids, Cabbage Looper, Armyworms, Leafhopper, Lygus Bugs, Leaf Spot, Downy Mildew, Powdery Mildew—40 lbs. per acre. May be used 1 day before harvest.

STRAWBERRIES: Red Spider Mites, Aphids. Do not apply within 4 days of harvest.

TOMATOES: Tomato Fruitworms, Hornworms, Leafminers, Late Blight—Use 25 lbs. per acre. Do not apply within 4 days of harvest. May starting before early fall rains are expected. Repeat if conditions according to local practices. May be used 1 day before harvest.

CONDITIONS OF SALE: 1. Chevron Chemical Company, Chevron warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability, except those assumed by Chevron under 1 above, resulting from handling, storage, and use of this material.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after using. Do not store or transport near feed or food. **Note to Physicians:** Atropine is antidotal. 2 PAM is also antidotal and may be administered in conjunction with atropine. Do not use on cherries as injury may result. May corrode metal dust equipment, clean equipment after use. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residue on plants. Protection of bees may be achieved from your Cooperative Agricultural Extension Service.