

239-2181

8-11-1977

1/2



ORTHO SEVIN* Garden Dust

KILLS PESTS

IN THE GARDEN:

Tomato Hornworms,
Armyworms, Cutworms,
Certain Beetles, Squash Bugs
and Certain Other Garden Pests

ON DOGS AND CATS:

Fleas, Ticks
on Animals and in Areas
Where Animals Live



| Active Ingredient | By Wt. |
|---|--------|
| Carbaryl (1-naphthyl N-methylcarbamate) | 5% |
| Inert Ingredients | 95% |

CAUTION: Keep out of reach of children.
See back panel for additional cautions.
NET WT. 1 LB.

2/2



EPA Est. 239-CA-1

ORTHO SEVIN* Garden Dust

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

Apply dust at first sign of insects. Repeat as necessary. For best results, dust in early morning or late evening when temperature is low and air is still. Apply dust so that all parts of plants are contacted, especially upper and lower leaf surfaces. Apply dust as lightly as possible. Avoid excessive applications.

| Plants | Insects | Comments |
|--|--|---|
| BEANS | Mexican Bean Beetle, Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Velvetbean Caterpillar, Flea Beetles, Japanese Beetle, Corn Earworm, Armyworms, Tarnished Plant Bug, Stink Bugs. | May be applied up to harvest. |
| CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI | Flea Beetles, Harlequin Bug, Armyworms, Cabbage Caterpillars, Corn Earworm. | Do not apply within 3 days of harvest. |
| CORN | Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetle, Leafhoppers, Flea Beetles. | For Corn Borer—Apply thoroughly into the whorls when plant is about knee-high. Repeat at weekly intervals until control is achieved. For Corn Earworm—Apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals. Re-apply immediately after rain. May be applied up to harvest. |
| CUCUMBERS, MELONS, PUMPKIN, SQUASH | Pickleworm, Melonworm, Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs. | Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not use SEVIN on watermelons in Florida. May be used up to harvest. |
| LETTUCE, CARROTS | Flea Beetles, Harlequin Bug, Leafhoppers, Six-spotted Leafhopper, Armyworms, Cabbage Caterpillars, Corn Earworm, Spittlebugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs. | May be used up to harvest on Carrots. Do not use on Head Lettuce within 3 days of harvest and on Leaf Lettuce within 14 days. |
| OKRA | Corn Earworm, Stink Bugs. | Repeat at 5 to 7 day intervals or as necessary. May be applied up to harvest. |
| POTATO, EGGPLANT, PEPPERS | Colorado Potato Beetle, Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus Bugs, Stink Bugs, Lace Bugs, Tarnished Plant Bugs. | May be applied up to harvest. |
| STRAWBERRIES | Meadow Spittlebug, Strawberry Leaf Roller, Strawberry Weevil. | May be applied up to 1 day of harvest. |
| TOMATO | Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer. | May be applied up to harvest. |

DOGS, CATS: Brown Dog Ticks, Fleas—Dust lightly over the animal and rub thoroughly into the skin. Apply also to the feet, between the toes and on the legs. Apply at weekly intervals if needed. Avoid treatment of kittens and puppies less than 4 weeks of age.

IN OR AROUND DOG AND CAT QUARTERS: Ticks, Fleas—Dust lightly in and around sleeping quarters, kennels and dog houses by means of a shaker can or hand duster. Force dust into cracks and crevices. Repeat as necessary. **DO NOT APPLY TO RUGS AND CARPETS AS DAMAGE MAY OCCUR.**

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Skin Contact. Wash Hands and Face Before Eating. Wear clean clothing.

Note to Physicians: Emergency Information—call (415) 233-3737. Carbaryl is a moderate, reversible Cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD. Dispose of container promptly when empty.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

Chevron Chemical Company

Ortho Division/San Francisco CA 94119 Richmond CA 94804
Form 7325-D Product 3819 Made in U.S.A.

EPA Reg. No. 239-2181-AA

ACCEPTED
AUG 11 1977
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide EPA Reg. No. 239-2181