## **SEVIN**SEVIN STHO Garden Dust

ACCEPTEDSE

CAUTION: stemilists and contact with ever stage and reference to restrict the stage of the stage PREADTROMANT STATEMENTS
CAUTION: Hamiltonian and And contect with 1945, state and the State of t

Apoly dust at first sup of inserts Repeat as necessary. For bass results, dust in early morning or in last eyeaking when temperature is the temperature of the temperature is the temperature of the temperature of the temperature is the temperature of temperature of the temperature of the temperature of the temperature of temperature of the temperature of temperature of the temperature of temperature OMECTABLE TOR USE It is a revolving of Sadara law is use the product in a manner inconsistent with its labelong. It is a revolving of Sadara law is use a Accompanies with Label, Precentification is statistically and Describes.

Z SEE	heacts	Caracti
BEANS	Mauran Been Beatle, Leefingopers, Sean Leaf Berslie. May be applied up to hervest. Cusumber Seetles, Wessen Been Cuswom, Valver. Dean Categolder, File Beetles, Appenses Beetle, Corn Envorm, Authorins, Tensahel Pant Bug, Sint Bugs.	May be applied up to hervest.
CABBAGE, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI	Fat Berties, Hariequio Beg, Armyweoms, Cabbage. Do not apply within 3 ders of harvest. Caterpiffest, Corn Earworm.	Do not apply within 3 days of harvest.
CORN	Order Revents Core Recoverent Adults, European Core for Care Boars - Apple Recognitive to the whole the Boars and Boars fail Annexem, Sport Feld Manneyorn. Sport Board St., Japanese Boarle, plant is about been been Report at weekly disterval Leathloppers. Fire Beetles, Jesus Holles and Sold St. Report St. R	For Core Bears — Apply thoroughly into the wheats when plant is about have been, Repeat at a weekly interval until control is achieved. For Core Enverses — Apply to silts as soon as they appear. Make 3 to 4 applications as 2 to 4 applications.
CUCUMBERS.	Pickleworm, Melonworm, Cedumber Beetles, Fles Avoid excessive applications, Same lesi insury may	Avoid excessive applications. Same lest injust may

Corn Earworm, Sint Bugs.

Caturade Peters Beetle, Rea Beetle, Lesthappers. May be applied up to harrest.

Caturade Peters Beetle, Rea Beetle, Lesthappers. May be applied up to harrest.

Total Destituteon, R.A. Armyworm, Total by horses.

Lest Bugs, Investing Stand Bugs.

May Be applied up to harvest.

May be applied up to ledy of harvest.

May be applied up to ledy of harvest.

Feet Seetles, Lesthoppers, Learnes Frithworm, Fall May be applied up to ledy of harvest.

Anymorm, Total to pers. Learnes Frithworm, Fall May be applied up to harvest.

TRAWBERRIES TOMATO DOGS, CATE. Take, These — Dots lightly over the annead and rot throughly ratio the slain. Apply size to take thest, becomes takes of the lightly over the annead and rot throughly ratio the slain. Apply size to take the slain of the slain throughly throughly throughly the slain throughly compared consistent. Annead assuments in several layers throughly th





Chevron Chemical Company one Other Contumer Pressurs Devision

The Treatest CA WIPP-14H Historial CA Minol LISTS

Fam 7255 B.7 Froduct 2019 Male m U.S.A.

EM. Neg. No. 236-2181-54 CFA Est. 228-NJ-1, 228-MG-1, 238-ML-1 Superscript used as first fetter of fet such

0

THEFT WEEK LED BY RDA

## COPY A

PRECAUTIONARY STATEMENTS 12 pt.

HAZARDS TO HUMANS & DOMESTIC ANIMALS 12 pt.

CAUTION 12 pt.

Harmful if swallowed. Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT 12 pt.

If swallowed, DO NOT make person vomit unless directed to do so by a doctor. Immediately take person and container of product to, or if not possible, call a doctor or an emergency hospital. In case of eye contact, immediately flush eyes with fresh water for 15 minutes. Get medical attention if irritation persists. For skin, wash with plenty of soap and water. Note to Physicians: Emergency Information - call (415) 233-3737. Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS 12 pt.

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

## COPY B

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

## COPY C

Do not use on pregnant dogs.

CONFIDENTIAL

PROPERTY OF

CHEVRON CHEMICAL COMPANY

UNTIL ACCEPTED BY EPA