

PM19

239-2181

1/4

[OUTDOOR ICON] OUTDOOR USE ONLY

ORTHO® SEVIN® GARDEN DUST

[ABN: ORTHO® SEVIN® brand carbaryl insecticide Garden Dust]

~~KILLS GARDEN PESTS IN THE GARDEN: Armyworms, Cutworms, Squash Bugs, Tomato Hornworms, Certain Beetles And Other Garden Pests As Listed.~~

~~KILLS FLEAS & TICKS ON DOGS AND CATS AND IN DOGHOUSES~~

~~Kills: Beetles, Bugs, Leafhoppers and Worms as Listed On Vegetables.
May be Used up to Harvest Unless Otherwise Noted.~~

KILLS PESTS

~~IN THE GARDEN: Tomato Hornworms, Armyworms Cutworms, Certain Beetles
Squash Bugs and Certain Other Garden Pests~~

~~ON DOGS AND CATS: Fleas, Ticks on Animals and in Areas Where Animals Live~~

Active Ingredient

Carbaryl (1-naphthyl N-methylcarbamate)..... 5%

Inert Ingredients.....95%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional precautionary statements.

~~Specially formulated for residential use.~~

NET WT 1 LB / 453 g

ACCEPTED
APR 02 1997
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 239-2181

96 DEC 31 AM 1:10

REC'D.../DP01

Kills Pests in the Garden: ~~Tomato Hornworms, Armyworms, Cutworms, Certain Beetles, Squash Bugs and Certain Other Garden Pests. Kills Fleas & Ticks on Animals and in Kennels and Dog Houses.~~

READ ENTIRE LABEL BEFORE EACH USE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

~~READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.~~

VEGETABLES & FRUITS

PLANTS	INSECTS/PESTS	WHEN TO HARVEST
BEANS	Mexican Bean Beetles, Leafhoppers, Bean Leaf Beetles, Cucumber Beetles, Western Bean Cutworms, Velvet Bean Caterpillars, Flea Beetles, Japanese Beetles, Corn Earworms, Armyworms, Tarnished Plant Bugs, Stink Bugs	May be applied up to day of harvest.
CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER KOHLRABI	Flea Beetles, Harlequin Bugs, Armyworms, Cabbage Caterpillars, Corn Earworms	Do not apply within 3 days of harvest.
CORN	Corn Earworms, Corn Rootworm Adults, European Corn Borers, Fall Armyworms, Sap Beetles, Japanese Beetles, Leafhoppers, Flea Beetles	May be applied up to day of harvest.
CUCUMBERS MELONS, PUMPKINS, SQUASH	Pickleworms, Melonworms, Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs	May be applied up to day of harvest.
Do not use this product on watermelons in Florida.		
LETTUCE, CARROTS	Flea Beetles, Harlequin Bugs, Leafhoppers, Sixspotted Leafhoppers, Armyworms, Cabbage Caterpillars, Corn Earworms, Spittlebugs, Tarnished Plant Bugs, Lygus Bugs, Stink Bugs	May be applied up to day of harvest to carrots. Do not apply to head lettuce within 3 days of harvest or to leaf lettuce within 14 days of harvest.
OKRA	Corn Earworms, Stink Bugs	May be applied up to day of harvest.
POTATOES, EGGPLANT, PEPPERS	Colorado Potato Beetles, Flea Beetles, Leafhoppers, Tomato Fruitworms, Fall Armyworms, Tomato Hornworms, European Corn Borers, Lygus Bugs, Stink Bugs, Lace Bugs, Tarnished Plant Bugs	May be applied up to day of harvest.
STRAWBERRIES	Meadow Spittlebugs, Strawberry Leafrollers, Strawberry Weevils	Do not apply within 1 day of harvest.

TOMATOES	Flea Beetles, Leafhoppers, Tomato Fruitworms, Fall Armyworms, Tomato Hornworms, European Corn Borers	May be applied up to day of harvest.
----------	--	--------------------------------------

HOW TO APPLY

- Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply dust so that all part of plants are contacted, especially upper and lower leaf surfaces. Apply dust as lightly as possible. Avoid excessive applications.
- Apply when air is calm to avoid drift and contact with eyes and skin.
- Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.
- For plants above chest height, consider using a liquid carbaryl product.

WHEN TO APPLY

- Apply dust at first sign of insects/pests.
- For best results, Dust in early morning or in late evening when temperature is low and air is still.
- Repeat/Reapply as necessary, but not more often than once weekly.
- CORN: For European Corn Borers, apply thoroughly into the whorls when plant is about knee high. For Corn Earworms, apply to silks as soon as they appear. Make 3 to 4 applications.
- CUCUMBERS, MELONS, PUMPKINS, SQUASH: Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

Plants	Insects	Comments
BEANS	Mexican Bean Beetle, Leafhoppers, Bean Leaf Beetle, Cucumber beetles, Western Bean Cutworm, Velvetbean Caterpillar, Flea Beetles, Japanese Beetle Corn Earworm, Armyworms, Tarnished Plant Bug, Stink Bugs.	May be applied up to harvest.
CABBAGE, BROCCOLI, BRUSSELS SPROUTS,	Flea Beetles, Harlequin Bug, Armyworms, Cabbage Caterpillars, Corn	Do not apply within 3 days of harvest.
CAULI FLOWER, KOHLRABI	Earworm.	
CORN	Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall	For Corn Borer Apply thoroughly into the whorls when plant is about knee high. Repeat at weekly intervals until control is

Armyworm, Sap ~~achieved. For Corn Earworm~~
Beetles, Japanese ~~Apply to silks as soon as they~~
Beetle, Leafhop ~~appear. Make 3 to 4 applications~~
pers, Flea Beetles. ~~at 2 to 3 day intervals. Re apply~~
~~immediately after rain. May be~~
~~applied up to harvest~~

CUCUMBER — Pickleworm, Melon — Avoid excessive applications.
 MELON — worm, Cucumber — Some leaf injury may occur if
 SQUASH — Beetles, Flea Beetles — treatments are made when tender
 PUMPKIN — Leafhoppers, Squash — foliage is wet or in the presence of
 Bugs. — prolonged high humidity. Do not
 use SEVIN Carbaryl insecticide on
 Watermelons in Florida. May be
 used up to harvest.

LETTUCE, — Flea Beetles, Harle- May be used up to harvest on
 CARROTS — quin Bug, Leafhop Carrots. Do not use on Head-
 per, Army worms, — Lettuce within 3 days of harvest
 Cabbage Caterpillars, — and on Leaf Lettuce within 14-
 Corn Earworm, Spittle — days.
 bugs, Tarnished Plant
 Bugs, Lygus Bugs,
 Six spotted leafhopper,
 Stink Bugs.

OKRA — Corn Earworm, Stink — Repeat at 5 to 7 day intervals or
 Bugs — as necessary. May be applied up
 to harvest.

POTATO, — Colorado Potato Beetle — May be applied up to harvest.
 PEPPERS, — Flea Beetles, Leaf-
 EGGPLANT — hoppers, Tomato Fruit
 worm, Fall Armyworm,
 Tomato Hornworm, Euro-
 pean Corn Borer, Lygus
 Bugs, Stink Bugs, Lace
 Bugs, Tarnished Plant
 Bugs.

STRAW — Meadow Spittlebug, — May be applied up to 1 day of
 BERRIES — Strawberry Leafroller — harvest.
 Strawberry Weevil.

TOMATO — Flea Beetles, Leaf — May be applied up to harvest.
 hoppers, Tomato
 Fruitworm, Fall Army
 worm, Tomato Horn-
 worm, European Corn
 Borer.

4/7

DOGS AND CATS

To control Ticks and Fleas: Use product only outside or in a well ventilated area. Wearing household latex or rubber gloves, lightly dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. This product should not be applied by children. Dust lightly over the animal and rub thoroughly into the skin. Apply also to the feet, between toes and on the legs. Repeat/Reapply as necessary, but not more often than once weekly. Apply at weekly intervals if needed. **DO NOT USE ON PREGNANT DOGS.** Avoid treatment of ~~Do not use on kittens or~~ and puppies less than 124 weeks of age. **NOTE:** Do not allow treated pets back on carpets or rugs in the house as this product may stain or discolor rugs and wall-to-wall carpets. Dust bedding and doghouse lightly in and around kennels and dog houses by means of using a shaker can or other dust applicator-hand duster. Force dust into cracks and crevices. Repeat/Reapply as necessary, but not more often than once weekly. **DO NOT USE IN THE HOME: DO NOT APPLY TO RUGS AND CARPETS AS STAINING MAY OCCUR.**

STORAGE AND DISPOSAL

STORAGE: Store in original container in a safe place. Keep pesticide in original container. Do not put into food or drink containers. Avoid contamination of food and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. **DISPOSAL:** Do not reuse container. Securely wrap partially filled or empty container in newspaper and discard in trash. **PRODUCT** Partially filled canister may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER** Do not reuse canister. Discard canister in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly after handling.

USER SAFETY REQUIREMENTS

- | |
|---|
| <ul style="list-style-type: none"> • Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product. • Change clothing as soon as possible after use. • Wash the outside of gloves before removing. As with any pesticide product, wash thoroughly immediately after handling and before eating, smoking or using the toilet. |
|---|

(reentry icon) Do not allow children or pets to contact treated area until dust has settled.

FIRST AID: If swallowed, immediately telephone a poison control center, emergency treatment center or a physician for advice. DO NOT make person vomit unless directed to do so by medical personnel a doctor. If medical advice cannot be obtained, then immediately take person and product container to an emergency treatment center or hospital of product to, or if not possible, call a doctor or an emergency hospital. If in eyes, flush eyes with plenty of water. In case of eye contact, immediately flush eyes with fresh water for 15 minutes. Call a physician Get medical attention if irritation persists. For skin, wash with plenty of soap and water. **Note to Physicians:** Emergency Information call 1-800-454-2333. Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming plants crops or weeds. Do not apply this product or allow it to drift to blooming plants crops or weeds if bees are visiting the treatment area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Medical Information call 1-800-454-2333
Product Information call 1-800-225-2883

ORTHO® SOLARIS® Registered Trademarks of Monsanto Company
SEVIN® is the registered trademark of Rhone-Poulenc for carbaryl insecticide.
Monsanto Company _____

Manufactured for
The SOLARIS Group of
Monsanto Company
P.O. Box 5008
San Ramon CA 94583-0808
Form _____
Product 3910
EPA Reg. No. 239-2181
EPA Est. 58996-MO-1^A, 769-GA-1^G
Superscript is first letter of lot number
MADE IN USA