



SEVIN^{*} Bait (Pelleted)

[INSECTICIDE]

NET WEIGHT 50 LBS.

Active Ingredient
Carbaryl (1-naphthyl N-methylcarbamate) 5%
Inert Ingredients 95%
U. S. Pat. Nos. 2,903,478 - 3,009,855

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

Read and apply in accordance with directions on label and government regulations. Repeat treatment as necessary.


CUTWORMS, ARMYWORMS, SOWBUGS, GRASSHOPPERS, DARKLING GROUND BEETLES AND CRICKETS IN CALIFORNIA, ARIZONA, NEVADA AND UTAH: Cucumbers, Squash, Melons. 20 lb. per acre. May be applied to alfalfa, Alfalfa, Clover, Sorghum, Cotton, Peas. 10 lb. per acre. May be applied to alfalfa, Alfalfa, Clover, Sorghum, Cotton, Peas. Do not apply to Sorghum within 21 days of harvest for use as forage. Asparagus, Beans, Broccoli, Beets, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Collards, Corn, Eggplants, Endive, Lettuce, Horseradish, Kale, Okra, Parsley, Parsnips, Peppers,



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeve Work Clothing. Change to Clean Clothing Daily. Note for Physicians: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES. This product is highly toxic to bees exposed to direct treatment or residue on crops. Protection information may be obtained from your Cooperator, Agricultural Extension Service. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.



Potatoes, Radishes, Rutabagas, Spinach, Strawberries, Sugar Beets, Swiss Chard, Tomatoes, Turnips. 40 lb. per acre. Apply during cutting season on Asparagus. Repeat application on Asparagus within 7 days and do not apply within 1 day of harvest. Do not apply within 1 day of harvest on Strawberries. Do not apply within 2 days of harvest on Broccoli, Brussels Sprouts, Cabbage, Endive, Flower, Head Lettuce, Horseradish, Parsnips, Swiss Chard, Rutabagas. Do not apply within 14 days of harvest on Sugar Beets, Cellulose, Beans, Kale, Leaf Lettuce, Peas, Spinach and Swiss Chard. Do not apply to beets within 3 days of harvest or within 14 days of harvest to be used for feed or food. Grapes. Do not apply within 1 day before harvest.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed thereon. Chevron neither makes, nor authorizes any agent or representative to make any claim with respect to the material. 2. Chevron and its agent, distributor, buyer and user's control prevent it from accepting all risks in connection with the use of chemical. Chevron and its agent, distributor, buyer and user's control prevent it from accepting all risks in connection with the use of chemical. Chevron and its agent, distributor, buyer and user's control prevent it from accepting all risks in connection with the use of chemical. Chevron and its agent, distributor, buyer and user's control prevent it from accepting all risks in connection with the use of chemical.

THE CHEVRON CHEMICAL COMPANY
SAN FRANCISCO, CALIFORNIA

CHEVRON CHEMICAL COMPANY

FORM 7428 D

USDA Reg. No. 239-2235

ORTHO DIVISION

San Francisco, California 94120

Product No. 3857

RICHMOND, CALIFORNIA

PORTLAND, OREGON

FRIEDLAND, CALIFORNIA



DES MOINES

IOWA

CHERRY HILL, NEW JERSEY

DAVIDSON, TEXAS