

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

READ DANGER AND FIRST AID TREATMENT ON FRONT PANEL AND WORKER SAFETY RULES ON BACK OF PAMPHLET

NOTE: Do Not Use Around Home Gardens, Schools, Recreational Parks or Playgrounds.

Do not combine ORTHO GRAMOXONE CL with liquid fertilizers except as specified below. DO NOT combine with granular or slow-type liquid fertilizers as the activity of ORTHO GRAMOXONE CL will be reduced.

DIRECTIONS

ORCHARDS, VINEYARDS, WINDBREAK, SHADE AND ORNAMENTAL TREES: For kill of annual broadleaf weeds and grasses and for top kill and suppression of perennials in the interspaces and around the base of trees and vines—Almonds, Apples, Apricots, Avocados, Bananas, Cherries, Citrus, Coffee, Figs, Filberts, Grapes, Macadamia Nuts, Nectarines, Olives, Papayas, Peaches, Pears, Plums, Prunes, Walnut and other Trees such as Arborvitae, Ash, Elm, Fir, Oak and Pine—Apply as a directed spray at 1 to 2 qts. per sprayed acre in 50 to 200 gals. water to thoroughly cover the weeds and grasses. Add ORTHO X-77 Spreader (non-ionic) at 8 oz. per 100 gals. Apply when the weeds and grasses are succulent and the new growth is from 1 to 6 inches high. For mature woody weeds, late germinating weeds and grasses, and for perennials—retreatment or spot treatment may be necessary.

CAUTION: Do not allow spray to contact green stems, fruit or foliage as injury may result. Do not spray under windy conditions and use a shield for young trees or vines. Do not allow animals to graze on treated areas.

GUAVA: Apply as a directed spray at 2 qts. per sprayed acre in 50 to 200 gals. to thoroughly cover the weeds and grasses. Add ORTHO X-77 Spreader (non-ionic) at 8 oz. per 100 gals. Apply when the weeds and grasses are succulent and the new growth is from 1 to 6 inches high. For mature woody weeds, late germinating weeds and grasses and for perennials—retreatment or spot treatment may be necessary. **CAUTION:** Do not allow spray to contact green stems, fruit or foliage as injury may result. Do not spray under windy conditions and use a shield for young trees. Do not allow animals to graze on treated areas.

PASSION FRUIT: Apply as a directed spray at 2 qts. per sprayed acre in 50 to 200 gals. to thoroughly cover the weeds and grasses. Add ORTHO X-77 Spreader (non-ionic) at 8 oz. per 100 gals. Apply when the weeds and grasses are succulent and the new growth is from 1 to 6 inches high. For mature woody weeds, late germinating weeds and grasses and for perennials—retreatment or spot treatment may be necessary. **CAUTION:** Do not allow spray to contact green stems, fruit or foliage as injury may result. Do not spray under windy conditions and use a shield for young vines. If bark is still green, wrap vine prior to application to prevent injury. If application is to be made during harvesting season, apply ORTHO GRAMOXONE CL only after picking fruits off the ground. Do not allow animals to graze on treated areas.

PREPLANT AND PREEMERGENCE USE—For Kill of Emerged Annual Broadleaf Weeds and Grasses and for Top Kill and Suppression of Perennials—Apply when the weeds and grasses are succulent and growth is from 1 to 6 inches high (danger plants are less affected by this treatment).

NOTE: To prevent injury to desirable plants, do not spray directly on or near them. Apply prior to preemergence treatment. Do not spray on roads, foot paths, bare soil, and on plants in flower.

CORN, LETTUCE, MELLONS, PEPPERS, SOYBEANS, SORGHUM, SUGAR BEETS, TOMATOES: Preplant or Preemergence Treatment—Apply 1 to 2 qts. per sprayed acre as a band treatment over the crop row, or as a broadcast treatment, prior to, during or after planting—but before emergence of the crop. Use 20 to 60 gals. diluted spray. Add ORTHO X-77 Spreader (non-ionic) at 8 oz. per 100 gals. of diluted spray. Seedbeds should be formed as far ahead of planting and treatment as possible to permit maximum weed and grass emergence. Seeding should be done with a minimum amount

of soil disturbance. Weeds and grasses emerging after application will not be controlled. Crop plants emerged at time of application will be killed.

NONCROP AREAS (Roadsides, Highway Margins, Rights-of-way, Around Commercial Buildings, Power Plants, Storage Yards and Other Installations, Fence Lines, Parkways and similar Noncrop Areas)—For the control of Annual Broadleaved Weeds and Grasses (such as Burdock, Chickweed, Filaree, Groundsel, Nettle, Pigweed, Plantain, Puncturevine, Purslane, Red Clover, Shepherdspurse, Thistle, White Mustard, Wild Radish, Wild Oats, Blue Grass, Cheat Grass, Cr. s) and Top Kill and suppression of Perennial Weeds (such as Bermudagrass, Johnsongrass and Morning-glory)—1 to 2 qts. per acre. Apply for full coverage and thorough weed contact (50 to 100 gals. dilute spray per acre). Best results are obtained when applications are made to young succulent weeds and grasses. Mature woody weeds are less susceptible. Repeat as needed. Add ORTHO X-77 Spreader (non-ionic) at 8 oz. per 100 gals. ORTHO GRAMOXONE CL is compatible with many residual herbicides. **CAUTION:** Avoid spray contact with foliage or fruit of food crops and ornamentals.

SUGARCANE: Preharvest Crop Desiccation—Apply 1 pt. per acre. Apply to mature plant or stubble cane using 8 to 10 gals. water per acre. ORTHO X-77 Spreader (a non-ionic surfactant) should be added to all sprays at the rate of 1 pt. per 100 gals. dilute spray. Apply with an aircraft sprayer 7 to 10 days before burning and harvest. For best results, burn desiccated sugarcane when foliage is dry. For rapid results apply when weather is clear and warm. Apply as a coarse spray at low pressure when air is calm to prevent spray drift.

WORKER SAFETY RULES

Avoid accidents. Follow these pesticide Safety rules when handling this product.

USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

WASH SPLASHES from skin and eyes immediately.

REMOVE and wash contaminated clothing.

WASH before eating, smoking and drinking.

CONCENTRATE/MIXING

Wear full face shield, rubber gloves and apron when handling concentrate.

SPRAY APPLICATION

Avoid working in spray mist. If there is risk of exposure wear goggles and approved face mask capable of filtering spray droplets. Wear waterproof footwear and clothing when spraying or when contacting vegetation wet with spray.

ENVIRONMENTAL SAFETY CAUTIONS

WILDLIFE

This product is toxic to wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur.

DRIFT

ORTHO GRAMOXONE CL is a contact herbicide that kills or injures plants through direct spray contact. It is not systemic and will not move through soil to damage other plants that may be growing nearby. Do not spray on or near desirable plants.

EQUIPMENT CONTAMINANTS

Do not use ORTHO GRAMOXONE CL in areas where it may contact desirable plants. Wash spray equipment and containers thoroughly with water immediately after use. WASH and decontaminate when empty. Never reuse.

DO NOT contaminate water when cleaning equipment or disposing of wastes. Apply the product only as specified on this label.

• STORE AT TEMPERATURES ABOVE 32°F.

Do not contaminate feed, foodstuffs or drinking water.

Do not store next to feed or food, or transport in or on vehicles containing foodstuffs or feeds.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

GRAMOXONE is a registered trademark of Plant Protection Limited, a subsidiary company of Imperial Chemical Industries Limited.

Manufactured for

Chevron Chemical Company

Ortho Division, San Francisco, CA 94104 • Richmond, CA 94701

Form PE-8914 • Product 1000-72

EPA Reg. No. 239-2286