

ORTHOCIDE[®] DIBROM[®]
10-4 Dust BEST DOCUMENT AVAILABLE



BEST DOG

GRAPES Leafhoppers, Bunch Rot—40 lbs per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically up to 3 applications before harvest. Do not use within 4 days of harvest.

MELONS (Cantaloupes, Honeydews, Watermelons), Cucumbers, Summer Squash, Winter Squash, Dainty Midew, Anthracnose, Cucumber Beetles, Aphids, Leafminers, Pickworms, Armyworms, loopers, Leafhoppers—25 lbs. per acre. Dig for signs of the first sign of a pest and report it to your neighbors.

Do not apply within 1 day of blouses, Honeysuckles, Watermelon, apply within 4 days of horseradish, Summer Squash

STRAWBERRIES: Red Spider *M. t.*
Aphids, Botrytis Rot—40 lbs. per
ton when new growth starts in
form. Repeat application week
days of harvest.

The main . . . days of re-education only, as the pro-
duct decreases.

Date of Manufacture

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE
WITH CAUTIONS AND DIRECTIONS, AND WITH APPLI-
CABLE STATE AND FEDERAL REGULATIONS

2. * * * * * IN ORIGINAL CONTAINER

STORE IN COOL, DRY PLACE,
OUT OF DIRECT SUNSHINE.

PROTECT FROM EXCESSIVE HEAT.

BURN BAG IMMEDIATELY WHEN EMPTY STAY OUT OF SMOKE

DIRECTIONS

This product is not to be used for application by, g. and on
9/1/02 6:50 PM

NET WEIGHT 50 LBS.

Chevron Chemical Company
 6001 Airport Blvd., Emeryville, CA 94608
 San Francisco, CA 94119 R. L. Hood, CA 94304
 Product 3510 Made in U.S.A.
 Form 76284 EPA Est. 2
 EPA Reg. No. 2-277-91-3A



READ THE LABEL

CAUTION: Harmful if swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after use.

Note to Physicians: Emergency Information: Call 415-233-3636.

Use only in a transport near feed or food. May corrode metal equipment, clean equipment after use. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale or consumption.

[illegible]

Insecticides slightly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

TOMATOES. Early Blight, Late blight, Leaf Spot, Fruit Flies (Drosophila), Aphids, leafminers, Tomato Pinworm. 25 lbs per acre. Begin application of insecticides Repeat at 7 day intervals until 1 day of harvest.

COMPATIBILITY: Do not use the following oil sprays:

CONDITIONS OF SALE: I warrant that this is a correct description of the lot for sale as directed herein. I authorize any agent or representative of FITNESS or of NEF or representation, express or implied.

2. Physical and informational control prevents tampering with the use of elements. Such tampering can do damage to plants, but this is a rapid, lack of consequences to other plants, even though the product is restored again and even though, least. Buyer and user attention and liability (except those above) resulting from handling material.

ACCEPTED

MAR 7 1965

234-1281

BEST DOCUMENT AVAILABLE

239-2291

THOCIDE® DIBROM

4 Dust

BEST DOCUMENT AVAILABLE



FUNGICIDE

Per Wt.
95%
0.5%
4.0%
88.0%
ethyl 1,1-dichloro-2,2-bis(4-chlorophenyl) ethane
2,2-dichloroethyl dimethyl phosphite
N: KEEP OUT OF REACH OF
days of manufacture only, as the pro

USE STRICTLY IN ACCORDANCE
AND DIRECTIONS, AND WITH APPLI-
CATION FEDERAL REGULATIONS.
ORIGINAL CONTAINER
COOL, DRY PLACE,
DIRECT SUNSHINE.
FROM EXCESSIVE HEAT.
WHEN EMPTY, STAY OUT OF SMOKE
insulated for application by ground or

EIGHT 50 LBS.

Ortho Company
Riverside, CA 92504
EPA Est. 209-CW-5
39-3291 AA

GRAPE: Leafhoppers, Bunch Rot-40 lbs. per acre. Make
2 applications before bloom and 1 application just after
bloom. Repeat period daily up to 3 applications before
bunches close. Do not use within 4 days of harvest.

MELONS (Cantaloupes, Honeydews, Watermelons), Cucum-
bers, Summer Squash, Winter Squash, Downy Mildew, An-
thrachnose, Cucumber Beetles, Aphids, Leafminers, Pickle-
worms, Armyworms, Looper, Leafhoppers-35 lbs. per acre.
Begin application at first signs of disease and repeat
at 7-10 day intervals.



READ THE LABEL

CAUTION: Harmful if swallowed. May cause
irritation of eyes, nose and throat. Avoid breathing dust.
Avoid contact with skin and eyes. Wash thoroughly after
using.
Note to Physicians: Emergency information call 411-233-
373.
Do not store or transport near food or feed. May corrode
metal dust equipment, clean equipment after use. Do not
apply under conditions involving possible drift to food,
forage, or other plantings that might be damaged or the
crops thereof rendered unfit for sale, use or consumption.
Do not use on food crops or feed crops. Do not use on
crops that are to be used for seed. Do not use on crops
that are to be used for seed. Do not use on crops that are
to be used for seed. Do not use on crops that are to be
used for seed. Do not use on crops that are to be used
for seed. Do not use on crops that are to be used for
seed. Do not use on crops that are to be used for seed.
This product is highly toxic to bees exposed to direct treat-
ment or residues on crops. Protective information may be
obtained from your Cooperative Agricultural Extension
Service.

BEST DOCUMENT AVAILABLE

Do not apply within 1 day of harvest on Melons (Canta-
loupes, Honeydews, Watermelons), Winter Squash. Do not
apply within 4 days of harvest on Cucumbers and Sum-
mer Squash.

STRAWBERRIES: Red Spider Mites, Spittle Bug, Leafhopper,
Aphids, Botrytis Rot-40 lbs. per acre. Make first appli-
cation when new growth starts in spring before fruit starts to
form. Repeat application weekly. Do not apply within 4
days of harvest.

TOMATOES: Early Blight, Late Blight, Anthracnose, Sep-
toria Leaf Spot, Fruit Flies (Drosophila sp.), Two-spotted
Mites, Aphids, Leafminers, Tomato Fruitworms, Hornworms
(Tent caterpillars). Begin applications at first signs of dis-
ease. Repeat at 7 to 10 day intervals. Do not
apply within 1 day of harvest.

COMPATIBILITY: Do not apply to produce on foliage closely
covered with spray.

CONDITIONS OF SALE: 1. Ortho Chemical Company
warrants that this material conforms to the
description on the label and is reasonably fit
for the uses intended. 2. Ortho Chemical Company makes no
warranty, expressed or implied, for any other
uses of this material. 3. Ortho Chemical Company, its
representatives, agents, dealers, and distributors
do not assume any liability for damage or injury
resulting from the use of this material. 4. The
material is sold as a fungicide, and its use is
limited to the control of fungi. 5. The material
is sold as a fungicide, and its use is limited to
the control of fungi. 6. The material is sold as
a fungicide, and its use is limited to the control
of fungi. 7. The material is sold as a fungicide,
and its use is limited to the control of fungi.
8. The material is sold as a fungicide, and its
use is limited to the control of fungi. 9. The
material is sold as a fungicide, and its use is
limited to the control of fungi. 10. The material
is sold as a fungicide, and its use is limited to
the control of fungi.

ACCEPTED

MAR 7 1980

Ortho Company
Riverside, CA 92504
EPA Est. 209-CW-5
39-3291 AA

ORTHOCIDE® DIBROM 10-4 Dust



ORTHO

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

(INSECTICIDE-FUNGICIDE)

Active Ingredients	By Wt.
*Captan	9.5%
*Related Derivatives	0.5%
*Naled	4.0%
Inert Ingredients	86.0%
*N-(1,1-dichloromethylthio)-4-cyclohexene-1,2-dicarboximide.	
**1,2-dibromo-2,2-dichloroethyl dimethyl phosphate	

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Use within _____ days of manufacture only, as this product decomposes.

Date of Manufacture _____

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL, DRY PLACE,
OUT OF DIRECT SUNSHINE.

PROTECT FROM EXCESSIVE HEAT.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

This product is formulated for application by ground or air equipment.

GRAPES: Leafhoppers, Bunch Rot—40 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat just before up to 3 applications before harvest. Do not use within 4 days of harvest.

MELONS: Cantaloupes, Honeydews, Watermelons, Cucumbers, Summer Squash, Winter Squash, Downy Mildew, Anthracnose, Cucumber Beetles, Aphids, Leafminers, Pickleworms, Armyworms, Loopers, Leafhoppers—35 lbs. per acre. Do not use within 4 days of harvest.



READ THE LABEL

CAUTION: Harmful if swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after using.

Note to Physicians: Emergency information—call 415-233-0300.

Do not store or transport near food or feed. May corrode metal dust equipment; clean equipment after use. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply within 1 day of harvest on Melons (Cantaloupes, Honeydews, Watermelons), Winter Squash. Do not apply within 4 days of harvest on Cucumbers and Summer Squash.

STRAWBERRIES: Red Spider Mites, Spittle Bug, Leafhopper, Aphids, Botrytis Rot—40 lbs. per acre. Make first application when new growth starts in spring before fruit starts to form. Repeat application weekly. Do not apply within 4 days of harvest.

TOMATOES: Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot, Fruit Flies (*Drosophila* sp.), Twospotted Mites, Aphids, Leafminers, Tomato Fruitworms, Hornworms—25 lbs. per acre. Begin applications at first sign of damage or insects. Repeat at 7 to 10 day intervals. Do not apply within 1 day of harvest.

COMPATIBILITY: Do not use this product on foliage closely following oil sprays.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit to use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Actual and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability, except those assumed by Chevron under 1 above, resulting from handling, storage, and use of this material.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Agricultural Chemicals Division
San Francisco, CA 94119 Richmond, CA 94804

Product 3918 Made in USA

Form 76/8 N

FPA Est. 209 CA-5

PA Reg. No. 239 0091 AA

ACCEPTED

MAR 7 1960

Handwritten notes and signature: 239-2281