

ORTHO^(R)CIDE-BOTRAN^(R) 10-5 Dust



ORTHO

ACCEPTED

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. 250 281

(FUNGICIDE)

Active Ingredients	By Wt.
Captan**	10%
2,6-Dichloro-4-nitroaniline	5%
Inert Ingredients	85%
**N-(trichloromethylthio)-4-cyclohexene-1,2-dicarboximide	
U. S. Patent No. 3,119,736	
BOTRAN [®] is a registered trademark of Upjohn Co.	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL, DRY PLACE.

PROTECT FROM EXCESSIVE HEAT.

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

This product is formulated for application by ground or air equipment.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119
 Richmond, California Fresno, California
 Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida
 Form 7850-F Product 5022 Made in U.S.A.
 EPA Reg. No. 239-2307-AA

APRICOTS: Fruit Decay (Brown Rot and Rhizopus Rot)—50 lbs. per acre. Apply 10 days and 1 day prior to harvest. Do not apply to Apricots within 1 day of harvest.

PEACHES, NECTARINES: Brown Rot, Peach Scab—50 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. **Fruit Decay (Brown Rot, Rhizopus Rot)**—50 lbs. per acre. Apply 18 days, 10 days and 1 day prior to harvest. Do not apply within 1 day of harvest to peaches or nectarines.

SWEET CHERRIES: Cherry Leaf Spot, Brown Rot—50 to 60 lbs. per acre. Apply at popcorn, blossom and petal fall stages, continuing at 10 to 14-day intervals. **Fruit Decay (Brown Rot, Rhizopus Rot)**—50 lbs. per acre. Apply 10 days and 1 day prior to harvest. Do not apply to Cherries within 1 day of harvest.

ORNAMENTAL CROPS

ROSES, HYDRANGEA: Blight (Botrytis)—30 lbs. per acre. **Field:** Apply prior to shipment or storage. **Stored Areas:** Thoroughly dust storage area and all canes and dead foliage; repeat application during long storage periods.

CHRYSANTHEMUM, GERANIUM, ROSES: Blight (Botrytis)—30 lbs. per acre. Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10-day intervals.

GLADIOLUS: Dry Rot (Stromatinia)—Soil—250 lbs. per 14,520 linear feet of furrow. Dust into furrow at planting time, so treated soil surrounds corms. **Foliage**—30 lbs. per acre, dust thoroughly to insure coverage between spikes. Apply to bunched flower spikes after harvest and before shipment or storage.

COMPATIBILITY: Do not use this product on foliage closely following oil sprays.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. 2. Natural and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated herein and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability, except those assumed by Chevron under 1 above, resulting from handling, storage, and use of this material.

CAUTION **READ THE LABEL**

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin and eyes. In case of contact, remove contaminated clothing and wash skin with soap and water, for eyes, flush with water and call a physician. Do not store or transport near feed or food. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.