

VOLCK[®] Supreme-Ethion Combination Northwest



ORTHO

ACCEPTED

FEB 25 1977

Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act as amended for the pesticide registration act.
EPA Reg. No. 239-2312

(INSECTICIDE-MITICIDE)

Active Ingredients	By Wt.
Petroleum Oil	95%
*Ethion	2%
Inert Ingredients	3%
*O,O,O',O' Tetraethyl S,S' methylene bisphosphorodithioate	

Contains 0.16 pound Ethion per gallon
Petroleum Oil Classification—Unclassified
Petroleum Oil Min. Unsulfonated Residue—92%
U.S. Pat. No. 2,873,228

WARNING: KEEP OUT OF REACH OF CHILDREN.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DO NOT STORE AT TEMPERATURES BELOW 32° F.

WHEN DRUM IS EMPTY, TRIPLE RINSE (OR EQUIVALENT) AND OFFER FOR RECYCLING, RECONDITIONING OR DISPOSAL IN APPROVED LANDFILL OR BURY IN A SAFE PLACE.

VOLCK Supreme-Ethion Combination Northwest is for use in dormant and delayed dormant periods for control of scale insects, aphids and mites. THOROUGH COVERAGE IS NECESSARY. This product is for use in the Pacific Coast States for application during dormant and late dormant periods only. Do not apply after foliage or edible parts begin to form. Do not apply to foliage crops. Do not use in and later form.

NET CONTENTS 28 GALS. LIQUID

Chevron Chemical Company

Ortho Division San Francisco CA 94119 Richmond CA 94804

Product 5028 Made in U.S.A.

Form 7879 H

EPA Reg. No. 239-2312

DIRECTIONS

ALMOND, NECTARINE, PEACH, PLUM, PRUNE San Jose Scale, Soft Scales, Overwintering Eggs of European Red Mite and Brown Almond Mite, Parlatoria (Olive) Scale. Use 1-2 gals. per 100 gals. water during the dormant period. Do not graze livestock in treated almond groves.

APPLES San Jose Scale, Soft Scales, Overwintering eggs of Rosy Apple Aphid and Green Apple Aphid, European Red Mite or Brown Almond Mite. Use 1-2 gals. per 100 gals. water in the late dormant period. Blister Mites—Use 2 gals. per 100 gals. water in the late dormant period. Do not use on Astrachan, Duchesne, Marquise, Pratt, Melba, Wealthy, Williams Red (Red Williams), Yellow Transparent or any variety maturing before Early McIntosh, as injury may result.

PEAR San Jose Scale, Soft Scales, Overwintering Eggs of European Red Mite and Brown Almond Mite. Use 1-2 gals. per 100 gals. water during dormant period. Pear Psylla—2 gals. per 100 gals. water in delayed dormant period. Blister Mites—Use 2 gals. per 100 gals. water shortly after harvest of fruit or late dormant period. Do not apply to trees lacking moisture.



DO NOT TAKE INTERNALLY
DO NOT INHALE
DO NOT GET ON SKIN

WARNING:

May be fatal if swallowed. Absorbed through skin. Handle concentrate in a well-ventilated area. Do not breathe vapor or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling.

Note to Physicians: Emergency information call 1-800-233-1717. Antidote: none. 2-PAM is a first aid aid that may be administered in conjunction with other first aid measures.

This product is toxic to fish and should not be applied when bees are actively visiting the area.

This product is toxic to fish and should not be applied when bees are actively visiting the area. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply to any surface not specified on the label. Read equipment and state of wastes by burying in a container. If S. away from water supply.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

PLUM, PRUNE, APRICOT, CHERRY Overwintering eggs of Mealy Plum Louse, Leaf Curl Plum Aphid, European Red Mite and Brown Almond Mite. Use 1-2 gals. in 100 gals. water in the late dormant period.

MIXING DIRECTIONS: With enough water in tank to cover agitator shaft and with agitator in motion, add desired amount of this material for full tank. Pump this mixture through the overflow and back into tank for about two minutes to insure proper emulsification. Fill tank with water, adding other desired materials as tank is filling.

NOTE: This product may be used in combination with ORTHO Copper Fungicide. It may also be used in combination with ORTHO Lime-Sulfur Solution if the total solids are greater than 4 lbs. per 100 gals. spray, add an additional 1 gal. of dormant oil per 100 gals. Exercise care to insure complete mixing. Do not use in combination with or closely following ORTHOCIDE 50 Wettable or other captan or PHALTAN containing products. Do not use if material does not emulsify. Avoid application of oil sprays just prior to or during hot or freezing weather, or during rainy weather or dry hot winds. When temperatures drop below 32° F. (usually night time lows), do not spray tree and vine crops until daytime temperatures have warmed to a point above 35° F. and are free from moisture due to melting frost or heavy dews. Avoid double spraying. Plants receiving oil sprays should be in healthy condition and proper moisture should be present.

Test this product for proper emulsification by mixing a small quantity of it with water in a small container before filling the entire spray tank. Correct emulsification is indicated when a uniform milk white spray mixture results while shaking. If the spray mixture is not milk white after shaking, do not use the remaining product.

CONDITIONS OF SALE: Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed herein. Chevron neither makes nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, expressed or implied, concerning this product.

This product is sold on a "no-objection" basis and Chevron's control product is not recommended for use in combination with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses intended. Buyer assumes all responsibility for use of this product. Buyer is advised that this product is not intended for use on plants or crops not listed on the label.