



ORTHO Poison Ivy Control

ACCEPTED
 MAY 18, 1971
 U.S. DEPARTMENT OF AGRICULTURE
 INSECTIONS
 FOR ECONOMIC POISON
 EC UNDER 10



ORTHO

Poison Ivy Control

\$1.98

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

ORTHO Poison Ivy Control contains 2,4-D and MCPP. This product is designed to give easy control of poison ivy and poison oak around homes and buildings, in areas such as fence rows and vacant lots and on tree trunks. A foam marker has been added to aid in the complete coverage of all leaves.

DIRECTIONS

1. Point arrow on button toward red mark on edge of metal rim at top of can to direct spray properly and provide complete emptying of container.
2. Shake can well before use and occasionally during use.
3. Point arrow on button towards poison ivy and poison oak and spray to cover each leaf thoroughly. Do not overspray. The white foam marker will indicate when leaves have been thoroughly covered.
4. Spray only when air is calm to avoid drift to desirable plants. Poison ivy and poison oak may be sprayed any time after leaves are fully developed in the spring.
5. When poison ivy is growing up poles, fence posts or trunks of trees, spray all poison ivy leaves around the base and up to a height of 3 to 4 feet. Avoid spraying the foliage of desirable trees or shrubs. If poison ivy is growing on privet or lilac, to avoid damage to these plants cut ivy vines at base and apply ORTHO Poison Ivy Control to cut.
6. Under ordinary summer weather conditions, poison ivy and poison oak leaves turn brown in 10 days to 2 weeks after spraying. The chemicals are translocated throughout the entire plant, killing roots and tops.

IMPORTANT: Dead poison ivy and poison oak can cause skin irritations. Avoid any contact with bare arms or face. Do not burn dead poison ivy stems or foliage. Smoke from burning poison ivy and poison oak has caused severe injury to skin or to lungs when inhaled.

CAUTION: Avoid contact with skin, eyes or clothing. Wash hands after use. Avoid spray drift as this product may injure desirable flowers, vegetables, shrubs and turf. Do not store near fertilizers, seeds, insecticides or fungicides. Do not use or store near heat or open flame.

WARNING: Contents under pressure. Do not puncture. Exposure to high temperature may cause bursting. Keep at room temperature—away from direct sunlight, radiators, stoves, hot water and other heat. Never throw container into fire or incinerator.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.


Manufactured for

Chevron Chemical Company

Ortho Division/San Francisco, California 94119

Richmond, Calif. Lafayette, Calif. Des Plaines, Ill. Perth Amboy, N. J. Atl 3a

Form #094 USDA Reg. No. 239.2335 Product 1010.17 Made in U.S.A.



KILLS POISON OAK
AND IVY
ROOTS AND TOPS

ACCEPTED

\$ 1.98

Chevron

ORTHO Poison Ivy Control

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS

Ortho Poison Ivy Control contains 2,4-D and MCP. This product is designed to kill poison ivy, poison oak, and poison sumac and to help them regrow in a temperate and subtropical climate. It kills the trunk and branches, but does not kill the complete coverage of all leaves.

DIRECTIONS

1. Point arrow on button toward red mark on edge of nozzle and try to aim to direct spray properly and provide complete emptying of tank.
2. Shake can well before use and occasionally during use.
3. Point arrow on button toward poison ivy and poison oak and spray to cover each leaf thoroughly. Do not overspray. The white foam marker will indicate when leaves have been thoroughly covered.
4. Spray only when air is calm to avoid drift to desirable plants. Poison ivy and poison oak may be sprayed any time after leaves are fully developed in the spring.
5. When poison ivy is growing up the stems or trunks of trees, spray all poison ivy leaves around the base and up to a height of 3 to 4 feet. Avoid spraying the foliage of desirable trees or shrubs. It may be necessary to wet or mist to avoid damage to the plants. At the base of trees, apply ORTHO Poison Ivy Control to the trunk and spray summer weather on the stem, branches and leaves. Repeat every 10 days for 2 weeks after spraying. In other areas, spray throughout the entire plant, including roots and top.

IMPORTANT

Ortho Poison Ivy Control kills poison ivy, poison oak, and poison sumac. It does not kill the trunk and branches, but does not kill the complete coverage of all leaves.

CAUTION

Keep out of reach of children. Avoid spraying on children's clothing. Avoid spraying on children's hands, face, or neck. Avoid spraying on children's hair.

WARNING

Do not spray on children's clothing. Avoid spraying on children's hands, face, or neck. Avoid spraying on children's hair.

NOTICE

Ortho Poison Ivy Control kills poison ivy, poison oak, and poison sumac. It does not kill the trunk and branches, but does not kill the complete coverage of all leaves.

ORTHO

Poison Ivy Control



Active Ingredients	By Wt
*Diethanolamine salt of 2,4-Dichlorophenoxyacetic acid	0.75%
**Diethanolamine salt of 2-(2-Methyl-4-chlorophenoxy) propionic acid	0.75%
Inert Ingredients	98.50%
*Equivalent to 0.51% of 2,4-Dichlorophenoxyacetic acid	
**Equivalent to 0.50% of 2-(2-Methyl-4-chlorophenoxy) propionic acid	

CAUTION: Keep out of reach of children. See back panel for additional cautions.

NET WEIGHT 1 POUND 3 OUNCES