

Rose & Floral Dust

PAGE 9

ORDINAL


Rose & Floral Dust

POLICE

CUT HERE TO OPEN EASY

Police

Common House Fly
Aphididae
Pond Skater
Common Flower Garden Flea





ACCEPTED

JUL 13 1971

Rose & Floral Dust

Controls Insects - Aphids, Japanese Beetles,
Powdery Mildew, Black Spot and
Certain Other Flower Garden Pests



Active Ingredients	By Wt.
*Carbaryl (1-naphthyl N-methylcarbamate)	3.0%
**Malathion	4.0%
***Folpet	5.0%
****1,1-bis (p-chlorophenyl)-2,2,2-trichloroethanol	1.5%
Inert Ingredients	86.5%
*SEVIN® - Union Carbide Corp. U.S. Pat. Nos. 2,903,478-3,009,855	
**O,O dimethyl dithiophosphate of diethyl mercaptosuccinate	
***N-(trichloromethylthio) phthalimide (PHALTAN®)	
****KELTHANE® - Rohm & Haas Co.	

CAUTION: Keep out of reach of children.
See side panel for additional cautions.

NET WEIGHT 4 POUNDS

Chevron



**Rose &
Floral
Dust**

CONTROLS
Insects and
Diseases
Aphids, Japanese
Beetles, Beetles,
Mildew, Black
Spot and Certain
Other Flower
Garden Pests



ONION

Rose & Floral Dust

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DIRECTIONS

Apply as a dust or as a spray. Apply by dusting or as a spray in early morning or late evening when the temperature is low and the air is still. Apply so all exposed parts are contacted, especially upper and lower leaf surfaces.

As a Spray: Mix 8 Tablespoons in a little water to make a thin paste, add to 1 gallon water and agitate during application. 1 pound makes up to 10 gallons diluted spray. Do not store diluted spray. Make new dilution for each use.

For Insect Control: Treat thoroughly when damage or insects first appear. Repeat as necessary.

For Disease Control: Start regular preventive program of treatments as soon as foliage appears. Under normal conditions, apply every 7 to 10 days during growing season. Treat at least weekly during periods of moderate infection and twice weekly during periods of rapid new growth and severe infection caused by continuous summer rain or high humidity.

NOTE: Do not use on certain ferns including Boston, Maidenhair and Pteris, as well as on some species of Dioscorea. Do not use on Boston Ivy. Do not use on Japanese Maple.

Plants	Insects	Diseases
Roses and Other Flowers	Aphids, Spider Mites, Japanese Beetles, Rose Chalcids, Rose Leafhoppers, Leaf Rollers, Rose Strips, Lacebugs, Boxwood Leaf Miner, Bag- worms and Scale Crawlers.	Powdery Mildew, <u>Leaf Spot</u> , <u>Black</u> <u>Spot</u>

CAUTION: Harmful if swallowed. Induce vomiting if swallowed. May be absorbed through skin! Avoid breathing dust or spray. Avoid contact with skin and eyes. Wash thoroughly after handling. Keep children and animals away from treated areas until these areas are dry. Avoid contamination of feed or foodstuffs. Do not use within 24 hours of an oil spray treatment. Dispose of container promptly when empty.

Keep out of any body of water. Fish may be killed. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE: Buyer assumes all responsibility for safety and use not in accordance with directions.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DIRECTIONS

Apply as a dust or as a spray. Apply by dusting or as a spray in early morning or late evening when the temperature is low and the air is still. Apply so all exposed parts are contacted, especially upper and lower leaf surfaces.

As a Spray: Mix 8 Tablespoonfuls in a little water to make a thin paste, add to 1 gallon water and agitate during application. 1 pound makes up to 10 gallons diluted spray. Do not store diluted spray. Make new dilution for each use.

For Insect Control: Treat thoroughly when damage or insects first appear. Repeat as necessary.

For Disease Control: Start regular preventive program of treatment as soon as foliage appears. Under normal conditions, apply every 7 to 10 days during growing season. Treat at least weekly during periods of moderate infection and twice weekly during periods of rapid new growth and severe infection caused by continuous summer rains or high humidity.

NOTE: Do not use on certain ferns including Boston, Maidenhair and Pteris, as well as on some species of Crassula. Do not use on Boston Ivy. Do not use on Japanese Maple.

Plants	Insects	Diseases
Roses and Other Flowers	Aphids, Spider Mites, Japanese Beetles, Rose Chafers, Rose Leafhoppers, Leaf Rollers, Rose Slugs, Lacebugs, Boxwood Leaf Miner, Bag- worms and Scale Crawlers.	Powdery Mildew, Leaf Spot, Black Spot

CAUTION: Harmful if swallowed. Induce vomiting if swallowed. May be absorbed through skin! Avoid breathing of dust or spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Keep children and animals away from treated areas until these areas are dry. Avoid contamination of feed or foodstuffs. Do not use within 2 weeks of an oil spray treatment. Dispose of container promptly when empty.

Keep out of any body of water. Fish may be killed. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

©1968 Bayer assumes all responsibility for safety and use not in accordance with directions.