

Chevron

ORTHO Fruit & Vegetable Insect Control

Directions: ORTHO Fruit & Vegetable Insect Control is made ready-to-use with water and may be used on all types of plants. Use 2 teasp. (containing 1/2 oz.) per gallon of water or 1 pint in 48 gallons of water. For best results, apply full coverage spray to all exposed parts of plants, especially upper and lower leaf surfaces. **Numbers in () indicate days required between last application and harvest.** Spray trees thoroughly to point of runoff. Stay out of spray mist. Start spraying after blossoming when pests first appear. See table for detailed spray information.

FRUITS & NUTS

Crop	Insects	When to apply	How many sprays
Apples (14) Pears (14) Cherries (11)	Leafing Moth	Make first spray when petals have fallen	3-5 additional sprays at 7-10 day intervals
	Aphids, Scale Crawlers, Fruit Tree Leaf Rollers, Mites	When pests first appear	Repeat at 7-10 day intervals as necessary
Peaches (20) Appl. (13) Plums (10) Nectarines (7)	Peach Twig Borer, Oriental Fruit Moth	Make first spray when all petals have fallen	2-3 additional sprays at 10-15 day intervals
	Aphids, Scale Crawlers, Mites	When pests first appear	Repeat at 7-10 day intervals as necessary
Strawberries (1) Blackberries (1) Raspberries (1) Loganberries (1) Raspberries (1)	Aphids, Leafhoppers, Mites, Leaf Rollers	When pests first appear	Repeat at 7-10 day intervals as necessary
Watermelons (3)	Aphids, Scale Crawlers, Fruit Tree Leaf Rollers	When pests first appear	Repeat at 7-10 day intervals as necessary
Pears (1)	Aphids, Mites, Pear Leaf Casebearer, Wax Moth, Pear Leaf Rollers	late Apr. to early May; when pests first appear	Repeat at 7-10 day intervals as necessary; but not after fruit begins to ripen
Walnuts	Aphids, Mites, Leafing Moth, Shell Grafting Moth, Walnut Leafhopper	When pests first appear	Repeat at 7-10 day intervals as necessary

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL WARNINGS AND DIRECTIONS.

KEEP PESTS OUT OF REACH OF CHILDREN. MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. DO NOT BREATHE SPRAY MIST. AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WASH THOROUGHLY AFTER HANDLING. **Note to Physicians: Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Keep children and pets away from treated area until it is completely dry. Keep out of reach of domestic animals. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not contaminate ornamental fishpools. Do not reuse container. Destroy when empty.**

FOR DISEASE CONTROL on Fruits and Vegetables: Apply ORTHO Fruit & Vegetable Insect Control to foliage and fruit.

*On apples, light russeting may occur on some varieties, such as Golden Delicious.

VEGETABLES

Crop	Insects	When to apply	How many sprays
Tomatoes (1) Potatoes (5)	Aphids, Flea Beetles, Colorado Potato Beetles, Leaf Miners, Spotted Cucumber Beetles, Fall Armyworms	When pests first appear	Repeat at 7-10 day intervals as necessary
Beans (7) Pole Snap Lima Peppers (5)	Aphids, Mexican Bean Beetles, Leaf Miners, Cucumber Beetles	When pests first appear	Repeat at 7-10 day intervals as necessary. Do not feed treated beans to livestock.
Cantaloupes (3) Muskmelons (3) Watermelon (3) Summer Squash (7)	Aphids, Leafhoppers, Mites, Leaf Miner	When pests first appear	Repeat at 7-10 day intervals as necessary

Note: Do not treat food crops grown in the greenhouse.

Warning: Keep out of reach of children. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. **Note to Physicians:** Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Keep children and pets away from treated area until it is completely dry. Keep out of reach of domestic animals. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not contaminate ornamental fishpools. Do not reuse container. Destroy when empty.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Notice: Buyer assumes all responsibility for safety, and is not in any way connected with directions.

©

Chevron Chemical Company

San Francisco, California 94104

For more information, contact your local Cooperative Agricultural Extension Service.

© 1975 Chevron Chemical Company. All rights reserved.


EPA Reg. No. 139-0351 AA

Chevron

ORTHO

Fruit & Vegetable Insect Control

Makes 96
Gallons
Diluted Spray



ORTHO Fruit & Vegetable
Insect Control

ORTHO

**Fruit &
Vegetable
Insect Control**



Active Ingredients	By Wt
*O,O diethyl O (2 isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	25%
Aromatic Petroleum Derivative Solvent	57%
Inert Ingredients	18%

*Diazinon - a registered trademark of CIBA-GEIGY Corp.
U.S. Patent No. 2,754,243 DO NOT USE IN THE HOME

WARNING: Keep out of reach of children.
See back panel for additional warnings.

NET CONTENTS 1 QUART LIQUID