

PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS & DOMESTIC ANIMALS** **DANGER**

Corrosive to eyes. Causes eye damage. Harmful if swallowed or absorbed through the skin. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin or clothing. Do not use on household pets or humans. Keep children and pets away from treated area until completely dry. Keep out of reach of domestic animals.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, wash immediately with fresh water for 15 minutes and see a doctor. For skin contact, wash thoroughly with soap and water. Remove and launder contaminated clothing. If swallowed, DO NOT make person vomit. Call a doctor immediately.

Note to Physicians: Emergency information - call (415) 233-3737. This product contains a light petroleum solvent and may present an aspiration problem. The organophosphorus pesticide also present in this product is a cholinesterase inhibitor. If signs and symptoms of cholinesterase inhibition develop, atropine is antidotal. 2-PAM may also be given in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate ornamental fish ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Application should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

DIRECTIONS: ORTHO Fruit & Vegetable Insect Control mixes readily with water and may be used in any type sprayer. Use 2 teaspoons (1 fl. oz.) per gallon water or 1 pint in 40 gallons water. For best results, apply full coverage spray to all exposed parts of plants, especially upper and lower leaf surfaces. Numbers in () indicate days required between last application and harvest. Spray trees thoroughly to point of runoff. Stay out of spray mist. Start spraying after blooming or when pests first appear. Consult tables for detailed spray information.

FOR DISEASE CONTROL: Use Fruits and Vegetables Apple DITHOCIDE Garden Fungicide according to label directions.

Chevron Chemical Company

Ortho Consumer Products Division
 San Francisco CA 94119 Richmond CA 94804
 Product 1900 50 Made in U.S.A.
 Form 0060 G
 EPA Reg. No. 239-2350-AA

EPA Reg. No. 239-CA-1-239-NJ-1-239-MO-1
 Superscript used is first letter of lot number



FRUITS & NUTS

Crop	Insects	When to apply	How many sprays
Apples (14) Pears (14) Cherries (10)	Codling Moth Aphids, Scale Crawlers, Fruit Tree Leaf Rollers, Mites	Make first spray when petals have fallen When pests first appear	3-5 additional sprays at 7-10 day intervals Repeat at 7-10 day intervals as necessary
Peaches (20) Apricots (10) Plums (10) Nectarines (10)	Peach Tree Borer, Oriental Fruit Moth Aphids, Scale Crawlers, Mites	Make first spray when all petals have fallen When pests first appear	2-3 additional sprays at 10-15 day intervals Repeat at 7-10 day intervals as necessary
Caneberries (2) Blackberries Boysenberries Loganberries Raspberries Strawberries (5)	Aphids, European Tarnips, Mites, Scale Crawlers	When pests first appear	Repeat at 7-10 day intervals as necessary
Citrus Fruits (21) (15 apply to Lemons, Limes, Tangerines, Grapefruit, Kumquats)	Aphids, Scale Crawlers, Leaf Rollers, Thrips	When pests first appear. Do not apply during bloom	Repeat at 7-10 day intervals as necessary
Pecans	Aphids, Mites, Pecan Nut Casebearers, Walnut Caterpillars	Late April or early May for Nut Casebearers. Other pests when they first appear	Repeat at 7-10 day intervals as necessary, but not after pecan husks crack or walnut husks open. Do not exceed 6000 per acre.
Walnuts	Aphids, Mites, Codling Moth, Scale Crawlers, Walnut Caterpillars	When pests first appear	Repeat at 7-10 day intervals as necessary

*On apples slight russeting may occur on some varieties such as Golden Delicious.

VEGETABLES

Crop	Insects	When to apply	How many sprays
Tomatoes (1) Potatoes (25)	Aphids, Flea Beetles, Colorado Potato Beetles, Leaf Miners, Spotted Cucumber Beetles, Fall Armyworms	When pests first appear	Repeat at 10-day intervals or as necessary
Beans (17) (Pole Snap Lima) Peppers (5)	Aphids, Mexican Bean Beetles, Leaf Miners, Cucumber Beetles	When pests first appear	Repeat at 10-day intervals as necessary. Do not feed bean leaves.
Cantaloupes (3) Watermelon (3) Summer Squash (12)	Aphids, Leafhoppers, Mites, Leaf Miners	When pests first appear	Repeat at 10-day intervals as necessary. Do not feed bean leaves.

Note: Do not treat food crops grown in the greenhouse.

STORAGE AND DISPOSAL

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Do not reuse empty container. Wrap container and put in trash collection.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.



OR **Fruit & Vegetable Insect Control**

Controls Most Fruit



ACCEPTED
 NOV 21 1980
 Under the Federal Insecticide, Fungicide, or Rodenticide Act, this pesticide is registered under EPA Reg. No. 239-2350-AA

Keep out of reach
DANGER
 See back panel for
NET CONTENTS

STATEMENTS DOMESTIC ANIMALS

...wallowed or absorbed through the skin. Do not...
 ...washed. Avoid contact with skin or clothing...
 ...and pets away from treated area until...

TREATMENT

...fresh water for 15 minutes and see a doctor...
 ...water. Remove and launder contaminated...
 ...Call a doctor immediately...
 ...1 (415) 233-3737...
 ...and may present an aspiration problem. The...
 ...product is a cholinesterase inhibitor. If signs...
 ...of atropine is anticholinergic. 2 PAM may also be...

...ing in treated areas may be killed. Keep out...
 ...of fish ponds. Do not apply when weather...
 ...contaminated water by cleaning of equipment...

...treatment or to residues remaining on...
 ...during the crop cover crop or weeds bloom...
 ...need to provide the maximum possible interval...
 ...vary.

HAZARDS

...in manner inconsistent with its labeling...
 ...INMANCE WITH LABEL PRECAUTIONARY...

FRUITS & NUTS

Crop	Insects	When to apply	How many sprays
Apples (14) Pears (14) Cherries (10)	Codling Moth	Make first spray when petals have fallen	3-5 additional sprays at 7-10 day intervals
	Aphids, Scale Crawlers, Fruit Tree Leaf Rollers, Mites	When pests first appear	Repeat at 7-10 day intervals as necessary
Peaches (20) Apricots (10) Plums (10) Nectarines (10)	Peach Twig Borer, Oriental Fruit Moth	Make first spray when all petals have fallen	2-3 additional sprays at 10-15 day intervals
	Aphids, Scale Crawlers, Mites	When pests first appear	Repeat at 7-10 day intervals as necessary
Caneberries (7) (Blackberries, Boysenberries, Loganberries, Raspberries)	Aphids, Leafhoppers, Thrips, Mites, Scale Crawlers	When pests first appear	Repeat at 7-10 day intervals as necessary
Strawberries (5)			
Citrus Fruits (21) (Grapefruit, Lemons, Limes, Oranges, Tangerines, Japaneze Kumquats)	Aphids, Scale Crawlers, Leaf Rollers, Thrips	When insects first appear. Do not apply during bloom	Repeat at 7-10 day intervals as necessary
Pecans	Aphids, Mites, Pecan Nut Casebearers, Walnut Caterpillars	Late April or early May for Nut Casebearers. Other pests when they first appear	Repeat at 7-10 day intervals as necessary, but not after pecan husks crack or walnut husks open. Do not exceed 6 qts. per acre
Walnuts	Aphids, Mites, Codling Moth, Scale Crawlers, Walnut Caterpillars	When pests first appear	

*On apples slight russeting may occur on some varieties such as Golden Delicious.

VEGETABLES

Crop	Insects	When to apply	How many sprays
Tomatoes (1) Potatoes (35)	Aphids, Flea Beetles, Colorado Potato Beetles, Leaf Miners, Spotted Cucumber Beetles, Fall Armyworms	When pests first appear	Repeat at 7-10 day intervals as necessary
Beans (7) (Pole Snap Lima) Peppers (5)	Aphids, Mexican Bean Beetles, Leaf Miners, Cucumber Beetles	When pests first appear	Repeat at 7-10 day intervals as necessary. Do not treat bean foliage after flowering.
Cantaloupe (3) Muskmelon (3) Watermelon (3) Summer Squash (7)	Aphids, Leafhoppers, Mites, Leaf Miners	When pests first appear	Repeat at 7-10 day intervals as necessary.

Note: Do not treat food crops grown in the greenhouse.

STORAGE AND DISPOSAL

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Do not reuse empty container. Wrap container and put in trash collection.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.



ORTHO

Makes 96
Gallons
Diluted Spray

**Fruit & Vegetable
Insect Control**

Controls Most Fruit Tree and Vegetable Insects



CONTAINS
25% Diazinon

Active Ingredients: 25% Diazinon (25% Diethyl 1,2,4,5-tetramethyl-6-isopropyl-1,3,5-triazine-2,6-dithiolate)
 Inert Ingredients: 75%
 By Weight: 100%
 DO NOT USE IN THE HOME

ACCEPTED

NOV 21 1980

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this pesticide is registered under EPA Reg. No. 234-236C

Keep out of reach of children
DANGER
 See back panel for additional precautionary statements
NET CONTENTS 1 QT.

