

JUN 4 1981
 259 2356

FRUITS & NUTS

Crop	Insects	When to Apply	How many sprays
*Apples (14) Pears (14) Cherries (10)	Leafhopper Aphids, Scale Crawlers, Fruit Tree Leafhoppers, Mites	Make first spray when petals have fallen When pests first appear	3 to 5 additional sprays at 7 to 10 day intervals Repeat at 7 to 10 day intervals as necessary
Peaches (20) Apricots (10) Plums (10) Nectarines (10)	Peach Twig Borer, Oriental Fruit Moth Aphids, Scale Crawlers, Mites	Make first spray when all petals have fallen When pests first appear	2 to 3 additional sprays at 10 to 15 day intervals Repeat at 7 to 10 day intervals as necessary
Caneberries (7) (Blackberries, Boysenberries, Loganberries, Raspberries) Strawberries (5)	Aphids, Leafhoppers, Thrips, Mites, Scale Crawlers	When pests first appear	Repeat at 7 to 10 day intervals as necessary
Citrus Fruits (21) (Grapefruit, Lemons, Limes, Oranges, Tangelos, Tangerines, Kumquats)	Aphids, Scale Crawlers, Leafhoppers, Thrips	When insects first appear Do not apply during bloom	Repeat at 7 to 10 day intervals as necessary
Pecans	Aphids, Mites, Pecan Nut Casebearer, Walnut Caterpillars	Late April or early May for Nut Casebearer. Other pests when they first appear	Repeat at 7 to 10 day intervals as necessary, but not after pecan husks crack or walnut husks open. Do not exceed 6 qts per acre
Walnuts	Aphids, Mites, Codling Moth, Scale Crawlers, Walnut Caterpillars	When pests first appear	6 qts per acre

*On apples slight russeting may occur on some varieties such as Golden Delicious

VEGETABLES

Crop	Insects	When to Apply	How many sprays
Tomatoes (11) Potatoes (35)	Aphids, Flea Beetles, Colorado Potato Beetles, Leafminers, Spotted Cucumber Beetles, Fall Armyworms	When pests first appear	Repeat at 7 to 10 day intervals or as necessary
Beans (7) (Pole, Snap, Lima) Peppers (5)	Aphids, Mexican Bean Beetle, Leafminers, Cucumber Beetles	When pests first appear	Repeat at 7 to 10 day intervals as necessary. Do not feed treated bean forage. Peppers - Do not exceed 1 qt per acre
Cantaloupe (3) Muskmelon (3) Watermelon (3) Summer Squash (1)	Aphids, Leafhoppers, Mites, Leafminers	When pests first appear	

Note: Do not treat food crops grown in the greenhouse

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS & DOMESTIC ANIMALS
 DANGER**

Causes eye damage. Harmful if swallowed or absorbed through skin. Avoid contact with skin or clothing. Do not eat or drink from treated area until completely dry. Keep out of reach of children.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, wash immediately with fresh water for 15 minutes with soap and water. Remove and launder contaminated clothing immediately.

Note to Physicians: Emergency information - call (415) 233-3737

This product contains a light petroleum solvent and may present a fire hazard. It is a cholinesterase inhibitor. If signs and symptoms of poisoning occur, give 1 mg atropine immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on treated areas may be poisoned. Do not apply when weather conditions favor drift or when bees are active. Do not apply to areas where bees are active.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL.

DIRECTIONS

ORTHO Fruit & Vegetable Insect Control mixes readily with water. Use 1/2 oz per gallon water or 1 pint in 40 gallons water. For best results, spray especially upper and lower leaf surfaces. Numbers in parentheses indicate trees thoroughly to control. Stay out of spray mist. Start spray 10 minutes before sunrise.

FOR DISEASE CONTROL on Fruits and Vegetables - Apply ORTHO

STORAGE AND DISPOSAL

Keep pesticide in original container. Do not put concentrate or contents in food or feed containers. Do not reuse empty container. Wash container and contents thoroughly before disposal.

Do not use or store near heat or open flame.

