

ORTHO Liquid SEVIN*

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DIRECTIONS: ORTHO Liquid SEVIN mixes readily with water and may be used in all type sprayers. Shake container well before using. Use 1 tablespoonful (½ fl. oz.) per gallon water or 1 pint in 32 gallons water. Do not store diluted spray. For best results, apply full coverage spray to all exposed parts of plants, especially upper and lower leaf surfaces.

ROSES, ORNAMENTALS, FLOWERS, SHRUBS, SHADE TREES: Japanese Beetles, Gypsy Moth, Aphids, Bagworm, Elm Leaf Beetle, Certain Leafminers, Rose Slug, Periodical Cicada (17-year locust), Scale Insects, Tent Caterpillars, Blister Beetles, Leafrollers, Exposed Thrips—Start spraying when insects or their damage appear. Repeat weekly or as necessary. Do not use on Boston Ivy. Do not use on Virginia Creeper. Injury to tender foliage may occur if plants are wet when treated or in the presence of high humidity.

FRUIT TREES (APPLES, PEARS AND PEACHES): Apple Maggot, Plum Curculio, Apple Aphid, Leaf Rollers, Peach Tree Borer, Oriental Fruit Moth, Codling Moth—Make one application between bud break and bloom, a second application at petal fall, and then at 3 to 4 week intervals for 3 to 4 months as necessary. **Apple Maggot**—For best control, apply at 10 day intervals. **Rosy Apple Aphid**—Apply before leaves are curled. **NOTE:** To avoid foliage injury on McIntosh apples and possible fruit thinning on all varieties of apples, delay use until at least 30 days after bloom. Do not apply within 1 day of harvest.

VEGETABLES (TOMATOES, EGGPLANT, PEPPERS, CUCUMBERS, SUMMER SQUASH): Pickleworm, Fruitworms, Melon Worms, Hornworms, Fall Armyworms,

Corn Borer—Apply when insects first appear. Repeat at 7 to 10 day intervals. May be applied up to day of harvest.

LAWN INSECTS: Sod Webworms, Fall Armyworms, Earwigs, Grasshoppers, Fleas and Ants—Use 8 fl. oz. (1 cup) in 15 gallons water and cover 750 sq. ft. of established lawn. For best results, use an ORTHO Lawn Sprayer. Mow lawn first and apply after rain or watering. Do not water for at least two days after application. Repeat in 2 or 3 weeks or as necessary.

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Spray. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Skin Contact. Wash Hands and Face Before Eating. Keep children and animals off treated lawns until these areas are dry. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with pesticides.

Note to Physicians: Emergency Information—call (415) 233-3737. Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. **AVOID CONTAMINATION OF FOOD.** Dispose of container promptly when empty.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

Chevron Chemical Company

Ortho Division/San Francisco CA 94119 Richmond CA 94804

Product 5104 Made in U.S.A.

Form 8492-B

EPA Reg. No. 239-2356-AA



ORTHO Liquid SEVIN*

Controls Insects on Ornamentals, Flowers, Lawns, Fruit Trees and Vegetables. Kills Gypsy Moth, Japanese Beetles, Armyworms and Certain Other Insects

Active Ingredient	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	. . . 27%
Inert Ingredients 73%

U.S. Pat. Nos. 2,903,478 - 3,009,855

CAUTION: Keep out of reach of children. See back panel for additional cautions.

NET CONTENTS 1 QUART LIQUID

